SINGULAR BOUNDARY VALUE PROBLEMS FOR QUASI-DIFFERENTIAL EQUATIONS

PAUL W. ELOE

Department of Mathematics
University of Dayton
Dayton, Ohio 45469 USA

JOHNNY HENDERSON

Center for Dynamical Systems and Nonlinear Studies
Georgia Institute of Technology, Atlanta, Georgia 30332 and
Department of Discrete and Statistical Sciences
Auburn University, Auburn, Alabama 36849 USA

(Received April 13, 1993)

ABSTRACT. Solutions are obtained of boundary value problems for $L_n y + f(x, L_0 y, \dots, L_{n-2} y)$, satisfying $L_i y(0) = L_{n-1} y(1) = 0$, $0 \leq i \leq n-2$, where L_i denotes the i^{th} quasiderivative, and where $f(x, y_1, \dots, y_{n-1})$ has singularities at $y_i = 0$, $1 \leq i \leq n-1$.

KEYWORDS AND PHRASES. Quasi-derivative, singular boundary value problem, cone.

1992 AMS CLASSIFICATION CODE. 34B15.

1. INTRODUCTION.

We first define the quasiderivative operators, L_i , $0 \leq i \leq n$, inductively by,

$$\begin{aligned} L_0 u &= p_0 u, \\ L_i u &= p_i (L_{i-1} u)', \quad 1 \leq i \leq n, \end{aligned}$$

where $p_i(x) \in C^{(n-i)}([0, 1], (0, \infty))$, $0 \leq i \leq n$, and we assume $p_{n-1}(x) \equiv 1$ on $[0, 1]$. We will be concerned with solutions of boundary value problems for the quasi-differential equation,

$$L_n y + f(x, L_0 y, L_1 y, \dots, L_{n-2} y) = 0, \quad (1.1)$$

satisfying the boundary conditions,

$$L_i y(0) = 0, 0 \leq i \leq n-2, \text{ and } L_{n-1} y(1) = 0, \quad (1.2)$$

where $f(x, y_1, \dots, y_{n-1}) = 0$ has singularities at $y_i = 0$, $1 \leq i \leq n-1$. For notation, we let

$$\delta_i = \inf_{0 \leq x \leq 1} p_i(x), 0 \leq i \leq n,$$

and

$$\Delta_i = \sup_{0 \leq x \leq 1} p_i(x), 0 \leq i \leq n,$$

and we assume throughout that,

- (A) $f(x, y_1, \dots, y_{n-1}) : (0, 1) \times (0, \infty)^{n-1} \rightarrow (0, \infty)$ is continuous;
- (B) $f(x, y_1, \dots, y_{n-1})$ is decreasing in y_i , for each fixed $(x, y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n)$, $1 \leq i \leq n-1$;
- (C) $\int_0^1 f(x, y_1, \dots, y_{n-1}) dx < \infty$, for each fixed $(y_1, \dots, y_{n-1})$;
- (D) $\lim_{y_i \rightarrow +0} f(x, y_1, \dots, y_{n-1}) = +\infty$ uniformly on compact subsets of $(0, 1) \times (0, \infty)^{n-2}$, $1 \leq i \leq n-1$;
- (E) $\lim_{y_i \rightarrow +\infty} f(x, y_1, \dots, y_{n-1}) = 0$ uniformly on compact subsets of $(0, 1) \times (0, \infty)^{n-2}$, $1 \leq i \leq n-1$.

We observe that, if y is a solution of (1.1), then by (A), $L_n y < 0$, so that $L_{n-2} y$ is a concave function.

The singular boundary value problem (1.1), (1.2) generalizes in some sense the second order nonlinear singular problems considered by Bobisud [1], Bobisud and Lee [2], Do and Lee [3], Garner and Shrivaji [4], O'Regan [5], Tinco [6], and Wang [7]-[8]. Among those works, [1], [4], and [8] used singular boundary value problems to model diffusion problems arising in physiology and physics, while in [5], singular problems included as special cases the Thomas-Fermi and Emden-Fowler equations. Moreover, in [1], [2], [3], [5], [6], and [7] - [8], *a priori* bounds are established on solutions, and then Schauder degree or Granas topological transversality applications yield solutions of the singular boundary value problems. Others have also used these methods (in the case $p_i(x) \equiv 1$, $0 \leq i \leq n$, so that $L_n = \frac{d^n}{dt^n}$), for singular boundary value problems, with the paper by Eloë and Henderson [9] containing many references to those works.

Our motivation for the techniques used in obtaining solutions of (1.1), (1.2) are the works by [10] and [11], followed by the papers by Eloë and Henderson [12] and Henderson and Yin [13]. These arguments involve concavity properties, an iterative technique, and a fixed point theorem due to [11] for mappings that are decreasing with respect to a cone in a Banach space. In Section 2, we state some properties of a cone in a Banach space, followed by the fixed point theorem. In Section 3, we construct a suitable cone and define a sequence of modifications of f , so that none of these modifications have the singularities of f . For this sequence, we construct a sequence of operators, each of which satisfies the hypotheses of the fixed point theorem, hence obtaining a sequence of iterates in the cone. The sequence is shown to converge to a solution of (1.1), (1.2) in the cone.

2. SOME PRELIMINARIES.

The following definitions and properties of cones in a Banach space can be found in Amann's [14] treatise.

Let $\mathcal{B}$ be a Banach space, and K a closed, nonempty subset of $\mathcal{B}$. K is a *cone* provided (i) $\alpha u + \beta v \in K$, for all $u, v \in K$ and all $\alpha, \beta \geq 0$, and (ii) $u, -u \in K$ imply $u = 0$. Given a cone K , a *partial order*, $\leq$, is induced on $\mathcal{B}$ by $x \leq y$, for $x, y \in \mathcal{B}$, iff $y - x \in K$. (We may sometimes write $x \leq y$ (wrt K).) If $x, y \in \mathcal{B}$ with $x \leq y$, let $\langle x, y \rangle$ denote the *closed order interval between x and y* given by $\langle x, y \rangle = \{z \in \mathcal{B} \mid x \leq z \leq y\}$. A cone K is *normal* in $\mathcal{B}$ provided there exists a $\delta > 0$ such that $\|e_1 + e_2\| \geq \delta$, for all $e_1, e_2 \in K$ with $\|e_1\| = \|e_2\| = 1$.

We remark that, if K is a normal cone in $\mathcal{B}$, then closed order intervals are norm bounded.

We now state a fixed point theorem due to Gatica, Olikar, and Waltman [11] for operators that are decreasing with respect to a cone.

THEOREM 1. *Let $\mathcal{B}$ be a Banach space, K a normal cone in $\mathcal{B}$, $E \subseteq K$ such that, if $x, y \in E$ with $x \leq y$, then $\langle x, y \rangle \subseteq E$, and let $T : E \rightarrow K$ be a continuous mapping that is decreasing with respect to K , and which is compact on any closed order interval contained in E . Suppose there exists $x_0 \in E$ such that $T^2 x_0 = T(Tx_0)$ is defined, and both Tx_0 and $T^2 x_0$ are order comparable to x_0 . If either, (i) $Tx_0 \leq x_0$ and $T^2 x_0 \leq x_0$ or (ii) $x_0 \leq Tx_0$ and $x_0 \leq T^2 x_0$, then T has fixed point in E .*

3. SOLUTIONS OF (1.1), (1.2).

In this section, we will apply Theorem 1 to a sequence of operators that are decreasing with respect to an appropriate cone. We then obtain a sequence of iterates from these fixed points which converges to a solution of (1.1), (1.2). Concavity of the $(n-2)^{\text{nd}}$ quasiderivative of a solution plays a role in this construction.

Let the Banach space $\mathcal{B} = C^{(n-2)}[0, 1]$ with norm

$$\|y\| = \max\{|L_0 y|_\infty, \dots, |L_{n-2} y|_\infty\},$$

where $|\cdot|_\infty$ denotes the supremum norm, and let

$$K = \{y \in \mathcal{B} \mid L_i y(x) \geq 0 \text{ on } [0, 1], 0 \leq i \leq n-2\}.$$

K is a normal cone in $\mathcal{B}$. We also note that, if $u, v \in \mathcal{B}$ and $u \leq v$ (wrt K), then

$$L_i u(x) \leq L_i v(x) \text{ on } [0, 1], 0 \leq i \leq n-2.$$

In addition, we will have need of the sign of the Green's function, $G(x, s)$, for the problem,

$$-L_n y = 0, L_i y(0) = 0, 0 \leq i \leq n-2, \text{ and } L_{n-1} y(1) = 0. \quad (3.1)$$

Eloe [15] has proved

$$(L_i)_x G(x, s) > 0 \text{ on } (0, 1) \times (0, 1), 0 \leq i \leq n-2. \quad (3.2)$$

By a solution, y , of (1.1), (1.2), we mean $y \in C^{(n)}(0, 1) \cap C^{(n-1)}[0, 1]$, y satisfies (1.1) on $(0, 1)$, y satisfies (1.2), and $L_i y(x) > 0$ on $(0, 1)$, $0 \leq i \leq n-2$. For such, we seek a fixed point of the integral operator,

$$T\varphi(x) = \int_0^1 G(x, s) f(s, L_0 \varphi(s), \dots, L_{n-2} \varphi(s)) ds.$$

But because of the singularities in f given by (D), we cannot define T on all of the cone K . We next let $g_i : [0, 1] \rightarrow [0, \infty)$ be the solution of

$$L_n u = 0, \quad (3.3)$$

$$\begin{aligned} L_i u(0) &= 0, \quad 0 \leq i \leq n-2, \\ L_{n-1} u(0) &= 1, \end{aligned} \tag{3.4}$$

and, for each $\theta > 0$, define

$$g_\theta(x) = \theta g_1(x).$$

We note that, for $\theta > 0$, and $0 \leq i \leq n-2$,

$$L_i g_\theta(x) = \theta \int_0^x \int_0^{\tau_{n-i-1}} \cdots \int_0^{\tau_i} \frac{1}{p_{i+1}(\tau_{n-i-1}) \cdots p_{n-2}(\tau_2) p_{n-1}(\tau_1)} d\tau_1 d\tau_2 \cdots d\tau_{n-i-1},$$

and

$$L_{n-1} g_\theta(x) \equiv \theta \text{ on } [0, 1],$$

so that $g_\theta \in K$, and in fact $L_i g_\theta(x) > 0$ on $(0, 1]$, $0 \leq i \leq n-2$.

Assume for the remainder of the paper,

$$(F) \text{ For each } \theta > 0, \int_0^1 f(x, L_0 g_\theta(x), L_1 g_\theta(x), \dots, L_{n-2} g_\theta(x)) dx < \infty.$$

Finally, we define $D \subseteq K$ by

$$D = \{\varphi \in B \mid \text{there exists } \theta(\varphi) > 0 \text{ such that } g_\theta \leq \varphi (\text{wrt } K)\},$$

and define $T : D \rightarrow K$ by

$$T\varphi(x) = \int_0^1 G(x, s) f(s, L_0 \varphi(s), L_1 \varphi(s), \dots, L_{n-2} \varphi(s)) ds, \quad 0 \leq x \leq 1, \varphi \in D.$$

It follows from (A) - (F) and properties of $G(x, s)$ that, if $\varphi \in D$, then $T\varphi \in C^{(n)}(0, 1) \cap C^{(n-1)}[0, 1]$, $T\varphi$ satisfies (1.2), $L_i(T\varphi)(x) > 0$ and increasing on $(0, 1]$, $0 \leq i \leq n-2$, $L_{n-2}(T\varphi)$ is concave on $[0, 1]$, and that $T\varphi \in D$. On the other hand, if $\varphi \in C^{(n)}(0, 1) \cap C^{(n-1)}[0, 1]$ is a solution of (1.1), (1.2), with $L_i \varphi(x) > 0$ on $(0, 1]$, $0 \leq i \leq n-2$, it again follows from the concavity of $L_{n-2} \varphi$ that $\varphi \in D$. Consequently, $\varphi \in D$ is a solution of (1.1), (1.2) iff $T\varphi = \varphi$.

Our first result of this section gives *a priori* bounds on $L_{n-2} \varphi$, for all solutions φ of (1.1), (1.2), that belong to D .

THEOREM 2. Assume that (A)-(F) are satisfied. Then, there exists an $R > 0$ such that $|L_{n-2} \varphi|_\infty \leq R$, for all solutions φ of (1.1), (1.2), that belong to D .

PROOF. Assume the conclusion to be false. Then, there is a sequence $\{\varphi_\ell\} \subseteq D$ of solutions of (1.1), (1.2), such that $\lim_{\ell \rightarrow \infty} |L_{n-2} \varphi_\ell|_\infty = \infty$. We may assume that, for each $\ell \geq 1$,

$$|L_{n-2} \varphi_\ell|_\infty \leq |L_{n-2} \varphi_{\ell+1}|_\infty. \tag{3.5}$$

From the equation (1.1), $L_{n-1} \varphi_\ell(x) > 0$ and decreasing on $[0, 1]$, and from (1.2), $L_i \varphi_\ell(x) > 0$ and increasing on $(0, 1]$, $0 \leq i \leq n-2$. It follows that, for each ℓ ,

$$0 \leq L_{n-2} \varphi_\ell(1) = |L_{n-2} \varphi_\ell|_\infty \leq |L_{n-2} \varphi_{\ell+1}|_\infty = L_{n-2} \varphi_{\ell+1}(1). \tag{3.6}$$

In addition, the concavity and positivity of $L_{n-2}\varphi_\ell$ imply that

$$(L_{n-2}\varphi_\ell(1)) \int_0^x \frac{1}{p_{n-1}(s)} ds = L_{n-2}\varphi_\ell(1) \cdot x \leq L_{n-2}\varphi_\ell(x), 0 \leq x \leq 1.$$

So, from the monotonicity in (3.5) and (3.6), if we set $\theta = L_{n-2}\varphi_1(1)$, then

$$L_{n-2}g_\theta(x) \leq L_{n-2}\varphi_\ell(x) \text{ on } [0, 1], \ell \geq 1.$$

From the conditions satisfied by g_θ and φ_ℓ at $x = 0$, upon multiplying successively by $(p_i(x))^{-1}$ and integrating, we obtain

$$g_\theta \leq \varphi_\ell (\text{wrt } K), \text{ for all } \ell \geq 1.$$

Now, set

$$0 < M = \sup_{[0,1] \times [0,1]} (L_{n-2})_x G(x, s).$$

Then (B) and (F) yield that, for $0 \leq x \leq 1$ and $\ell \geq 1$,

$$\begin{aligned} L_{n-2}\varphi_\ell(x) &= L_{n-2}(T\varphi_\ell)(x) \\ &= \int_0^1 (L_{n-2})_x G(x, s) f(s, L_0\varphi_\ell(s), \dots, L_{n-2}\varphi_\ell(s)) ds \\ &\leq M \int_0^1 f(s, L_0g_\theta(s), \dots, L_{n-2}g_\theta(s)) ds \\ &= N, \end{aligned}$$

for some $0 < N < \infty$. But $0 \leq x \leq 1$ and $\ell \geq 1$ were arbitrary. So

$$|L_{n-2}\varphi_\ell|_\infty \leq N, \text{ for all } \ell \geq 1,$$

which contradicts $\lim_{\ell \rightarrow \infty} |L_{n-2}\varphi_\ell|_\infty = \infty$. The proof is complete. $\square$

COROLLARY . Assume (A)-(F) are satisfied. Then there exists an $R > 0$ such that $0 \leq L_i\varphi(x) \leq \left(\frac{R}{\delta_{n-2} \cdots \delta_{i+1}}\right) \frac{x^{n-i-2}}{(n-i-2)!}$ on $[0, 1]$, for $0 \leq i \leq n-2$, and $\|\varphi\| \leq \sup_{0 \leq i \leq n-2} \left\{ \frac{R}{\delta_{n-2} \cdots \delta_{i+1}} \right\} \equiv \tilde{R}$, for all solutions φ of (1.1), (1.2) that belong to D . In particular, $\varphi \leq \frac{\tilde{R}}{(n-2)!} x^{n-2}$ (wrt K), for all solutions $\varphi \in D$ of (1.1), (1.2).

Next, for each $\ell \geq 1$, let $\psi_\ell : [0, 1] \rightarrow [0, \infty)$ be defined by

$$\psi_\ell(x) = \int_0^1 G(x, s) f(s, \ell, \dots, \ell) ds.$$

With assumptions (A) - (E), we have

$$0 < L_i\psi_{\ell+1}(x) \leq L_i\psi_\ell(x) \text{ on } (0, 1), 0 \leq i \leq n-2.$$

Furthermore,

$$\lim_{\ell \rightarrow \infty} L_i\psi_\ell(x) = 0 \text{ uniformly on } [0, 1], 0 \leq i \leq n-2.$$

Now, define a sequence of functions, $f_\ell(x, y_1, \dots, y_{n-1}) : (0, 1) \times [0, \infty)^{n-1} \rightarrow (0, \infty)$ by

$$f_\ell(x, y_1, \dots, y_{n-1}) = f(x, \max\{y_1, L_0\psi_\ell(x)\}, \dots, \max\{y_{n-1}, L_{n-2}\psi_\ell(x)\}).$$

For $\ell \geq 1$, f_ℓ is continuous and satisfies (B). Also, from (B), we have, for each $\ell \geq 1$,

$$f_\ell(x, y_1, \dots, y_{n-1}) \leq f(x, y_1, \dots, y_{n-1}) \text{ on } (0, 1) \times (0, \infty)^{n-1},$$

and

$$f_\ell(x, y_1, \dots, y_{n-1}) \leq f(x, L_0\psi_\ell(x), \dots, L_{n-2}\psi_\ell(x)) \text{ on } (0, 1) \times (0, \infty)^{n-1}.$$

THEOREM 3. *Assume that conditions (A)-(F) are satisfied. Then the boundary value problem (1.1), (1.2) has a solution, y , such that $L_i y(x) > 0$ on $(0, 1)$, $0 \leq i \leq n-2$.*

PROOF. We begin by defining a sequence of compact mappings $T_\ell : K \rightarrow K$ by

$$T_\ell \varphi(x) = \int_0^1 G(x, s) f_\ell(s, L_0 \varphi(s), \dots, L_{n-2} \varphi(s)) ds, 0 \leq x \leq 1, \varphi \in K.$$

For $\ell \geq 1$ and $\varphi \in K$, $L_{n-2}(T_\ell \varphi) > 0$ is concave on $(0, 1)$, $T_\ell \varphi$ satisfies the boundary conditions (1.2), and from $(L_i)_x G(x, s) > 0$, $0 \leq i \leq n-2$, we have $L_i(T_\ell \varphi) > 0$ and increasing on $(0, 1)$, $0 \leq i \leq n-2$.

Since each f_ℓ satisfies (B), it follows that T_ℓ is decreasing with respect to the cone K , for each $\ell \geq 1$. Also, $0 \leq T_\ell(0)$ (wrt K) and $0 \leq T_\ell^2(0)$ (wrt K), and so by Theorem 1, for each ℓ , there exists a $\varphi_\ell \in K$ such that $T_\ell \varphi_\ell = \varphi_\ell$. From our observations above,

$$|L_{n-2} \varphi_\ell|_\infty = L_{n-2} \varphi_\ell(1).$$

By essentially the same arguments as in Theorem 2, it follows that there is an $R > 0$ such that, for each $\ell \geq 1$,

$$|L_{n-2} \varphi_\ell|_\infty \leq R \text{ and } \|\varphi_\ell\| \leq \tilde{R},$$

where $\tilde{R}$ is given in the Corollary.

Our next claim is that there exists $k > 0$ such that $k \leq |L_{n-2} \varphi_\ell|_\infty$, all $\ell \geq 1$. Assume the claim is false. Then passing to a subsequence and relabeling, we have without loss of generality that $\lim_{\ell \rightarrow \infty} |L_{n-2} \varphi_\ell|_\infty = 0$, which implies, along with the boundary conditions (1.2),

$$\lim_{\ell \rightarrow \infty} L_i \varphi_\ell(x) = 0 \text{ uniformly on } [0, 1], 0 \leq i \leq n-2. \quad (3.7)$$

Let $0 < \delta < \frac{1}{2}$ be fixed and let

$$0 < m = \inf_{[\delta, 1-\delta] \times [\delta, 1-\delta]} L_{n-2} G(x, s).$$

By (D), there exists $\eta > 0$ such that, if $\delta \leq x \leq 1 - \delta$ and $0 < y_i < \eta$, for $1 \leq i \leq n-1$, then

$$f(x, y_1, \dots, y_{n-1}) > \frac{2}{m}.$$

From (3.7), there exists $\ell_0 \geq 1$ such that, for $\ell \geq \ell_0$,

$$0 < |L_i \varphi_\ell(x)| < \eta/2, 0 < x < 1, 0 \leq i \leq n-2.$$

Also, for some $\ell_1 \geq \ell_0$ it follows that, if $\ell \geq \ell_1$,

$$0 < |L_i \psi_\ell(x)| < \eta/2, 0 < x < 1, 0 \leq i \leq n-2.$$

So, for $\ell \geq \ell_1$ and $\delta \leq x \leq 1 - \delta$,

$$\begin{aligned} L_{n-2}\varphi_\ell(x) &= \int_0^1 L_{n-2}G(x,s)f_\ell(s, L_0\varphi_\ell(s), \dots, L_{n-2}\varphi_\ell(s))ds \\ &\geq \int_\delta^{1-\delta} G(x,s)f_\ell(s, L_0\varphi_\ell(s), \dots, L_{n-2}\varphi_\ell(s))ds \\ &\geq m \int_\delta^{1-\delta} f(s, \eta/2, \dots, \eta/2)ds \\ &\geq 1. \end{aligned}$$

This is a contradiction to (3.6). Thus, there is a $k > 0$ such that $k \leq |L_{n-2}\varphi_\ell|_\infty$, for all ℓ . With $\theta = \frac{k}{2}$, and applying (1.2), we obtain

$$L_i g_\theta(x) \leq L_i \varphi_\ell(x) \text{ on } [0, 1], \quad 0 \leq i \leq n-2, \quad \ell \geq 1.$$

We have

$$g_\theta \leq \varphi_\ell \leq \frac{\tilde{R}}{(n-2)!} x^{n-2} (\text{wrt } K), \quad \ell \geq 1,$$

or, in particular,

$$\{\varphi_\ell\} \subseteq \langle g_\theta, \frac{\tilde{R}}{(n-2)!} x^{n-2} \rangle \subseteq D.$$

When restricted to the closed order interval, $\langle g_\theta, \frac{\tilde{R}}{(n-2)!} x^{n-2} \rangle$, T is a compact mapping. So, there is a subsequence of $\{T\varphi_\ell\}$, which we relabel as the original sequence, which converges to some $\varphi^* \in K$; that is, $\|T\varphi_\ell - \varphi^*\| \rightarrow 0$, as $\ell \rightarrow \infty$.

To complete the proof, we show that $\|T\varphi_\ell - \varphi_\ell\| \rightarrow 0$, as $\ell \rightarrow \infty$. With $\theta = \frac{k}{2}$, let $\epsilon > 0$ be given, and let

$$P = \max_{0 \leq i \leq n-2} \left\{ \sup_{[0,1] \times [0,1]} L_i G(x,s) \right\}.$$

The integrability condition (F) and the absolute continuity of the integral imply there exists $0 < \delta < 1$ such that

$$\begin{aligned} 2P \left[\int_0^\delta f(s, L_0 g_\theta(s), \dots, L_{n-2} g_\theta(s)) ds + \right. \\ \left. \int_{1-\delta}^1 f(s, L_0 g_\theta(s), \dots, L_{n-2} g_\theta(s)) ds \right] < \epsilon. \end{aligned}$$

Also, there exists ℓ_0 such that, for $\ell \geq \ell_0$,

$$L_i \psi_\ell(x) \leq L_i g_\theta(x) \leq L_i \varphi_\ell(x) \text{ on } [\delta, 1 - \delta], \quad 0 \leq i \leq n-2.$$

Observe also that $f_\ell(s, L_0\varphi_\ell(s), \dots, L_{n-2}\varphi_\ell(s)) = f(s, L_0\varphi_\ell(s), \dots, L_{n-2}\varphi_\ell(s))$, for $\delta \leq s \leq 1 - \delta$ and $\ell \geq \ell_0$. Thus, for $0 \leq i \leq n-2$, $\ell \geq \ell_0$, and $0 \leq x \leq 1$,

$$\begin{aligned} |L_i(T\varphi_\ell)(x) - L_i\varphi_\ell(x)| &\leq 2P \left[\int_0^\delta f(s, L_0\varphi_\ell(s), \dots, L_{n-2}\varphi_\ell(s)) ds \right. \\ &\quad \left. + \int_{1-\delta}^1 f(s, L_0\varphi_\ell(s), \dots, L_{n-2}\varphi_\ell(s)) ds \right] \\ &\leq 2P \left[\int_0^\delta f(s, L_0 g_\theta(s), \dots, L_{n-2} g_\theta(s)) ds \right. \\ &\quad \left. + \int_{1-\delta}^1 f(s, L_0 g_\theta(s), \dots, L_{n-2} g_\theta(s)) ds \right] \\ &< \epsilon. \end{aligned}$$

Therefore,

$$\lim_{t \rightarrow \infty} \|T\varphi_t - \varphi_t\| = 0.$$

It follows, in turn, that $\lim_{t \rightarrow \infty} \|\varphi_t - \varphi^*\| = 0$, so that

$$\varphi^* \in \langle g_\theta, \frac{\tilde{R}}{(n-2)!} x^{n-2} \rangle \subseteq D,$$

and

$$\varphi^* = \lim_{t \rightarrow \infty} T\varphi_t = T(\lim_{t \rightarrow \infty} \varphi_t) = T\varphi^*.$$

and the proof is complete. $\square$

REFERENCES

1. BOBISUD, L. E. Existence of solutions of some nonlinear diffusion problems, *J. Math. Anal. Appl.* **168** (1992), 413-424.
2. BOBISUD, L. E. and LEE, Y. S. Existence for a class of nonlinear singular boundary value problems, *Appl. Anal.* **38** (1990), 45-67.
3. DO, T. S. and LEE, H. H. Existence of solutions of singular nonlinear two-point boundary value problems, *Kyungpook Math. J.* **31** (1991), 113-118.
4. GARNER, J. B. and SHIVAJI, R. Existence and uniqueness of solutions of a class of singular diffusion problems, *Appl. Anal.* **30** (1988), 101-114.
5. O'REGAN, D. Existence theorems for certain classes of singular boundary value problems, *J. Math. Anal. Appl.* **168** (1992), 523-539.
6. TINEO, A. Existence theorems for a singular two-point Dirichlet problem, *Nonlin. Anal.* **19** (1992), 323-334.
7. WANG, H. Y. Solvability of some boundary value problems for the singular equation $(p(t)r(t,y)y')'/q(t)f(t,y,pry')$, *Chinese Ann. Math. Ser. A* **12** (1991), 87-91 (Chinese).
8. WANG, H. Y. Solvability of some singular boundary value problems for the equation $(p(t)r(t,u)u')'/q(t)f(t,u,pru')$, *Nonlin. Anal.* **18** (1992), 649-655.
9. ELOE, P. W. and HENDERSON J. Existence of solutions of some singular higher order boundary value problems, *Zeit. Angew. Math. Mech.* **72** (1992).
10. GATICA, J. A., HERNANDEZ, G. E., and WALTMAN, P. Radially symmetric solutions of a class of singular elliptic problems, *Proc. Edinburgh Math. Soc.* **33** (1990), 169-180.
11. GATICA, J. A., OLIKER, V., and WALTMAN, P. Singular nonlinear boundary value problems for second-order ordinary differential equations, *J. Diff. Eqs.* **79** (1989), 62-78.
12. ELOE, P. W. and HENDERSON, J. Singular nonlinear boundary value problems for higher order ordinary differential equations, *Nonlin. Anal.* **17** (1991), 1-10.
13. HENDERSON, J. and YIN, K. C. Singular boundary value problems, *Bull. Inst. Math. Acad. Sinica* **19** (1991), 229-242.
14. AMANN, H. Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces, *SIAM Rev.* **18** (1976), 620-709.
15. ELOE, P. W. Sign properties of Green's functions for two classes of boundary value problems, *Canad. Math. Bull.* **30** (1987), 28-35.