TWO CRITERIA FOR UNIVALENCY

N. SAMARIS

Department of Mathematics
University of Patras
Patras 26110, Greece

(Received January 11, 1994 and in revised form February 28, 1995)

ABSTRACT. In the present paper we give two criteria for the functions $f(z) = z + \alpha_2 z^2 + \dots$ to be univalent in $|z| < 1$

KEY WORDS AND PHRASES. Univalent functions, univalence.

1990 MATHEMATICS SUBJECT CLASSIFICATION. Primary 30C45.

Let A denote the class of functions which are analytic in the unit disk $U = \{z : |z| < 1\}$ and $f'(0) - 1 = 0$. By B we denote the class of functions $f \in A$ which are univalent, convex and bounded in U . In the present paper we prove the following theorems.

THEOREM 1. Let $f \in A$, satisfy the condition

$$\left| \frac{q_0^2}{q'} \left[\frac{z^2 f'}{f^2} - \frac{z^2 q'}{q^2} \right] \right| \leq 1 \quad \text{in } U. \quad (1)$$

for some $q \in B$, where $q_0 = \sup\{|q(z)|; z \in U\}$. Then f is univalent in U .

THEOREM 2. Let $f \in A, q \in B$ satisfy the condition

$$\left| \frac{1}{q'} \left[\frac{z}{f(z)} - \frac{z}{q(z)} \right] \right| \leq \lambda \quad \text{in } U, \quad (2)$$

for some $q \in B$, where

$$\lambda = \inf\left\{ \left| \frac{q'(z)}{q_0^2 q(z)} \right|; z \in U \right\}.$$

Then f is univalent in U .

If we put $q(z) = z$ in the Th.1 we get the Ozaki and Nunokawa theorem [2].

If we put $q(z) = z$ in the Th.2 we get the Nunokawa, Obradovic and Owa theorem [1].

PROOF OF THEOREM 1. If

$$\phi = \frac{q_0^2}{q'} \left[\frac{f'}{f^2} - \frac{q'}{q^2} \right]$$

then ϕ is analytic in U and

$$-\frac{1}{f(z)} + \frac{1}{q(z)} = \frac{1}{q_0^2} \int_0^z q'(\omega) \phi(\omega) d\omega + c.$$

If we put $q(\omega) = \xi$ we get

$$-\frac{1}{f(z)} + \frac{1}{q(z)} = \frac{1}{q_0^2} \int_0^{q(z)} \phi(q^{-1}(\xi)) d\xi + c \quad (3)$$

From the condition $q \in B$ and the relation (3) we get

$$\frac{1}{f(z_1)} - \frac{1}{f(z_2)} + \frac{1}{q(z_2)} - \frac{1}{q(z_1)} = \frac{1}{q_0^2} \int_{q(z_1)}^{q(z_2)} \phi(q^{-1}(\xi)) d\xi \quad (4)$$

From Schwarz's Lemma and condition (1) we get $|\phi(z)| \leq 1$ in U . Now from the relation (4) we get

$$\left| \frac{1}{f(z_1)} - \frac{1}{f(z_2)} + \frac{1}{q(z_2)} - \frac{1}{q(z_1)} \right| \leq \frac{|q(z_1) - q(z_2)|}{q_0^2} \quad (5)$$

If $f(z_1) = f(z_2)$ then it is obvious that $q(z_1) = q(z_2)$ or $z_1 = z_2$.

PROOF OF THEOREM 2. If we put

$$P(z) = \frac{z^2 f'}{f^2} - \frac{z^2 q'}{q^2}$$

then we get

$$P'(z) = -z \left[\frac{z}{f(z)} - \frac{z}{q(z)} \right]'' \quad \text{and} \quad \frac{|P'(z)|}{|q'(z)|} \leq \lambda \quad \text{in } U.$$

From the relation

$$P(z) = \int_0^{q(z)} \frac{P'(q^{-1}(\xi))}{q'(q^{-1}(\xi))} d\xi$$

we get

$$|P(z)| \leq \lambda |q(z)| \quad \text{in } U.$$

Now, the condition (1) of Th.1 is obvious.

REFERENCES

1. NUNOKAWA, M., OBRADOVIC, M. and OWA, S. One criterion for univalence. Proc. Amer. Math. Soc. **106** (1989) 1035-1037.
2. OZAKI, S. and NUNOKAWA, M. The Scharzian derivative and univalent functions, Proc. Amer. Math. Soc. **33** (1972), 392-394."