NEW CHARACTERIZATIONS FOR HANKEL TRANSFORMABLE SPACES OF ZEMANIAN

J. J. BETANCOR

Departamento de Analisis Matematico
Facultad de Matematicas
Universidad de La Laguna
38271 La Laguna, Tenerife
Islas Canarias, SPAIN

(Received April 11, 1994 and in revised form February 25, 1995)

ABSTRACT. In this paper we obtain new characterizations of the Zemanian spaces H_μ and H'_μ

KEY WORDS AND PHRASES. Hankel transform, distribution, Zemanian spaces

AMS SUBJECT CLASSIFICATION CODE. 46F12

A. H. Zemanian [7, Ch. 5] introduced the space H_μ ($\mu \in \mathbb{R}$) of functions as follows: a complex valued smooth function $\phi(x)$, $x \in I = (0, \infty)$, is in H_μ if, and only if, the quantity

$$\gamma_{n,k}^\mu(\phi) = \sup_{x \in I} |x^n (x^{-1} D)^k (x^{-\mu-1/2} \phi(x))| < \infty$$

is finite, for every $n, k \in \mathbb{N}$. This space endowed with the topology generated by $\{\gamma_{n,k}^\mu\}_{n,k \in \mathbb{N}}$ is a Fréchet space. In the sequel we will refer to the above topology as the usual topology of H_μ . Zemanian introduced the space H'_μ to extend the Hankel integral transformation defined by

$$(h_\mu \phi)(x) = \int_0^\infty (xt)^{1/2} J_\mu(xt) \phi(t) dt,$$

where J_μ denotes the Bessel function of the first kind and order μ , to generalized functions. He proved that h_μ is an automorphism of H_μ provided that $\mu \geq -\frac{1}{2}$. The generalized Hankel transform $\mathcal{H}$ for $f \in H'_\mu$, the dual space of H_μ , is defined as the transposed of H_μ through

$$\langle h'_\mu f, \phi \rangle = \langle \mathcal{H} f, h_\mu \phi \rangle \quad \text{for } \phi \in H_\mu.$$

Thus if $\mu \geq -\frac{1}{2}$, $\mathcal{H}$ is an automorphism of H'_μ when this space is equipped with the weak* topology or with the strong topology.

In [2] J. J. Betancor and I. Marrero have studied the main topological properties of the spaces H_μ and H'_μ . Amongst other results, it is established (Theorem 3.3) that the space H_μ , $\mu \geq -\frac{1}{2}$, is constituted by all those complex valued smooth functions $\phi(x)$, $x \in I$, such that

$$\tau_{n,k}^\mu(\phi) = \sup_{x \in I} |x^n N_{\mu+k-1} \dots N_\mu \phi(x)| < \infty$$

for every $n, k \in \mathbb{N}$. Moreover, the system of seminorms $\{\tau_{n,k}^\mu\}_{n,k \in \mathbb{N}}$ generates of H_μ its usual topology. Moreover in [4] they gave new descriptions for the usual topology of H_μ through L_2 -norms.

A. H. Zemanian [7, p. 134] defined the space $\mathcal{O}$ formed by all those complex valued smooth functions $v(x)$, $x \in I$, satisfying that for every $k \in \mathbb{N}$ there exists $n_k \in \mathbb{N}$ such that $(1+x^2)^{n_k} (x^{-1} D)^k v(x)$ is a bounded function on I . He proved that $\mathcal{O}$ is a space of multiplier of H_μ . Recently J. J. Betancor and I. Marrero [2, Theorems 2.3 and 4.9] have characterized $\mathcal{O}$ as the space of multipliers of H_μ and H'_μ .

In this paper we characterize the smooth complex valued functions in H_μ , $\mu \geq -\frac{1}{2}$, as the ones satisfying

$$Z_n(\phi) = \sup_{x \in I} |x^n \phi(x)| < \infty \quad (1)$$

and

$$y_n^\mu(\phi) = \sup_{x \in I} |N_{\mu+n-1} \dots N_\mu \phi(x)| < \infty \quad (2)$$

for every $n \in \mathbb{N}$. Moreover we prove that the usual topology of H_μ can be defined by the family of seminorms $\{Z_n, y_n^\mu\}_{n \in \mathbb{N}}$ and a new characterization for the elements of H'_μ is obtained. In the sequel we will assume that $\mu \geq -\frac{1}{2}$.

PROPOSITION 1. A complex valued smooth function $\phi(x)$, $x \in I$, is in H_μ if, and only if, ϕ satisfies (1) and (2) for every $n \in \mathbb{N}$.

PROOF. It is clear that if $\phi \in H_\mu$ then ϕ satisfies (1) and (2) for every $n \in \mathbb{N}$.

Let now ϕ be a complex valued smooth function defined on I . To see that (1) and (2) ($n \in \mathbb{N}$) are sufficient conditions for ϕ belongs to H_μ we proceed by induction. Suppose, as induction hypothesis, that

$$\sup_{x \in I} |x^m N_{\mu+n-1} \dots N_\mu \phi(x)| < \infty, \quad m \in \mathbb{N} \quad \text{and} \quad n \in \mathbb{N}, \quad 0 \leq n < \ell$$

for certain $\ell \in \mathbb{N}$, $\ell \geq 1$.

By using partial integration we can obtain

$$\begin{aligned} \|x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2^2 &= \int_0^\infty |x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)|^2 dx \\ &= \int_0^\infty x^{2m} N_{\mu+\ell-1} \dots N_\mu(\phi(x)) N_{\mu+\ell-1} \dots N_\mu(\overline{\phi}(x)) dx \\ &= \int_0^\infty (Dx^{-1})^\ell (x^{2m+\mu+\ell+1/2} N_{\mu+\ell-1} \dots N_\mu(\phi(x))) x^{-\mu-1/2} \overline{\phi}(x) dx \end{aligned}$$

for every $m \in \mathbb{N}$, $\ell < 2m+2$, because

$$\left[(Dx^{-1})^i (x^{2m+\mu+\ell+1/2} N_{\mu+\ell-1} \dots N_\mu(\phi(x))) (x^{-1} D)^{\ell-i-1} (x^{-\mu-1/2} \overline{\phi}(x)) \right]_0^\infty = 0 \quad (3)$$

for each $i, m \in \mathbb{N}$, $0 \leq i < \ell < 2m+2$. In effect, if $m, i \in \mathbb{N}$, $0 \leq i < \ell < 2m+2$ then Leibniz's rule leads to

$$\begin{aligned} &(Dx^{-1})^i (x^{2m+\mu+\ell+1/2} N_{\mu+\ell-1} \dots N_\mu(\phi(x))) (x^{-1} D)^{\ell-i-1} (x^{-\mu-1/2} \overline{\phi}(x)) \\ &= \sum_{j=0}^i a_j x^{2m+2\ell+2\mu+1-2j} (x^{-1} D)^{\ell+i-j} (x^{-\mu-1/2} \phi(x)) (x^{-1} D)^{\ell-i-1} (x^{-\mu-1/2} \overline{\phi}(x)) \\ &= \sum_{j=0}^i a_j x^{2m+1-j} N_{\mu+\ell+i-j-1} \dots N_\mu(\phi(x)) N_{\mu+\ell-i-2} \dots N_\mu(\phi(x)) \end{aligned}$$

where a_j , $j \in \mathbb{N}$, $0 \leq j \leq i$, are suitable real numbers, and by virtue of induction hypothesis (3) follows.

Most straightforward manipulations allow us to write

$$(Dx^{-1})^\ell (x^{2m+\mu+\ell+1/2} N_{\mu+\ell-1} \dots N_\mu(\phi(x))) x^{-\mu-1/2} \overline{\phi}(x) = \sum_{j=0}^\ell a_j x^{2m-j} \overline{\phi}(x) N_{\mu+2\ell-j-1} \dots N_\mu \phi(x)$$

with $m \in \mathbb{N}$ and $a_j \in \mathbb{R}$, $j \in \mathbb{N}$, $0 \leq j \leq \ell$.

Hence we can establish

$$\begin{aligned} \|x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2^2 &\leq C_1 \sum_{j=0}^\ell \int_0^\infty |x^{2m-j} \overline{\phi}(x)| |N_{\mu+2\ell-j-1} \dots N_\mu \phi(x)| dx \\ &\leq C_2 \sum_{j=0}^\ell \sup_{x \in I} |(1+x^2) x^{2m-j} \phi(x)| \sup_{x \in I} |N_{\mu+2\ell-j-1} \dots N_\mu \phi(x)| < \infty, \end{aligned} \quad (4)$$

provided that $m \in \mathbb{N}$, $2m \geq \ell$. Here C_i , $i = 1, 2$, denotes suitable positive constants.

Assume now that $m \in \mathbb{N}$, $2m < \ell$. We have

$$\begin{aligned} \|x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2^2 &= \left(\int_0^1 + \int_1^\infty \right) |x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)|^2 dx \\ &\leq \int_0^1 |N_{\mu+\ell-1} \dots N_\mu \phi(x)|^2 dx + \int_0^\infty |x' N_{\mu+\ell-1} \dots N_\mu \phi(x)|^2 dx. \end{aligned}$$

Therefore, by invoking (4) and the induction hypothesis we infer that

$$\|x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2 < \infty, \quad \text{when} \quad m \in \mathbb{N}, \quad 2m \leq \ell.$$

Thus it is concluded that $\|x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2 < \infty$, $m \in \mathbb{N}$.

Also, for every $m \in \mathbb{N}$, $m \geq 1$, and $x \in I$,

$$\begin{aligned} (x^m N_{\mu+\ell-1} \dots N_\mu \phi(x))^2 &= \int_0^x D_t(t^m N_{\mu+\ell-1} \dots N_\mu \phi(t))^2 dt \\ &= \int_0^x 2t^m N_{\mu+\ell-1} \dots N_\mu(\phi(t)) \left((m + \mu + \frac{1}{2} + \ell)t^{m-1} N_{\mu+\ell-1} \dots N_\mu(\phi(t)) + t^m N_{\mu+\ell} \dots N_\mu(\phi(t)) \right) dt. \end{aligned}$$

Hence if $m \in \mathbb{N}$, $m \geq 1$, and $x \in I$ by using Holder's inequality we can find $C \geq 0$ such that

$$\begin{aligned} |x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)|^2 &\leq C \left(\|x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2 \|x^{m-1} N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2 \right. \\ &\quad \left. + \sup_{x \in I} |N_{\mu+\ell} \dots N_\mu \phi(x)| \left[\|x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2 + \|x^{m+1} N_{\mu+\ell-1} \dots N_\mu \phi(x)\|_2 \right] \right) \end{aligned}$$

and then $\sup_{x \in I} |x^m N_{\mu+\ell-1} \dots N_\mu \phi(x)| < \infty$, $m \in \mathbb{N}$.

Thus the proof is finished.

The last proposition allows us to define the usual topology of H_μ through a family of seminorms simpler than $\{\gamma_{m,k}^\mu\}_{m,k \in \mathbb{N}}$.

PROPOSITION 2. The usual topology of H_μ is defined by the system of seminorms $\{Z_n, y_n^\mu\}_{n \in \mathbb{N}}$.

PROOF. It is clear that the topology generated by $\{\gamma_{m,k}^\mu\}_{m,k \in \mathbb{N}}$ is finer than the one defined by $\{Z_n, y_n^\mu\}_{n \in \mathbb{N}}$ on H_μ . Moreover by proceeding in a way similar to A. H. Zemanian [7, Lemma 5.2-2] we can prove that H_μ endowed with the topology generated by $\{Z_n, y_n^\mu\}_{n \in \mathbb{N}}$ is a Fréchet space. Hence the desired result is an immediate consequence of the Open Mapping Theorem [6, Corollary 2.12].

We now prove a new characterization for the elements of H'_μ the dual space of H_μ . The procedure employed is analogous to the one used by the author [1] and by J. J. Betancor and I. Marrero [2].

PROPOSITION 3. Let f be a linear functional defined on H_μ . Then f is in H'_μ if, and only if, there exist $r \in \mathbb{N}$ and $f_k, g_k \in L_\infty(0, \infty)$ (the space of essentially bounded functions on $(0, \infty)$), $k \in \mathbb{N}$, $0 \leq k \leq r$, such that

$$f = \sum_{k=0}^r h'_\mu(x^k f_k + x^{-\mu+1/2} (x^{-1} D)^k x^{k+\mu-1/2} g_k). \quad (5)$$

PROOF. Let $f \in H'_\mu$. By virtue of a well-known result ([7, Theorem 1.8-1]) there exist $r \in \mathbb{N}$ and $C > 0$ such that

$$|\langle f, \phi \rangle| \leq C \max_{0 \leq k \leq r} \{Z_k(\phi), y_k^\mu(\phi)\}, \quad \phi \in H_\mu. \quad (6)$$

According to [7, Lemma 5.4-1(2), (3) and Theorem 5.4-1] and since $z^{1/2} J_\mu(z)$ is a bounded function on I for every $k \in \mathbb{N}$ one has

$$\sup_{x \in I} |x^k \phi(x)| = \sup_{x \in I} |x^k h_\mu(h_\mu \phi)(x)| \leq C \int_0^\infty |N_{\mu+k-1} \dots N_\mu(h_\mu \phi)(t)| dt \quad (7)$$

and

$$\sup_{x \in I} |N_{\mu+k-1} \dots N_\mu \phi(x)| = \sup_{x \in I} |N_{\mu+k-1} \dots N_\mu h_\mu(h_\mu \phi)(x)| \leq C \int_0^\infty |t^k (h_\mu \phi)(t)| dt \quad (8)$$

for a suitable $C > 0$.

The linear mapping

$$j: H_\mu \rightarrow JH_\mu \subset L_1(0, \infty)^{2r+2} \\ \phi \rightarrow (x^k h_\mu \phi, N_{\mu+k-1} \dots N_\mu h_\mu \phi)_{k=0}^r$$

is one to one because h_μ is an automorphism of H_μ ([7, Theorem 5.4-1]). Here $L_1(0, \infty)$ denotes the usual Lebesgue space of order 1.

On the other hand, the inequalities (6), (7) and (8) imply that the linear mapping

$$L: JH_\mu \subset L_1(0, \infty)^{2r+2} \rightarrow \mathbb{C} \\ (x^k h_\mu \phi, N_{\mu+k-1} \dots N_\mu h_\mu \phi)_{k=0}^r \rightarrow \langle f, \phi \rangle$$

is continuous when JH_μ is endowed with the topology induced by $L_1(0, \infty)^{2r+2}$. Hence, by invoking the Hahn-Banach Theorem L can be extended to $L_1(0, \infty)^{2r+2}$ as a member of $(L_1(0, \infty)^{2r+2})'$, the dual space of $L_1(0, \infty)^{2r+2}$. Since, as it is well known, $L_1(0, \infty)' = L_\infty(0, \infty)$ there exist $f_k, g_k \in L_\infty(0, \infty)$, $k \in \mathbb{N}$, $0 \leq k \leq r$, such that

$$\langle f, \phi \rangle = \sum_{k=0}^r (\langle f_k, x^k h_\mu \phi \rangle + \langle g_k, x^{k+\mu+1/2} (x^{-1} D)^k (x^{-\mu-1/2} \phi) \rangle), \quad \phi \in H_\mu.$$

Therefore

$$f = \sum_{k=0}^r h'_\mu (x^k f_k + (-1)^k x^{-\mu+1/2} (x^{-1} D)^k x^{k+\mu-1/2} g_k).$$

Thus the proof of necessity is finished.

Conversely, if f is a linear functional defined on H_μ by (5) for certain $r \in \mathbb{N}$ and $f_k, g_k \in L_\infty(0, \infty)$, $k \in \mathbb{N}$, $0 \leq k \leq r$, then

$$|\langle f, \phi \rangle| \leq C \sum_{k=0}^r \left(\|f_k\|_\infty \sup_{x \in I} |(1+x^2)x^k (h_\mu \phi)(x)| + \|g_k\|_\infty \sup_{x \in I} |(1+x^2)N_{\mu+k-1} \dots N_\mu (h_\mu \phi)(x)| \right)$$

for $\phi \in H_\mu$, where $\|\cdot\|_\infty$ denotes the usual norm in $L_\infty(0, \infty)$. Hence, according to [7, Theorem 5.4-1] and [2, Theorem 3.3], f is in H'_μ .

REFERENCES

- [1] BETANCOR, J.J., Characterization of Hankel transformable generalized functions, *Internat. J. Math. & Math. Sci.*, **14** (2) (1991), 269-274.
- [2] BETANCOR, J.J. and MARRERO, I., Some linear topological properties of the Zemanian space H_μ , *Bull. de la Soc. Roy. des Sci. de Liege*, **61** (3-4) (1992), 299-314.
- [3] BETANCOR, J.J. and MARRERO, I., Multipliers of Hankel transformable generalized functions, *Comment. Math. Univ. Carolinae*, **33** (3) (1992), 389-401.
- [4] BETANCOR, J.J. and MARRERO, I., A Hilbert space approach to Hankel transformable distributions, *Applicable Analysis*, **52** (1994), 103-124.
- [5] CHUNG, J., CHUNG, S-Y. and KIM, D., Une caractérisation de l'espace S de Schwartz, *C. R. Acad. Sci. Paris*, 316 Serie I (1993), 23-25.
- [6] RUDIN, W., *Functional Analysis* (Second Edition), McGraw-Hill, Inc., New York, 1991.
- [7] ZEMANIAN, A. H., *Generalized Integral Transformations*, Dover Publications, Inc., New York, 1987.