**STRICTLY BARRELLED DISKS IN INDUCTIVE LIMITS
OF QUASI-(LB)-SPACES**

CARLOS BOSCH

Department of Mathematics
I.T.A.M.

Río Hondo #1, Col. Tizapán San Angel
México 01000, D.F., Mexico

THOMAS E. GILSDORF

Department of Mathematics
University of North Dakota
Grand Forks, ND 58202-8376, USA

(Received June 9, 1995)

ABSTRACT. A strictly barrelled disk B in a Hausdorff locally convex space E is a disk such that the linear span of B with the topology of the Minkowski functional of B is a strictly barrelled space. Valdivia's closed graph theorems are used to show that closed strictly barrelled disk in a quasi-(LB)-space is bounded. It is shown that a locally strictly barrelled quasi-(LB)-space is locally complete. Also, we show that a regular inductive limit of quasi-(LB)-spaces is locally complete if and only if each closed bounded disk is a strictly barrelled disk in one of the constituents.

KEY WORDS AND PHRASES. Quasi-(LB)-space, strictly barrelled space, inductive limit.

1991 AMS SUBJECT CLASSIFICATION CODE. Primary 46A13, 46A08. Secondary 46A30.

1. INTRODUCTION.

Throughout this paper, we use the word *space* to denote a Hausdorff locally convex space. An absolutely convex set will be called a *disk*. If A is a disk in a space E , its linear span E_A may be endowed with the semi-normed topology

given by the Minkowski functional of A . When distinction is needed, we will denote this topology by ρ_A . When A is a bounded disk, it is easy to see that E_A is normed and that $\text{id}: E_A \rightarrow E$ is continuous. If E_A is a Banach space (resp. Baire space), we call A a Banach (resp. Baire) disk. If every bounded subset of E is contained in a bounded Banach (resp. Baire) disk, we say that E is locally complete (resp. locally Baire). Locally complete spaces are also called fast complete, and according to [1; 5.1.6, pg. 152], a space is locally complete if and only if every closed bounded disk is already a Banach disk.

DEFINITION 1.1: Following [2], a space E is strictly barrelled if given any ordered absolutely convex web $\mathcal{W}$ on E there exists a strand $(W(k)) = \{W(k) : k \in \mathbb{N}\}$ of $\mathcal{W}$ such that for each positive integer k , the closure $\overline{W(k)}$ is a zero neighborhood in E , where $W(k)$ denotes the k th member of a strand $(W(k))$.

DEFINITION 1.2: Let A be a disk. If E_A is a strictly barrelled space, we will say that A is a strictly barrelled disk. If every bounded set is contained in a strictly barrelled disk, we say that E is locally strictly barrelled.

REMARK 1.3: Using [1; chapt. 9] and [2; Prop. 6.17, pg. 160],
locally complete $\Rightarrow$ locally Baire $\Rightarrow$ locally strictly barrelled.

These implications cannot be reversed; the first by [1; 1.2.12 pg. 17], the second by [2; Prop. 17, pg. 160 & Note 4, pg. 162]. Valdivia defines quasi-(LB)-spaces in [2], and proves a webbed-space equivalence in [2; Th. 4.1, pg. 153]. We will use this equivalence as our definition below.

DEFINITION 1.4: A space with an ordered, absolutely convex strict web is called a quasi-(LB)-space.

2. QUASI-(LB)-SPACES AND STRICTLY BARRELLED DISKS.

The following generalizes [3; Th. 3, pg. 173] and [4; Th. 1, pg. 222].

THEOREM 2.1: Let B be a closed strictly barrelled disk in a quasi- (LB)-space. Then B is bounded.

PROOF: Let (E, τ) be the quasi- (LB)-space that contains B . Denote by η the topology induced on E_B by the following system of neighborhoods: $\{(n^{-1}B) \cap V : V \text{ is a } \tau\text{-closed zero neighborhood, } n \in \mathbb{N}\}$. Using the ordered strict web on (E, τ) and the

construction in [4; Th. 1, pg. 222], we have that (E_B, η) is a quasi-(LB)-space. The map $id: (E_B, \eta) \rightarrow (E_B, \rho_B)$ is continuous and (E_B, ρ_B) is strictly barrelled. Therefore, by [2; Th. 6.5(a), pg. 163], this map is open, implying that for any τ -zero neighborhood V , $id(B \cap V)$ is a neighborhood of zero in (E_B, ρ_B) . In particular, there exists $a > 0$ such that $aB \subset B \cap V \subset V$. We conclude that B is τ -bounded. ♦

The result that follows uses the closed graph theorem of Valdivia [2].

THEOREM 2.2: *Any locally strictly barrelled quasi-(LB)-space is locally complete.*

PROOF: Assume (E, ι) is such a space and suppose A is bounded in E . There is a bounded disk $B \supset A$ such that (E_B, ρ_B) is strictly barrelled. Because $id: (E_B, \rho_B) \rightarrow (E_B, \iota)$ is continuous, [2; Th. 7.6 pg. 164] shows that there is a Fréchet space F for which $E_B = id(E_B) \subset F$ and the following injections are continuous: $(E_B, \rho_B) \rightarrow F \rightarrow (E_B, \iota)$. Hence, there is a bounded Banach disk D in F , with $A \subset B \subset D$, and D is a bounded Banach disk in E as well. ♦

3. INDUCTIVE LIMITS.

In this section we consider sequences (E_n, ι_n) , $n \in \mathbb{N}$ of spaces with $E_1 \subset E_2 \subset \dots$, and for every positive integer n , E_n injects continuously into E_{n+1} . We put $E = \text{ind}_n E_n$ for the inductive limit. Recall that an inductive limit is called regular if for any of its bounded subsets, there is a constituent space such that the subset is contained in and bounded in that constituent.

THEOREM 3.1: *Let $E = \text{ind}_n E_n$ be an inductive limit of quasi-(LB)-spaces. Suppose B is a disk in (E_n, ι_n) . Then:*

- (a) *If there exists $m \geq n$ such that B is a closed strictly barrelled disk in (E_m, ι_m) , then B is a closed bounded strictly barrelled disk in both (E_n, ι_n) and (E_m, ι_m) . Moreover, B is contained in a bounded Banach disk in (E_n, ι_n) and (E_m, ι_m) .*
- (b) *If (a) holds for every bounded disk in E_n , then E_n is locally complete.*

(c) If E is regular and locally complete, then E_n is locally complete for every positive integer n .

PROOF: (a): If the assumptions are satisfied, then from the continuity of $\text{id}: (E_n, t_n) \rightarrow (E_m, t_m)$, B is t_n -closed. As a strictly barrelled, closed disk in (E_m, t_m) , B is t_n -bounded by Theorem 2.1. We use Theorem 2.2 in both (E_m, t_m) and (E_n, t_n) to conclude that B is contained in a bounded Banach disk in both spaces.

(b): Obvious consequence of (a).

(c): Let E be any fixed natural number and let $A \subset E_n$ be bounded. By the assumptions and topology on E , A is bounded in E , and contained in an E -closed, bounded Banach disk D , where D itself is contained in and bounded in some (E_m, t_m) ; clearly $m \geq n$. As $\text{id}: (E_m, t_m) \rightarrow E$ is continuous, D is t_m -closed and of course is a bounded Banach disk there. We apply part (a) to the disk $D \cap E_n$ and we are done. ♦

In [5] we have that if each (E_n, t_n) is webbed and locally complete, then is $E = \text{ind}_n E_n$ regular if and only if it is locally complete. One can ask what happens if the inductive limit is regular but the spaces (E_n, t_n) are not locally complete; see for example [6] and [7]. It is not difficult to prove a similar type of result using quasi-(LB)-spaces; the details follow. Compare also [4; Th. 3, pg 223] and [3; Th.5, pg. 174].

THEOREM 3.2: *Suppose each (E_n, t_n) is a quasi-(LB)-space and $E = \text{ind}_n E_n$ is regular. Then E is locally complete if and only if for each closed, bounded disk $B \subset E_n$, there is an $m \in \mathbb{N}$ such that B is a strictly barrelled disk in (E_m, t_m) .*

PROOF: If E is locally complete, the conclusion follows directly from 3.1 (c). Conversely, take a closed, bounded disk B in E . There is an $n \in \mathbb{N}$ such that $B \subset E_n$ and is t_n -bounded, and there is an $m \in \mathbb{N}$ with $B \subset E_m$ and B is a strictly barrelled disk. If $m > n$, we use 3.1 (a). On the other hand, if $n \geq m$, then 2.1 tells us that $n = m$ and (a) of 3.1 applies. In either case, E is locally complete.

♦

We want to construct a regular inductive limit of non-locally complete quasi-(LB)-spaces, but first we need:

LEMMA 3.3: *A finite product of locally convex spaces is locally complete if and only if each space is locally complete.*

PROOF: One may use bornologies, [8; 3.2(3), pg 43], to prove that any product of locally complete spaces is locally complete. Conversely, let $E = F \times G$, and assume that E is locally complete. Suppose, without loss of generality, that F is not locally complete. This means there is a disk B , closed and bounded in F , and B is not a Banach disk in F . Then $B' = B \times \{0\}$ is an E -closed and bounded disk that is not a Banach disk, a contradiction. Hence, F is locally complete. The proof for general finite products can be done by induction. ♦

EXAMPLE 3.4: Let E_0 be an non-regular (LB)-space. Then E_0 is a quasi-(LB)-space by [2; Prop 3.5, pg 152]. For each positive integer n , put $E_n = \oplus \{E_0 : i = 1, 2, \dots\} \cong \prod \{E_0 : i = 1, 2, \dots\}$. The lemma, the non-regularity of E_0 and [2; Prop 3.3, pg 151] imply that each E_n is a non-locally complete quasi-(LB)-space.

Set $E = \text{ind}_n E_n = \oplus \{E_0 : n \in \mathbb{N}\}$. As a direct sum, if $A \subset E$ is bounded, then there is a finite subset I of $\mathbb{N}$ such that A is bounded in $\oplus \{E_0 : i \in I\}$. If $n = \max\{i : i \in I\}$, then A is bounded in E_n , and E is therefore regular. Next, we use 3.2. Let $B \subset E_I = E_0$ be a closed, bounded disk that is not a Banach disk. Using the definition of the direct sum topology t of E and the fact that t induces on E_0 its own topology, we have that B is a closed bounded disk in E , also. The disk B cannot be a Banach disk in E , so E is not locally complete. From 3.2, we see that B is in fact a really bad disk; not only is it a non-Banach disk in E , it cannot be a strictly barrelled disk in any E_n . ♦

ACKNOWLEDGEMENT. Research for the second author was supported as part of a Solomon Lefshetz Fellowship at el Centro de Investigaciones y Estudios Avanzados, Mexico City, Mexico, 1992-1993.

REFERENCES

- [1] PEREZ-CARRERAS, P., BONET, J.; Barrelled Locally Convex Spaces, North Holland Math. Studies, 131, (1987).
- [2] VALDIVIA, M.; *Quasi-(LB)-spaces*, J. London Math. Soc., (2) 35, (1987), pp. 149-168.
- [3] QIU, J. H.; *Dieudonné-Schwartz theorem in inductive limits of metrizable spaces II*, Proc. AMS, 108, no. 1, Jan. (1990), pp. 171-175.
- [4] BOSCH, C., KUCERA, J.; *Bounded sets in inductive limits of $\mathcal{K}$ -spaces*, Czech J. Math., 43 (118), (1993), pp. 221-223.
- [5] BOSCH, C., KUCERA, J.; *Bounded sets in fast complete inductive limits*, Int. J. Math. & Math. Sci., 7, no. 3, (1984), pp. 615-617.
- [6] GILSDORF, T.; *Local Baire-like properties and inductive limits*, preprint.
- [7] GILSDORF, T.; *Regular inductive limits of $\mathcal{K}$ -spaces*, Collect. Math., 42, no. 1, 1991 (1992), pp. 45-49.
- [8] HOGBE-NLEND, H.; Bornologies and Functional Analysis, North Holland Math. Studies, 26, (1977).