FRACTIONAL DERIVATIVES OF HOLOMORPHIC FUNCTIONS ON BOUNDED SYMMETRIC DOMAINS OF $\mathbb{C}^n$

ZENGJIAN LOU

Department of Mathematics
Qufu Normal University
Qufu Shandong, 273165
P R CHINA

(Received March 8, 1994 and in revised form January 13, 1995)

ABSTRACT. Let $f \in H(B_n)$ $f^{[\beta]}$ denotes the β th fractional derivative of f If $f^{[\beta]} \in A^{p,q,\alpha}(B_n)$, we show that

- (I) If $\beta < \frac{\alpha+1}{p} + \frac{n}{q} = \delta$, then $f \in A^{s,t,\alpha}(B_n)$, and $\|f\|_{s,t,\alpha} \leq C \|f^{[\beta]}\|_{p,q,\alpha}$, $s = \frac{\delta p}{\delta - \beta}$, $t = \frac{\delta q}{\delta - \beta}$
- (II) If $\beta = \frac{\alpha+1}{p} + \frac{n}{q}$, then $f \in B(B_n)$ and $\|f\|_B \leq C \|f^{[\beta]}\|_{p,q,\alpha}$
- (III) If $\beta > \frac{\alpha+1}{p} + \frac{n}{q}$, then $f \in \wedge_{\beta - \frac{\alpha+1}{p} - \frac{n}{q}}(B_n)$ especially If $\beta = 1$ then $\|f\|_{\wedge_{1 - \frac{\alpha+1}{p} - \frac{n}{q}}} \leq C \|f^{[1]}\|_{p,q,\alpha}$ where B_n is the unit ball of $\mathbb{C}^n$

KEY WORDS AND PHRASES. Fractional derivative, Bergman space, Bloch space, Lipschitz space
1991 AMS SUBJECT CLASSIFICATION CODE. 32A10

Let Ω be a bounded symmetric domain in the complex vector space $\mathbb{C}^n$, $o \in \Omega$, with Bergman-Silov boundary b , Γ the group of holomorphic automorphisms of Ω and Γ_0 its isotropy group. It is known that Ω is circular and star-shaped with respect to o and b is circular. The group Γ_0 is transitive on b and b has a unique normalized Γ_0 -invariant measure σ with $\sigma(b) = 1$. Hua [2] constructed by group representation theory a system $\{\phi_{kv}\}$ of homogeneous polynomials, $k = 0, 1, \dots$, $v = 1, \dots, m_k$, $m_k = \binom{n+k-1}{k}$, complete and orthogonal on Ω and orthonormal on b .

By $H(\Omega)$ we denote the class of all holomorphic functions on Ω . Every $f \in H(\Omega)$ has a series expansion

$$f(z) = \sum_{k,v} a_{kv} \phi_{kv}(z), \quad a_{kv} = \lim_{r \rightarrow 1} \int_b f(r\xi) \overline{\phi_{kv}(\xi)} d\sigma(\xi) \quad (0)$$

where $\sum_{k,v} = \sum_{k=0}^{\infty} \sum_{v=1}^{m_k}$ and the convergence is uniform on a compact subset of Ω .

Let $f \in H(\Omega)$ with the expansion (0) and $\beta > 0$. The β th fractional derivatives of f are defined, respectively, by

$$f^{[\beta]}(z) = \sum_{k,v} \frac{\Gamma(k+1+\beta)}{\Gamma(k+1)} a_{kv} \phi_{kv}(z)$$

$$f_{[\beta]}(z) = \sum_{k,v} \frac{\Gamma(k+1)}{\Gamma(k+1+\beta)} a_{kv} \phi_{kv}(z)$$

It is known that $f^{[\beta]}, f_{[\beta]} \in H(\Omega)$ and

$$f(r\xi) = \frac{1}{\Gamma(\beta)} \int_0^1 (1-\rho)^{\beta-1} f^{[\beta]}(r\rho\xi) d\rho. \quad (1)$$

Let $f \in H(\Omega)$. It will be said that f belongs to the Bergman spaces $A^{p,q,\alpha}(\Omega)$, $0 < p, q \leq \infty$, $\alpha > -1$ if

$$\|f\|_{p,q,\alpha} = \begin{cases} \left(\int_0^1 (1-r)^\alpha M_q(r, f)^p dr \right)^{\frac{1}{p}}, & p < \infty \\ \sup_{0 < r < 1} (1-r)^\alpha M_q(r, f), & p = \infty \end{cases}$$

is finite, where

$$M_q(r, f) = \left(\int_b |f(r\xi)|^q d\sigma(\xi) \right)^{1/q}, \quad 0 < q < \infty$$

and

$$M_\infty(r, f) = \sup_{\xi \in b} |f(r\xi)|$$

see [1,3,5,6,7] for more on $A^{p,q,\alpha}(\Omega)$. For $0 < p \leq \infty$, let $A^p(\Omega)$ denote $A^{p,p,0}(\Omega)$ (see [10,12]), $H^p(\Omega)$ denote $A^{\infty,p,0}(\Omega)$ (see [9]).

Let B_n denote the unit ball in C^n . A function $f \in H(B_n)$ is called a Bloch function, that is $f \in B(B_n)$, if ([8,11])

$$\|f\|_B = \sup_{z \in B_n} (1 - |z|)|f^{[1]}(z)| < \infty$$

For $0 < \alpha < \infty$, the definition of Lipschitz space $\wedge_\alpha(B_n)$ can be found in [4, §8.8]

In [10] and [12], Watanabe and Stojan considered the problem: If $f' \in A^p(D)$ (D is the unit disc of C^1), then $q = ?$ such that $f \in A^q(D)$. In this paper we consider and solve the same problem in $A^{p,q,\alpha}(\Omega)$.

The main results of this paper are the following

THEOREM 1. Let $0 < p, q \leq \infty, \alpha > -1, 0 < \beta < \delta \leq \frac{\alpha+1}{p} + \frac{n}{q}$, if $f^{[\beta]} \in A^{p,q,\alpha}(\Omega)$ and $f^{[\beta]}(r\xi) = O\left(\|f^{[\beta]}\|_{p,q,\alpha}(1-r)^{-\delta}\right)$, then $f \in A^{s,t,\alpha}(\Omega)$ and $\|f\|_{s,t,\alpha} \leq C\|f^{[\beta]}\|_{p,q,\alpha}$, where $s = \frac{\delta p}{\delta - \beta}, t = \frac{\delta q}{\delta - \beta}$.

THEOREM 2. Let $0 < p, q \leq \infty, \alpha > -1, 0 < \beta < \infty, f^{[\beta]} \in A^{p,q,\alpha}(B_n)$.

(I) If $\beta < \frac{\alpha+1}{p} + \frac{n}{q} = \delta$, then $f \in A^{s,t,\alpha}(B_n)$, and $\|f\|_{s,t,\alpha} \leq C\|f^{[\beta]}\|_{p,q,\alpha}$, where s, t are the same as above.

(II) If $\beta = \frac{\alpha+1}{p} + \frac{n}{q}$, then $f \in B(B_n)$ and $\|f\|_B \leq C\|f^{[\beta]}\|_{p,q,\alpha}$.

(III) If $\beta > \frac{\alpha+1}{p} + \frac{n}{q}$, then $f \in \wedge_{\beta - \frac{\alpha+1}{p} - \frac{n}{q}}(B_n)$, especially If $\beta = 1$, then

$$\|f\|_{\wedge_{\beta - \frac{\alpha+1}{p} - \frac{n}{q}}} \leq C\|f^{[1]}\|_{p,q,\alpha}$$

REMARK. (i) Theorem 2(I) ($p = q, \alpha = 0, \beta = n = 1$) extends the results of Watanabe's and Stojan's. (ii) Theorem 1 ($p = \infty$) extends the results of Shi's ([9]) and Lou's ([6,7]).

REFERENCES

- [1] AHERN, P AND JEVITIC, M, Duality and multipliers for mixed norm spaces, *Mich. Math. J.*, **30** (1983), 53-63
- [2] HUA, L K, Harmonic analysis of functions of several variables in the classical domains, *Trans. Amer. Math. Soc.*, **6** (1963).
- [3] JEVITIC, M, Projection theorems, fractional derivatives and inclusion theorems for mixed-norm spaces on the ball, *Analysis*, **9** (1989), 83-105
- [4] KRANTZ, S.G, *Function Theory of Several Complex Variables*, John Wiley & Sons, New York, 1982
- [5] LOU, Z J, A note on a conjecture of S. Axler, *J. Math. Res. & Exp.*, **11** (4) (1991), 629-630 (Chinese)
- [6] LOU, Z J, Hardy-Littlewood type theorem on Bergman spaces, *J. Math. Res. & Exp.*, **11** (3) (1991) (Chinese)
- [7] LOU, Z J, Hardy-Littlewood type theorem on Bergman spaces of several complex variables, *J. Qufu Normal Univ.*, **1** (1992), 111-112 (Chinese)
- [8] LOU, Z J, Characterizations of Bloch functions on the unit ball of C^n , *Kodai Math. J.*, **16** (1993), 74-78
- [9] SHI, J H, Hardy-Littlewood theorems on bounded symmetric domains, *Science in China*, **4** (1988), 366-375
- [10] STOJAN, D, Some new properties of the spaces A^p , *Mat. Vesnik*, **5** 18,33 (1981), 151-157
- [11] TIMONEY, R M, Bloch functions in several complex variables, *Bull. London. Math. Soc.*, **12** (1980), 241-267