ELLIPTIC RIESZ OPERATORS ON THE WEIGHTED SPECIAL ATOM SPACES

KUANG JICHANG

Department of Mathematics
Hunan Normal University
Changsha, Hunan, 410081
P.R. CHINA

(Received July 14, 1993 and in revised form March 3, 1995)

ABSTRACT. In this paper we study the boundedness and convergence of $\sigma_r^s(f)$ and $\tilde{\sigma}_r^s(f)$, the elliptic Riesz operators and the conjugate elliptic Riesz operators of order $s > 0$, on the weighted special atom space $B(\omega)$.

KEY WORDS AND PHRASES. Elliptic Riesz operators, weighted special atom space, Lorentz spaces.

1991 AMS SUBJECT CLASSIFICATION CODES. 42B30, 42B08.

1. INTRODUCTION.

Let R^n be n -dimensional Euclidean space and Z^n be the unit lattice in R^n . The n -Torus T^n is the coset space $R^n/(2\pi Z^n)$, $Q^n = \{x = (x_1, \dots, x_n) : 0 < x_k \leq 2\pi, 1 \leq k \leq n\}$. Let $A(D)$ be a self-adjoint elliptic differential operator with real coefficients defined on $C_0^\infty(R^n)$, $A(D) = \sum_{|\alpha|=m} a_\alpha D^\alpha$, where $D^\alpha = \partial^{|\alpha|}/\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}$, $\alpha = (\alpha_1, \dots, \alpha_n)$ is multi index and $|\alpha| = \alpha_1 + \dots + \alpha_n$. We always assume that the set $\{x \in R^n : A(x) < 1\}$ is convex and its boundary has non-vanishing Gaussian curvature everywhere.

The elliptic Riesz operators and the conjugate elliptic Riesz operators of order $s > 0$ are defined respectively by

$$\sigma_r^s(f, x) = \sum_{m \in Z^n} (1 - A(m/r))_+^s \hat{f}(m) e^{imx} \quad (1.1)$$

$$\tilde{\sigma}_r^s(f, x) = \sum_{m \in Z^n} (1 - A(m/r))_+^s \hat{f}(m) \hat{K}(m) e^{imx} \quad (1.2)$$

where

$$\hat{f}(m) = (2\pi)^{-n} \int_{Q^n} f(x) e^{-imx} dx$$

are the multiple Fourier coefficients of f , $K(x) = P(x)/|x|^{n+m}$ ($x \neq 0$) is a kernel with a homogeneous and harmonic polynomial $P(x)$ of order m , and $\tilde{f}$ is the conjugate function of f with respect to the kernel $K(x)$. $\beta_+ = \max\{0, \beta\}$. If $A(\xi) = |\xi|^2$, $\sigma_r^s(f)$, $\tilde{\sigma}_r^s(f)$ is just the usual Bochner-Riesz mean.

The maximal elliptic Riesz operators defined by

$$\sigma^s(f, x) = \sup_{r > 0} |\sigma_r^s(f, x)|, \tilde{\sigma}^s(f, x) = \sup_{r > 0} |\tilde{\sigma}_r^s(f, x)|.$$

In this paper, using the weighted special atom space $B(\omega)$, we will study the boundedness and convergence of $\sigma_r^s(f)$ and $\tilde{\sigma}_r^s(f)$ for all $s > 0$ and $n = 1$.

We rewrite $B(\omega)$ which was introduced in [4]:

$$B(\omega) = \left\{ f : T \rightarrow R', f(t) = \sum_{k=0}^{\infty} C_k b_k(t), \sum_{k=0}^{\infty} |C_k| < \infty \right\},$$

each b_k is a weighted special atom, that is, a real valued function b , defined on $T = [0, 2\pi]$, which is either $b(t) = 1/(2\pi)$ or $b(t) = \omega(|Q|)^{-1/q}$. $[\chi_R(t) - \chi_L(t)]$, $1 \leq q < \infty$, where Q is an interval in T , L is the left half of Q and R is the right half, $|Q|$ denotes the length of Q , χ_Q the characteristic function of Q and ω is a non-negative real valued function which is increasing, and $\omega(0) = 0$. $B(\omega)$ is endowed with the norm $\|f\|_{B(\omega)} = \inf \left\{ \sum_{k=0}^{\infty} |C_k| \right\}$, where the infimum is taken over all possible representations of f .

$B(\omega)$ is a Banach space.

A function $\omega : [0, \infty) \rightarrow [0, \infty)$ is said to be in the class b_λ ($0 < \lambda < \infty$), if it satisfies

- (1) $\omega(0) = 0$,
- (2) ω is non-decreasing,
 $\omega(t)/t$ is decreasing,
- (4) $\int_0^h \omega(t)/t dt \leq C\omega(h)$, C an absolute constant,
- (5) $\int_h^{2\pi} \omega(t)/t^{\lambda+1} dt \leq C\omega(h)/h^\lambda$ with C independent of h and ω .

Example of functions in the class b_λ are $\omega(t) = t^\alpha$ ($0 < \alpha < 1$) and $\omega(t) = t^\alpha (\log(e/t))^\beta$, ($0 < \alpha < \lambda$, $\beta \geq 0$).

We also define the space $L(\phi)$ be $L(\phi) = \{f : T \rightarrow R', \|f\|_\phi < \infty\}$, where $\|f\|_\phi = (\int_T (f^*(t))^q \phi(t) dt)^{1/q}$, $1 \leq q < \infty$ and f^* is the decreasing rearrangement of f , defined by $f^*(t) = \inf\{y : |\{x : |f(x)| > y\}| \leq t\}$, the outside bars means the Lebesgue measure of the set $\{x : |f(x)| > y\}$, ϕ is a non-negative decreasing function. $\|\cdot\|_\phi$ is a norm if and only if ϕ is a non-negative decreasing function. $L(\phi)$ is a Banach space. If $\omega(t) = (q/p)t^{q/p}$, $1 \leq q \leq p < \infty$, $\phi(t) = \omega(t)/t$, then the space $L(\phi)$ is the Lorentz space $L(p, q)$ in [6,7].

The main result of this paper is stated as follows:

THEOREM 1. Suppose $\omega \in b_\lambda$, $1 \leq \lambda < \infty$, $\phi(t) = \omega(t)/t$, then $\sigma^s(f)$ is of type $(B(\omega), L(\phi))$ for all $s > 0$, that is,

$$\|\sigma^s(f)\|_\phi \leq C\|f\|_{B(\omega)}, \quad f \in B(\omega).$$

COROLLARY 1. Suppose $\omega \in b_\lambda$, $1 \leq \lambda < \infty$, and $f \in B(\omega)$, then $\sigma_r^s(f, x)$ converges to $f(x)$ almost everywhere for all $s > 0$.

THEOREM 2. Suppose $\omega \in b_\lambda$, $1 \leq \lambda < \infty$, $\phi(t) = \omega(t)/t$, then $\tilde{\sigma}^s(f)$ is of type $(B(\omega), L(\phi))$ for all $s > 0$, that is,

$$\|\tilde{\sigma}^s(f)\|_\phi \leq C\|\tilde{f}\|_{B(\omega)} \leq C\|f\|_{B(\omega)}, \quad f \in B(\omega).$$

COROLLARY 2. Suppose $\omega \in b_\lambda$, $1 \leq \lambda < \infty$, and $f \in B(\omega)$, then $\tilde{\sigma}_r^s(f, x)$ converges to $\tilde{f}(x)$ almost everywhere for all $s > 0$.

REMARK 1. When $n = 1$, $A(\xi) = |\xi|^2$, $\sigma_r^s(f, x)$ become

$$\sigma_r^s(f, x) = \sum_{|k| < r} (1 - (|k|/r)^2)^s \hat{f}(k) e^{ikx}. \quad (1.3)$$

As $s \rightarrow 0$, (1.3) become the partial sums of Fourier series of f , when $s = 1/2$, (1.3) are essentially equivalent to the classical Cesàro means. Consequently, the main result in [5,6] become a special case of our results.

REMARK 2. For the maximal (C, α) operators T are defined by

$$T(f, x) = \sup_n |\sigma_n^\alpha(f, x)| \quad (1.4)$$

where

$$\sigma_n^\alpha(f, x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) K_n^\alpha(x-t) dt,$$

since (C, α) kernels

$$K_n^\alpha(t) = \sum_{k=0}^n A_{n-k}^{\alpha-1} D_k(t) / A_n^\alpha \quad (1.5)$$

satisfies

$$|K_n^\alpha(t)| \leq \begin{cases} \frac{An}{(1+nt)(1+(nt)^\alpha)} \leq \frac{An}{1+nt}, & 0 < \alpha < 1, \ 0 \leq t \leq \pi, \\ C/|t|, & \alpha = 1, \ 0 < |t| \leq \pi, \end{cases}$$

thus using the same methods for $\phi(t) = \omega(t)/t$ we can prove

$$\|Tf\|_\phi \leq C\|f\|_{B(\omega)}, \quad f \in B(\omega), \quad 0 < \alpha \leq 1.$$

2. PROOFS OF THEOREMS

PROOF OF THEOREM 1. Let $f^\alpha(x) = f(x-a)$, then the operator $T_a f = f^\alpha$ is of type $(B(\omega), B(\omega))$. Consequently, we just need to prove the result for $f_h(t) = [\omega(2h)]^{-1/q} [\chi_{[-h,0)}(t) - \chi_{[0,h]}(t)]$, $h > 0$ which will follow from the estimate for $g(t) = \chi_{[0,h]}(t)$. Let $H(x) = (2\pi)^{-1} \int_{\mathbb{R}^1} (1-A(y))_+^s e^{ixy} dy$, $s > 0$, $H_{1/r}(x) = rH(rx)$, then $\sigma_r^s(f, x) = (f * K_{1/r})(x)$, where

$$K_{1/r}(x) = \sum_{k=-\infty}^{\infty} H_{1/r}(x+2k\pi).$$

We may assume $r > 1$. By the inequality (see [2]):

$$|H(x)| \leq C(1+|x|)^{-s-1},$$

we get

$$|K_{1/r}(x)| \leq Cr \sum_{k=-\infty}^{\infty} (1+r|2k\pi+x|)^{-(s+1)} \leq Cr(1+r|x|)^{-(s+1)}.$$

Thus

$$\begin{aligned} |\sigma_r^s(g, x)| &= \left| \int_{-\pi}^{\pi} g(y) K_{1/r}(x-y) dy \right| = \left| \int_0^h K_{1/r}(x-y) dy \right| \leq \int_{x-h}^x |K_{1/r}(t)| dt \\ &\leq C \int_{x-h}^x r(1+rt)^{-1} dt \leq Ch(x-h)^{-1} < 2Ch/x, \end{aligned}$$

for $x > 2h$, and $|\sigma_r^s(g, x)| \leq -2Ch/x$ for $x < -2h$. On the other hand, we have

$$\begin{aligned} |\sigma_r^s(g, x)| &\leq \int_0^h |K_{1/r}(x-y)| dy \leq \int_{-\pi}^{\pi} \left(\sum_{k=-\infty}^{\infty} |H_{1/r}(x+2k\pi-y)| \right) dy \\ &= \int_{-\infty}^{\infty} |H_{1/r}(y)| dy \leq C \int_{-\infty}^{\infty} (1+|t|)^{-(s+1)} dt < \infty. \end{aligned}$$

Consequently, we have

$$|\sigma^s(g, x)| \leq \begin{cases} A, & \text{for all } x, \\ 2Ch/|x|, & \text{for } |x| > 2h. \end{cases} \quad (2.1)$$

Let $\phi(t) = \omega(t)/t$. By (2.1) and the conditions on ω , we get

$$\begin{aligned}\|\sigma^s(g)\|_\phi^q &= \int_0^{2\pi} ((\sigma^s(g))^*(x))^q \omega(x)/x dx \leq A^q \int_0^{2h} \omega(x)/x dx \\ &+ (2Ch)^q \int_{2h}^{2\pi} \omega(x)/x^{(q+1)} dx \leq CA^q \omega(2h) + (2Ch)^q (\omega(2h)/(2h)^q) = C\omega(2h).\end{aligned}$$

The constant C may not be the same at every occurrence in this paper. Thus $\|\sigma^s(f_h)\|_\phi \leq 2\omega(2h)^{-(1/q)} \cdot \|\sigma^s(g)\|_\phi \leq C$ and so if $f \in B(\omega)$, then $f(t) = \sum_{k=0}^{\infty} C_k b_k(t)$, where

$$b_k(t) = \omega(|Q_k|)^{-1/q} [\chi_{R_k}(t) - \chi_{L_k}(t)]$$

and $\sum_{k=0}^{\infty} |C_k| < \infty$, we have $\|\sigma^s(f)\|_\phi \leq C \sum_{k=0}^{\infty} |C_k|$, which implies $\|\sigma^s(f)\|_\phi \leq C \|f\|_{B(\omega)}$. The proof is complete.

PROOF OF COROLLARY 1. Let

$$\omega(f, x) = \limsup_{r \rightarrow \infty} |\sigma_{r_1}^s(f, x) - \sigma_{r_2}^s(f, x)|, r_1, r_2 > r, \quad (2.2)$$

then $\omega(f, x) \leq 2\sigma^s(f, x)$ and so

$$\begin{aligned}\|\omega(f)\|_\phi^q &= \int_0^{2\pi} ((\omega(f))^*(x))^q \omega(x)/x dx \leq 2 \int_0^{2\pi} ((\sigma^s(f))^*(x))^q \omega(x)/x dx \\ &= 2\|\sigma^s(f)\|_\phi^q.\end{aligned} \quad (2.3)$$

Since $f \in B(\omega)$, then $f(x) = \sum_{k=0}^{\infty} C_k b_k(x)$, where $\sum_{k=0}^{\infty} |C_k| < \infty$ and the b_k are weighted special atoms.

By Theorem 1 and (2.3), $\sigma^s(f) \in L(\phi)$ which implies $\omega(f) \in L(\phi)$ for $\phi(t) = \omega(t)/t$. On the other hand, we see that $\omega(f) = \omega(f - f_m)$ where $f_m(x) = \sum_{k=0}^m C_k b_k(x)$ and $\|f_m - f\|_{B(\omega)} \rightarrow 0$ as $m \rightarrow \infty$. Then

$$\omega(f, x) = \omega(f - f_m, x) \leq 2\sigma^s(f - f_m, x).$$

By Theorem 1,

$$\|\omega(f)\|_\phi \leq 2\|\sigma^s(f - f_m)\|_\phi \leq 2C\|f - f_m\|_{B(\omega)}.$$

So letting $m \rightarrow \infty$, we get $\|\omega(f)\|_\phi = 0$. Thus $\omega(f, x) = 0$ almost everywhere, which implies $\sigma_r^s(f, x)$ converges to $f(x)$ almost everywhere. The proof is complete.

Let $f \in B(\omega)$, then $f(x) = \sum_{k=0}^{\infty} C_k b_k(x)$, where $\sum_{k=0}^{\infty} |C_k| < \infty$ and the b_k are weighted special atoms. Thus $\tilde{f}(x) = \sum_{k=0}^{\infty} C_k \tilde{b}_k(x)$ and so $\|\tilde{f}\|_{B(\omega)} \leq \|f\|_{B(\omega)}$. Now using $\tilde{\sigma}_r^s(f, x) = \sigma_r^s(\tilde{f}, x)$ and Theorem 1, we can similarly show that Theorem 2 and Corollary 2. The details will be omitted.

REMARK 3. Theorem 1 and 2 are also true if we replace the above $\omega(|Q|)$ by a weight $\omega(Q) = \int_Q \omega(x) dx$, where ω in A_∞ , the proofs are the same.

REFERENCES

- [1] LU, S.Z. and WANG, K.Y., *Bochner-Reisz Means*, Beijing, 1989.
- [2] JIANG, Y.S., LIU, H.P. and LU, S.Z., Some properties of elliptic Riesz means at critical index on $H^p(T^n)$, *Approx. Theory & Its Appl.* 6 (1990), 2, 28-37.
- [3] KUANG JICHANG, Approximation of elliptic Riesz operators on $H^p(T^n)$, *Acta Sci. Nat. Univ. Norm. Hunan* 12 (1989), 1, 1-6, MR90g:41052.
- [4] BLOOM, S. and DeSOUZA, G.S., Atomic decomposition of the generalized Lipschitz spaces, *Illinois J. of Math.* 33 (1989), 2, 181-209.
- [5] DeSOUZA, G.S., Fourier series and the maximal operator on the weighted special atom spaces, *Internat. J. Math. and Math. Sci.* 12 (1989), 3, 579-582.
- [6] DeSOUZA, G.S., On the convergence of Fourier series, *Internat. J. Math and Math. Sci.* 7 (4) (1984), 817-820.
- [7] LORENTZ, G.G., Some new functional spaces, *Ann. of Math.* 51 (1950), 37-55.