A FIXED POINT THEOREM FOR NON-SELF SET-VALUED MAPPINGS

B.E. RHOADES

Department of Mathematics
Indiana University
Bloomington, Indiana 47405, U.S.A.

(Received September 29, 1994)

ABSTRACT. Let X be a complete, metrically convex metric space, K a closed convex subset of X , $CB(X)$ the set of closed and bounded subsets of X . Let $F : K \rightarrow CB(X)$ satisfying definition (1) below, with the added condition that $Fx \subseteq K$ for each $x \in \partial K$. Then F has a fixed point in K . This result is an extension to multivalued mappings of a result of Ćirić [1].

KEY WORDS AND PHRASES. Fixed point, multivalued map, non-self map

1991 AMS SUBJECT CLASSIFICATION CODE. 47H10

Let X be a complete metrically convex metric space. This means that, for each x, y in X , $x \neq y$, there exists a z in ∂X such that $d(x, y) = d(x, z) + d(z, y)$. Let $CB(X)$ denote the set of closed and bounded subsets of X , H denote the Hausdorff metric on $CB(X)$. Let K be a nonempty closed, convex subset of X .

Let $F : K \rightarrow CB(X)$ satisfying: for each x, y in K ,

$$H(Fx, Fy) \leq h \max \left\{ \frac{d(x, y)}{a}, D(x, Fx), D(y, Fy), \frac{[D(x, Fy) + D(y, Fx)]}{a + h} \right\}, \quad (1)$$

where $0 \leq h < (-1 + \sqrt{5})/2$, $a \geq 1 + (2h^2/(1 + h))$, and $F(x) \subseteq K$ for each $x \in \partial K$.

Ćirić [1] proved a fixed point theorem for the single-valued version of (1). He also established a multivalued version. However, he used the δ -distance, instead of the Hausdorff distance, so that the result and proof are identical to the single-valued case. It is the purpose of this paper to prove a multivalued version. For the single-valued version of (1), one can allow h to satisfy $0 \leq h < 1$. However, the multivalued proof requires smaller values of h .

THEOREM. Let X be a complete metrically convex metric space, K a nonempty closed, convex subset of X . Let $F : K \rightarrow CB(X)$ satisfying (1), and the condition that $Fx \subseteq K$ for each $x \in \partial K$. Then F has a fixed point in K .

PROOF. We shall need the following lemma of Nadler [2].

LEMMA. Let $A, B \in CB(X)$, $x \in A$. Then, for each positive number α , there exists a $y \in B$ such that

$$d(x, y) \leq H(A, B) + \alpha.$$

We shall assign $\alpha = h(1 + h)$. We shall now construct a sequence $\{x_n\}$ in K in the following way. Let $x_0 \in K$ and define $x'_1 \in Fx_0$. If $x'_1 \in K$, set $x_1 = x'_1$. If not, then select a point $x_1 \in \partial K$ such that $d(x_0, x_1) + d(x_1, x'_1) = d(x_0, x'_1)$. Then $x_1 \in K$. By the Lemma, choose $x'_2 \in Fx_1$ such that $d(x'_1, x'_2) \leq H(Fx_0, Fx_1) + \alpha$. If $x'_2 \in K$, set $x_2 = x'_2$. Otherwise, choose x_2 so

that $d(x_1, x_2) + d(x_2, x'_2) = d(x_1, x'_2)$. By induction we obtain sequences $\{x_n\}, \{x'_n\}$ such that, for $n = 1, 2, \dots$,

- (i) $x'_{n+1} \in Fx_n$,
- (ii) $d(x'_{n+1}, x'_n) \leq H(Fx_n, x_{n-1}) + \alpha^n$,

where

- (iii) $x'_{n+1} = x_{n+1}$ if $x'_{n+1} \in K$, or
- (iv) $d(x_n, x_{n+1}) + d(x_{n+1}, x'_{n+1}) = d(x_n, x'_{n+1})$ if $x'_{n+1} \notin K$ and $x_{n+1} \in \partial K$.

Now define

$$P := \{x_i \in \{x_n\} : x_i = x'_i, i = 1, 2, \dots\};$$

$$Q := \{x_i \in \{x_n\} : x_i \neq x'_i, i = 1, 2, \dots\}.$$

Note that, if $x_n \in Q$, for some n , then $x_{n-1} \in P$.

For $n \geq 2$ we shall consider $d(x_n, x_{n+1})$. There are three possibilities.

Case 1. $x_n, x_{n+1} \in P$. Then, from (1),

$$\begin{aligned} d(x_n, x_{n+1}) &= d(x'_n, x'_{n+1}) \leq H(Fx_{n-1}, Fx_n) + \alpha^n \\ &\leq h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, D(x_{n-1}, Fx_{n-1}), D(x_n, Fx_n), \frac{D(x_{n-1}, Fx_n) + D(x_n, Fx_{n-1})}{a+h} \right\} + \alpha^n \\ &\leq h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, d(x_{n-1}, x_n), d(x_n, x_{n+1}), \frac{d(x_{n-1}, x_{n+1}) + d(x_n, x_n)}{a+h} \right\} + \alpha^n \\ &\leq \max \left\{ hd(x_{n-1}, x_n) + \alpha^n, \frac{\alpha^n}{1-h}, \frac{hd(x_{n-1}, x_n) + \alpha^n(a+h)}{a} \right\} \\ &\leq hd(x_{n-1}, x_n) + \max \left\{ \frac{1}{1-h}, \frac{a+h}{a} \right\} \alpha^n = hd(x_{n-1}, x_n) + \frac{\alpha^n}{1-h}. \end{aligned} \quad (2)$$

Case 2. $x_n \in P, x_{n+1} \in Q$. Then, from (1),

$$\begin{aligned} d(x_n, x_{n+1}) &\leq d(x_n, x'_{n+1}) \leq H(Fx_{n-1}, Fx_n) + \alpha^n \\ &\leq h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, D(x_{n-1}, Fx_{n-1}), D(x_n, Fx_n), \frac{D(x_{n-1}, Fx_n) + D(x_n, Fx_{n-1})}{a+h} \right\} + \alpha^n \\ &\leq h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, d(x_{n-1}, x'_n), d(x_n, x'_{n+1}), \frac{d(x_{n-1}, x'_{n+1})}{a+h} \right\} + \alpha^n \\ &\leq \max \left\{ hd(x_{n-1}, x_n) + \alpha^n, \frac{\alpha^n}{1-h}, \frac{hd(x_{n-1}, x_n) + \alpha^n(a+h)}{a} \right\} \\ &\leq hd(x_{n-1}, x_n) + \max \left\{ \frac{1}{1-h}, \frac{a+h}{a} \right\} \alpha^n = hd(x_{n-1}, x_n) + \frac{\alpha^n}{1-h}. \end{aligned} \quad (3)$$

Case 3. $x_n \in Q, x_{n+1} \in P$. Note, that $x_n \in Q$ implies that $x_{n-1} \in P$. Using the convexity of X ,

$$d(x_n, x_{n+1}) \leq \max \{d(x_{n-1}, x_{n+1}), d(x'_n, x_{n+1})\} \quad (4)$$

Suppose that the maximum of the right hand side of (4) is $d(x'_n, x_{n+1})$. Then, from (1),

$$\begin{aligned} d(x_n, x_{n+1}) &\leq d(x'_n, x_{n+1}) \leq H(Fx_{n-1}, Fx_n) + \alpha^n \\ &\leq h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, D(x_{n-1}, Fx_{n-1}), D(x_n, Fx_n), \frac{D(x_{n-1}, Fx_n) + D(x_n, Fx_{n-1})}{a+h} \right\} + \alpha^n \\ &\leq h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, d(x_{n-1}, x'_n), d(x_n, x_{n+1}), \frac{d(x_{n-1}, x_{n+1}) + d(x_n, x'_n)}{a+h} \right\} + \alpha^n \end{aligned}$$

Recall that $d(x_{n-1}, x_n) \leq d(x_{n-1}, x'_n)$ and that $d(x_n, x'_n) \leq d(x_{n-1}, x'_n)$. Also, $d(x_{n-1}, x_{n+1}) + d(x_n, x'_n) \leq d(x_{n-1}, x_n) + d(x_n, x_{n+1}) + d(x_n, x'_n) = d(x_{n-1}, x'_n) + d(x_n, x_{n+1})$. Therefore,

$$\begin{aligned} d(x_n, x_{n+1}) &\leq h \max \left\{ d(x_{n-1}, x'_n), d(x_n, x_{n+1}), \frac{d(x_{n-1}, x'_n) + d(x_n, x_{n+1})}{a + h} \right\} + \alpha^n \\ &\leq \max \left\{ hd(x_{n-1}, x'_n) + \alpha^n, \frac{\alpha^n}{1-h}, \frac{hd(x_{n-1}, x'_n) + \alpha^n(a+h)}{a} \right\} \\ &\leq hd(x_{n-1}, x'_n) + \frac{\alpha^n}{1-h}. \end{aligned}$$

Since $x_{n-1} \in P$ and $x_n \in Q$, it follows from Case 2, that

$$d(x_n, x_{n+1}) \leq h^2 d(x_{n-2}, x_{n-1}) + \frac{h\alpha^{n-1}}{1-h} + \frac{\alpha^n}{1-h}. \quad (5)$$

If the maximum of the right hand side of (4) is $d(x_{n-1}, x_{n+1})$, then, from (1),

$$\begin{aligned} d(x_n, x_{n+1}) &\leq d(x_{n-1}, x_{n+1}) \leq d(x_{n-1}, x'_n) + d(x'_n, x_{n+1}) \\ &\leq d(x_{n-1}, x'_n) + H(Fx_{n-1}, Fx_n) + \alpha^n \\ &\leq d(x_{n-1}, x'_n) + h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, D(x_{n-1}, Fx_{n-1}), D(x_n, Fx_n), \right. \\ &\quad \left. [D(x_{n-1}, Fx_n) + D(x_n, Fx_{n-1})] / (a+h) \right\} + \alpha^n \\ &\leq d(x_{n-1}, x'_n) + h \max \left\{ \frac{d(x_{n-1}, x_n)}{a}, d(x_{n-1}, x'_n), d(x_n, x_{n+1}), \right. \\ &\quad \left. [d(x_{n-1}, x_{n+1}) + d(x_n, x'_n)] / (a+h) \right\} + \alpha^n \\ &\leq \max \left\{ (1+h)d(x_{n-1}, x'_n) + \alpha^n, \frac{\alpha^n}{1-h}, \right. \\ &\quad \left. [d(x_{n-1}, x_{n+1}) + d(x_n, x'_n)] / (a+h) \right\} + \alpha^n. \end{aligned} \quad (6)$$

Using (6), if the maximum of the quantity in braces is the third term, then

$$d(x_{n-1}, x_{n+1}) \leq \frac{hd(x_n, x'_n) + (a+h)\alpha^n}{a} \leq \frac{hd(x_{n-1}, x'_n) + (a+h)\alpha^n}{a}.$$

Therefore, by Case 2,

$$\begin{aligned} d(x_n, x_{n+1}) &\leq \max \left\{ (1+h)d(x_{n-1}, x'_n) + \alpha^n, \frac{\alpha^n}{1-h}, \frac{hd(x_{n-1}, x'_n) + (a+h)\alpha^n}{a} \right\} \\ &\leq (1+h)d(x_{n-1}, x'_n) + \frac{\alpha^n}{1-h} \\ &\leq h(1+h)d(x_{n-2}, x_{n-1}) + \frac{h\alpha^{n-1}}{1-h} + \frac{\alpha^n}{1-h}. \end{aligned} \quad (7)$$

Define $\delta = \alpha^{-1/2} \max \{d(x_0, x_1)d(x_1, x_2)\}$. We shall now show that

$$d(x_n, x_{n+1}) \leq \alpha^{n/2}(\delta + 3n), \quad n > 1. \quad (8)$$

The proof is by induction. Note that, for $0 \leq h < (-1 + \sqrt{5})/2$, $(1+h)/(1-h) < 3$, and $1/(1-h) < 3$.

If x_2 and x_3 are such that (3) or (4) is satisfied, then

$$d(x_2, x_3) \leq hd(x_1, x_2) + \frac{\alpha^2}{1-h} \leq h\alpha^{1/2}\delta + 3\alpha^2 < \alpha(\delta + 3),$$

since $h < h(1+h) = \alpha$.

Note that (5) implies (7). If x_2 and x_3 are such that (7) is satisfied, then

$$\begin{aligned} d(x_2, x_3) &\leq h(1+h)d(x_1, x_2) + \frac{(1+h)\alpha}{1-h} + \frac{\alpha^2}{1-h} \\ &\leq \alpha^{3/2}\delta + 3\alpha + 3\alpha^2 \leq \alpha(\delta + 6). \end{aligned}$$

Therefore, in all cases, $d(x_2, x_3) \leq \alpha(\delta + 6)$. Assume the induction hypothesis. If (3) or (4) are satisfied, then

$$\begin{aligned} d(x_n, x_{n+1}) &\leq hd(x_{n-1}, x_n) + \frac{\alpha^n}{1-h} \leq h\alpha^{(n-1)/2}(\delta + 3(n-1)) + 3\alpha^n \\ &\leq \alpha^{n/2}(\delta + 3n) \end{aligned}$$

If (7) is satisfied, then

$$\begin{aligned} d(x_n, x_{n+1}) &\leq h(1+h)d(x_{n-2}, x_{n-1}) + \frac{(1+h)\alpha^{n-1}}{1-h} + \frac{\alpha^n}{1-h} \\ &\leq \alpha^{n/2}(\delta + 3(n-2)) + 3\alpha^{n-1} + 3\alpha^n \leq \alpha^{n/2}(\delta + 3n). \end{aligned}$$

From (8) it follows that, for $m > n$,

$$d(x_n, x_m) \leq \sum_{i=n}^{m-1} d(x_i, x_{i+1}) \leq \delta \sum_{i=n}^{m-1} \alpha^{i/2} + 3 \sum_{i=n}^{m-1} \alpha^{i/2} i,$$

and $\{x_n\}$ is Cauchy, hence convergent. Call the limit p .

Let $\{x_{n_k}\}$ denote the subsequence of $\{x_n\}$ with the property that each term of the subsequence belongs to P . Then

$$\begin{aligned} H(Fx_{n_k-1}, Fp) &\leq h \max\{d(x_{n_k-1}, p)/a, D(x_{n_k-1}, Fx_{n_k-1}), D(p, Fp), \\ &\quad [D(x_{n_k-1}, Fp) + D(p, Fx_{n_k-1})]/(a+h)\} \\ &\leq h \max\{d(x_{n_k-1}, p)/a, d(x_{n_k-1}, x_{n_k}), D(p, Fp), \\ &\quad [D(x_{n_k-1}, Fp) + d(p, x_{n_k})]/(a+h)\}. \end{aligned}$$

Taking the limit as $k \rightarrow \infty$ yields

$$H(p, Fp) \leq hD(p, Fp),$$

which implies, since $H(p, Fp) = D(p, Fp)$, that $p \in Fp$.

REFERENCES

1. ĆIRIĆ, Lj. B., A remark on Rhoades fixed point theorem for non-self mappings, *Internat. J. Math. & Math. Sci.* **16**(1993), 397-400.
2. NADLER, S.B.JR., Multi-valued contraction mappings, *Pacific J. Math.* **30**(1969), 475-488.