SUBRINGS OF I-RINGS AND S-RINGS

MAMADOU SANGHARE

Département de Mathématiques et Informatiques
Faculté des Sciences et Techniques

UCAD

DAKAR (SENEGAL)

e-mail sanghare@ucad.refer.sn

(Received May 6, 1993 and in revised form February 13, 1997)

ABSTRACT. Let R be a non-commutative associative ring with unity $1 \neq 0$, a left R -module is said to satisfy property (I) (resp. (S)) if every injective (resp. surjective) endomorphism of M is an automorphism of M . It is well known that every Artinian (resp. Noetherian) module satisfies property (I) (resp. (S)) and that the converse is not true. A ring R is called a left I-ring (resp. S-ring) if every left R -module with property (I) (resp. (S)) is Artinian (resp. Noetherian). It is known that a subring B of a left I-ring (resp. S-ring) R is not in general a left I-ring (resp. S-ring) even if R is a finitely generated B -module, for example the ring $M_3(K)$ of 3×3 matrices over a field K is a left I-ring (resp. S-ring), whereas its subring

$$B = \left\{ \begin{bmatrix} \alpha & 0 & 0 \\ \beta & \alpha & 0 \\ \gamma & 0 & \alpha \end{bmatrix} / \alpha, \beta, \gamma \in K \right\}$$

which is a commutative ring with a non-principal Jacobson radical

$$J = K \cdot \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} + K \cdot \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

is not an I-ring (resp. S-ring) (see [4], theorem 8). We recall that commutative I-rings (resp. S-rings) are characterized as those whose modules are a direct sum of cyclic modules, these rings are exactly commutative, Artinian, principal ideal rings (see [1]). Some classes of non-commutative I-rings and S-rings have been studied in [2] and [3]. A ring R is of finite representation type if it is left and right Artinian and has (up to isomorphism) only a finite number of finitely generated indecomposable left modules. In the case of commutative rings or finite-dimensional algebras over an algebraically closed field, the classes of left I-rings, left S-rings and rings of finite representation type are identical (see [1] and [4]). A ring R is said to be a ring with polynomial identity (P. I-ring) if there exists a polynomial $f(X_1, X_2, \dots, X_n)$, $n \geq 2$, in the non-commuting indeterminates $X_1, X_2, \dots, X_n$ over the center Z of R such that one of the monomials of f of highest total degree has coefficient 1, and $f(a_1, a_2, \dots, a_n) = 0$ for all $a_1, a_2, \dots, a_n$ in R . Throughout this paper all rings considered are associative rings with unity, and by a module M over a ring R we always understand a unitary left R -module. We use M_R to emphasize that M is a unitary right R -module.

KEY WORDS AND PHRASES: Left I-ring, left S-ring, ring with polynomial identity, ring of finite representation type.

1991 AMS SUBJECT CLASSIFICATION CODES: 16D70, 16P10, 16L60.

1. THE MAIN RESULT

THEOREM. Let R be a left I-ring (resp. S-ring), and B be a sub-ring of R contained in the center Z of R . Suppose that R is a finitely generated flat B -module. Then B is an I-ring (resp. S-ring).

To prove this theorem we need some results.

It is easy to see that

LEMMA 1. Every homomorphic image of a left I-ring (resp. S-ring) is a left I-ring (resp. S-ring).

LEMMA 2. Let P_1 and P_2 be two prime ideals of a ring R . If P_1 is not contained in P_2 then $\text{Hom}_R(R/P_1, R/P_2) = \{0\}$.

PROOF. Let $f: R/P_1 \rightarrow R/P_2$ be an R -homomorphism, and set $f(1 + P_1) = t + P_2$, where $t \in R$. Let $x \in P_1 \setminus P_2$, and let r be any element in R . We have $P_2 = f(xr + P_1) = xrt + P_2$. Thus $xrt \in P_2$. Since P_2 is prime, we have $t \in P_2$, and hence $f = 0$.

LEMMA 3. Let R be a prime ring with polynomial identity. If R is a left I-ring (resp. S-ring), then R is simple Artinian.

PROOF. Let R' be the total ring of fractions of R [5]. It is known that R' is simple Artinian [5], so the R -module R' satisfies (I) (resp. (S)). Since R is a left I-ring (resp. S-ring), then R' is an Artinian (resp. Noetherian) R -module and hence $R' = R$.

LEMMA 4. Let R be a semi-prime ring with polynomial identity. If R is a left I-ring (resp. S-ring), then R is semi-simple Artinian.

PROOF. Let $(P_\ell)_{\ell \in L}$ be a family pairwise distinct minimal prime ideals of R such that

$$\bigcap_{\ell \in L} P_\ell = \{0\}.$$

By Lemma 1 the quotient rings R/P_ℓ ($\ell \in L$) are left I-rings (resp. S-rings) with polynomial identity. Then it follows from Lemma 3 that the rings R/P_ℓ ($\ell \in L$) are simple Artinian, so the left R -modules R/P_ℓ ($\ell \in L$) satisfy (I) (resp. (S)). Following Lemma 1, $\text{Hom}_R(R/P_\ell, R/P_{\ell'}) = \{0\}$ for $\ell \neq \ell'$, so the left R -module $M = \bigoplus_{\ell \in L} R/P_\ell$ satisfies (I) (resp. (S)). Since R is a left I-ring (resp. S-ring), then M is Artinian. But R regarded as left R -module is isomorphic to a submodule of the semi-simple Artinian left R -module M , hence R is semi-simple Artinian.

PROPOSITION 5. Let R be a ring with polynomial identity. If R is a left S-ring (resp. I-ring), then R is left Artinian.

PROOF. Suppose that R is a left S-ring (resp. I-ring) then the quotient ring $R/\text{rad}(R)$, where $\text{rad}(R)$ is the prime radical of R , is a left S-ring (resp. I-ring), so, following Lemma 4, the ring $R/\text{rad}(R)$ is semi-simple Artinian. This fact implies that R is semi-perfect and hence $\text{rad}(R) = J(R)$, where $J(R)$ is the Jacobson radical of R . Let e be a primitive idempotent of R . Since the endomorphism ring of the R -module Re is isomorphic to the local ring eRe with a nil Jacobson radical $eJ(R)e$, then the R -module Re satisfies property (I) (resp. (S)). It follows that the R -module Re is Noetherian (resp. Artinian). Since R regarded as R -module is a direct sum of finitely many left R -modules of the form Re , where e is a primitive idempotent of R , then R is Noetherian. Let P now be a prime ideal of R . Since the prime ring R/P is simple in virtue of Lemma 3, then R is left Artinian.

PROOF OF THE MAIN THEOREM. Since R is a finitely generated Z -module, then R is a ring with polynomial identity (see [6]). So by Proposition 5 R is a left Artinian ring. Thus by [7] the ring B is Artinian. Let $e_1, \dots, e_n$ be primitive idempotents of B such that $B = \bigoplus_{i=1}^n e_i B e_i$. For every i , $1 \leq i \leq n$, $B_i = e_i B e_i$ is a local Artinian ring. To show that B is a left I-ring (resp. S-ring) it is enough to show that for every i , $1 \leq i \leq n$, B_i is a left I-ring (resp. S-ring). We have $A = \bigoplus_{i=1}^n A_i$, where $A_i = e_i A e_i$, $1 \leq i \leq n$. By hypothesis the left B -module $\bigoplus_{i=1}^n A_i = A$ is flat and finitely generated, so the B_i -module

$$A_i = e_i A e_i \cong e_i A e_i \otimes_B B = A \otimes_B e_i B e_i = A \otimes_B B_i$$

is also flat and finitely generated. Since B_i is an Artinian local ring then the B_i -module A_i is faithfully flat (see [8] proposition 1, p. 44).

Suppose now that B_i is not an I-ring (resp. S-ring) for some i , $1 \leq i \leq n$. Then by Proposition 2 of [2], there exists a B_i -module M of infinite length such that, for every integer $n \geq 1$, the B_i -module M^n satisfies both properties (I) and (S). Following [8] (corollary 2, p. 107), the B_i -module A_i is a free module. Let $M' = M \otimes_{B_i} A_i$. Since the B_i -module M is of infinite length and A_i is a faithfully flat B_i -module, then M' is an A_i -module of infinite length. On the other hand, since A_i is a free B_i -module, there exists an integer $s \geq 1$ such that $A_i = B_i^s$. We have then the B_i -module isomorphism

$$M' = M \otimes_{B_i} A_i = M \otimes_{B_i} B_i^s \cong M^s.$$

Hence the B_i -module $M' \cong M^s$ satisfies both properties (I) and (S) and therefore M' , regarded as A_i -module, satisfies properties (I) and (S). This fact implies that the homomorphic image A_i of the left I-ring (resp. S-ring) A is not a left I-ring (resp. S-ring), in contradiction with Lemma 1.

COROLLARY. Let R be a left I-ring (resp. S-ring). If R is a finitely generated flat module over its center Z , then Z is an I-ring (resp. S-ring).

The following example shows that the converse of the theorem above is not true. Let K be a field. The commutative ring $A = K[X, Y]/(X^2, XY, Y^2)$ is not an I-ring (resp. S-ring) because its Jacobson radical $J = K\bar{X} + K\bar{Y}$ is not principal (see [1], theorem 8). On the other hand K is an I-ring (resp. S-ring) and A is a finite-dimensional K -vector space.

ACKNOWLEDGEMENT. The author would like to thank the referee for his valuable suggestions and numerous very useful remarks about the text.

REFERENCES

- [1] KAIDI, A.M. and SANGHARE, M., Une caractérisation des anneaux artiniens à idéaux principaux, *Lec. Notes in Math.*, Vol. 1328, Springer-Verlag, Berlin (1988), 245-254.
- [2] SANGHARE, M., Sur quelques classes d'anneaux liées au lemme de Fitting, *Rend. Sem. Math. Padova*, **87** (1992), 29-37.
- [3] SANGHARE, M., On S-duo-rings, *Comm. in Algebra* **20** (8) (1992), 2183-2189.
- [4] SANGHARE, M., Characterizations of algebras whose modules with Fitting's property are of finite length, *Ext. Mat.* **7** (2) (1992), 1-2.
- [5] POSNER, E.C., Prime rings satisfying a polynomial identity, *Proc. Amer. Math. Soc.* **11** (1960), 180-183.
- [6] RENAULT, G., *Algèbre non-commutative*, Gauthier-Villars, Paris (1975).
- [7] EISENBUD, E., Subrings of Artinian and Noetherian rings, *Math. Ann.* **185** (1970), 247-249.
- [8] BOURBAKI, N., *Algèbre commutative chap. I*, Hermann, Paris (1961).