n -COLOR PARTITIONS WITH WEIGHTED DIFFERENCES EQUAL TO MINUS TWO

A.K. AGARWAL

Mathematical Sciences Division
Institute of Advanced Study in
Science and Technology
Jawahar Nagar, Khanapara
Guwahati - 781022, INDIA

R. BALASUBBRANIAN

The Institute of Mathematical Sciences
C I T. Campus
Madras - 600 113, INDIA

(Received November 6, 1995)

ABSTRACT. In this paper we study those n -color partitions of Agarwal and Andrews, 1987, in which each pair of parts has weighted difference equal to -2 . Results obtained in this paper for these partitions include several combinatorial identities, recurrence relations, generating functions, relationships with the divisor function and computer produced tables. By using these partitions an explicit expression for the sum of the divisors of odd integers is given. It is shown how these partitions arise in the study of conjugate and self-conjugate n -color partitions. A combinatorial identity for self-conjugate n -color partitions is also obtained. We conclude by posing several open problems in the last section.

KEY WORDS AND PHRASES: Partitions, combinatorial identities, recurrence relations, generating functions.

1991 AMS SUBJECT CLASSIFICATION CODES: Primary: 05A15, 05A17, 05A19, Secondary 11B37, 11P81

1. INTRODUCTION, DEFINITIONS AND NOTATIONS

n -color partitions (also called partitions with " n copies of n ") were introduced by Agarwal and Andrews in [2]. These are the partitions in which a part of size n , can come in n different colors denoted by subscripts $n_1, n_2, \dots, n_n$. Thus, for example, the n -color partitions of 3 are

$$\begin{array}{ccc} 3_1 & 3_2 & 3_3 \\ 2_1 1_1 & 2_2 1_1 & \\ 1_1 1_1 1_1 & & \end{array}$$

If $P(\nu)$ denote the number of n -color partitions of ν , then it was shown in [2] that

$$1 + \sum_{\nu=1}^{\infty} P(\nu)q^{\nu} = \prod_{n=1}^{\infty} (1 - q^n)^{-n}. \quad (1.1)$$

It was pointed out in [2] that since the right hand side of (1.1) is also a generating function for the MacMahon's plane partition function so the number of n -color partitions of ν equals the number of plane partitions of ν . In terms of n -color partitions a class of new Rogers-Ramanujan type identities was given in the same paper. Further Rogers-Ramanujan Type identities using n -color partitions were found in [1]. This was one of the advantages of studying n -color partitions as there are no Rogers-Ramanujan Type identities for plane partitions. In this paper we define conjugate and self-conjugate n -color partitions and obtain various combinatorial identities using these definitions. This will give another advantage of studying n -color partitions since there are no self-conjugate plane partitions. It is worthwhile to remark

here that conjugate and self-conjugate $d(n)$ -color partitions, where $d(n)$ is the number of positive divisors of n have been studied by Agarwal and Mullen in [3]

We now give some definitions which we shall use in this paper

DEFINITION 1. Let $\Pi = (a_1)_{b_1} + (a_2)_{b_2} + \dots + (a_r)_{b_r}$ be an n -color partition of ν . We call $(a_i)_{a_i-b_i+1}$ the conjugate of $(a_i)_{b_i}$. An n -color partition of ν obtained from Π by replacing each of its parts by its conjugate will be called the conjugate of Π and will be denoted by Π^c . For example, if we consider $\Pi = 5_2 + 3_1$, an n -color partition of 8, then $\Pi^c = 5_{5-2+1} + 3_{3-1+1} = 5_4 + 3_3$

DEFINITION 2. We shall call an n -color partition to be self-conjugate if it is identical with its conjugate Π^c . Thus $5_3 + 3_2 + 1_1$ is a self-conjugate n -color partition of 9.

DEFINITION 3 (see [2]). The weighted difference of any pair of parts m_i, n_j is defined by $m - i - n - j$.

Throughout this paper $A(\nu)$ will denote the number of n -color partitions of positive integer ν where the weighted difference of each pair of parts is -2 . Thus $A(\nu) = 0$ if $\nu \leq 0$. $A(m, \nu)$ will denote the number of partitions of ν enumerated by $A(\nu)$ with the added restriction that there be exactly m parts. Obviously, $A(m, \nu) = 0$ for $\nu < m$.

We shall also use nearest integer functions $\lfloor x \rfloor$. We define $\lfloor x \rfloor$ as follows

$$\lfloor x \rfloor = r, \quad (1.2)$$

where r is the unique integer such that

$$x - \frac{1}{2} \leq r < x + \frac{1}{2}.$$

2. COMBINATORIAL IDENTITIES

In this section we shall prove several combinatorial identities. Our first identity is an easy consequence of the definition of conjugacy.

THEOREM 2.1. Let $B(\nu)$ denote the number of n -color partitions of ν such that in each pair of parts m_i, n_j ($m \geq n$) n is the arithmetic mean of the subscripts i and j . Then

$$A(\nu) = B(\nu), \quad \forall \nu.$$

EXAMPLE. $A(5) = 11$. The relevant partitions in this case are $5_1, 5_2, 5_3, 5_4, 5_5, 4_1 1_1, 3_1 2_2, 3_2 2_1, 3_3 1_1 1_1, 2_2 1_1 1_1 1_1, 1_1 1_1 1_1 1_1 1_1$.

$B(5)$ is also equal to 11, since in this case the relevant partitions are $5_1, 5_2, 5_3, 5_4, 5_5, 4_1 1_1, 3_3 2_1, 3_2 2_2, 3_1 1_1 1_1, 2_1 1_1 1_1 1_1, 1_1 1_1 1_1 1_1 1_1$.

PROOF. Conjugacy is the natural bijection between the two classes. To see this, let Π be a partition enumerated by $A(\nu)$. That is, each pair of parts m_i, n_j in Π satisfies the condition

$$m - n - i - j = -2. \quad (2.1)$$

We claim that each pair of parts p_q, r_s in Π^c satisfies the condition

$$r = \frac{q+s}{2}. \quad (2.2)$$

We see that

$$\begin{aligned} p_q, r_s \in \Pi^c &\Rightarrow p_{p-q+1}, r_{r-s+1} \in \Pi \\ &\Rightarrow p - (p-q+1) - r - (r-s+1) = -2, \\ &\quad (\text{by (2.1)}) \end{aligned}$$

which is the same as (2.2).

To see the reverse implication let σ be a partition enumerated by $B(\nu)$. That is, each pair of parts $p_q, r_s \in \sigma$ satisfies (2.2). We want to prove that each pair of parts m_i, n_j in σ^c satisfies (2.1).

Now

$$\begin{aligned}
 m_i, n_j \in \sigma^c &\Rightarrow m_{m-i+1}, n_{n-j+1} \in \sigma \\
 &\Rightarrow \frac{(m-i+1) + (n-j+1)}{2} = n
 \end{aligned}$$

which is the same as 2.1 This completes the proof of Theorem 2.1 (by (2.2)).

To illustrate the bijection of Theorem 2.1, we provide the example for $\nu = 6$

Partitions enumerated by $A(6)$	Conjugacy	Partitions enumerated by $B(6)$
6_1	$\longleftrightarrow$	6_6
6_2	$\longleftrightarrow$	6_5
6_3	$\longleftrightarrow$	6_4
6_4	$\longleftrightarrow$	6_3
6_5	$\longleftrightarrow$	6_2
6_6	$\longleftrightarrow$	6_1
$5_5 1_1$	$\longleftrightarrow$	$5_1 1_1$
$4_2 2_2$	$\longleftrightarrow$	$4_3 2_1$
$4_3 2_1$	$\longleftrightarrow$	$4_2 2_2$
$4_4 1_1 1_1$	$\longleftrightarrow$	$4_1 1_1 1_1$
$3_1 3_1$	$\longleftrightarrow$	$3_3 3_3$
$3_3 1_1 1_1 1_1$	$\longleftrightarrow$	$3_1 1_1 1_1 1_1$
$2_1 2_1 2_1$	$\longleftrightarrow$	$2_2 2_2 2_2$
$2_2 1_1 1_1 1_1 1_1$	$\longleftrightarrow$	$2_1 1_1 1_1 1_1 1_1$
$1_1 1_1 1_1 1_1 1_1 1_1$	$\longleftrightarrow$	$1_1 1_1 1_1 1_1 1_1 1_1$

REMARK. Using the idea of conjugacy, partition identities have been obtained for ordinary partitions [4, Theorem 343, p. 274] and for $d(n)$ -color partitions [3, Theorem 4.1, p. 128]. Following the method of proof of Theorem 2.1, one can prove its following generalization.

THEOREM 2.2. Let $A^k(\nu)$ denote the number of n -color partitions of ν such that each pair of parts has a weighted difference equal to $k - 2$. Let $B^k(\nu)$ denote the number of n -color partitions of ν such that each pair of parts m_i, n_j satisfies the condition $i + j - 2n = k$. Then $A^k(\nu) = B^k(\nu)$.

REMARK. Theorem 2.1 is a particular case $k = 0$ of Theorem 2.2.

Theorem 2.1 is a combinatorial identity between two n -color partition functions. Our next theorem is a combinatorial identity between an n -color partition function on one side and an ordinary partition function on the other side.

THEOREM 2.3. Let $C(m, \nu)$ denote the number of ordinary partitions of all numbers $\leq \nu$ with minimum part m and the differences between parts 0 or $m - 1$. Then

$$A(m, \nu) = C(m, \nu). \quad (2.3)$$

EXAMPLE. Consider the case in which $\nu = 11$ and $m = 3$. We see that $A(3, 11) = 5$, since the relevant partitions are $9_9 1_1 1_1, 7_6 2_1 2_1, 6_4 3_1 2_2, 5_3 3_1 3_1, 4_1 4_1 3_2$; also $C(3, 11) = 5$; in this case the relevant partitions are: $3, 3 + 3, 3 + 5, 3 + 3 + 3, 3 + 3 + 5$.

PROOF. For $m = 1$, the theorem is obviously true since $A(1, \nu) = \nu$, the relevant partitions are $\nu_1, \nu_2, \dots, \nu_\nu$; also $C(1, \nu) = \nu$, in this case the relevant partitions are

$$1, 1 + 1, 1 + 1 + 1, \dots, \underbrace{1 + 1 + \dots + 1}_\nu.$$

Now we consider the case when $m \geq 2$. In this case we shall first prove that

$$A(m, \nu) = A(m, \nu - m) + A(m, \nu - 2m + 1) - A(m, \nu - 3m + 1) + 1. \quad (2.4)$$

To prove (2.4), we split the partitions enumerated by $A(m, \nu)$ into three classes

- (i) those that do not contain $k_k (k \geq 1)$ as a part,
- (ii) those that contain 1_1 as a part, and
- (iii) those that contain $k_k, (k > 1)$ as a part but not 1_1

We now transform the partitions in class (i) by deleting 1 from each part ignoring the subscripts. Obviously, the transformed partition still satisfies the weighted difference condition so it will be a partition enumerated by $A(m, \nu - m)$. In class (ii) we observe that 1_1 can appear only with $k_k (k \geq 1)$ in order to satisfy the weighted difference condition. Also, no two parts k_k and l_l with $k > 1$ and $l > 1$ can appear together. This implies that corresponding to each value of $m (m > 1)$, there is one and only one partition in class (ii). Finally, we transform the partitions in class (iii) by replacing k_k by $(k+1)_{k-1}$ and then subtracting 2 from each part. This will give a partition of $\nu - 2m + 1$ into m parts with $(k-1)_{k-1}$ as a part. Therefore, the actual number of partitions which belong to class (iii) is $A(m, \nu - 2m + 1) - A(m, \nu - 3m + 1)$, where $A(m, \nu - 3m + 1)$ is the number of partitions of $\nu - 2m + 1$ into m parts which are free from the parts like $k_k (k \geq 1)$. The above transformations clearly prove (2.4).

Now we set

$$D(m, \nu) = A(m, \nu) - A(m, \nu - m), \quad \forall \nu. \quad (2.5)$$

Then

$$D(m, \nu) = \begin{cases} 0 & \text{if } \nu < m \\ D(m, \nu - 2m + 1) + 1, & \text{if } \nu \geq m \end{cases}. \quad (2.6)$$

Next, we set

$$E(m, \nu) = D(m, \nu) - D(m, \nu - 2m + 1), \quad \forall \nu. \quad (2.7)$$

Then

$$E(m, \nu) = \begin{cases} 0 & \text{if } \nu < m \\ 1 & \text{if } \nu \geq m \end{cases}. \quad (2.8)$$

Hence

$$\begin{aligned} \sum_{\nu=2}^{\infty} E(m, \nu) q^{\nu} &= 0 + \sum_{\nu=m}^{\infty} E(m, \nu) q^{\nu} \\ &= \sum_{\nu=m}^{\infty} q^{\nu} \\ &= \frac{q^m}{1 - q}. \end{aligned}$$

Using the extreme of the foregoing string of equations in (2.7) and then the resulting equation in (2.5), we get

$$\begin{aligned} \sum_{\nu=2}^{\infty} A(m, \nu) q^{\nu} &= \frac{q^m}{(1 - q)(1 - q^m)(1 - q^{2m-1})} \\ &= \sum_{\nu=2}^{\infty} C(m, \nu) q^{\nu} \end{aligned} \quad (2.9)$$

comparing the coefficients of q^{ν} , we get

$$A(m, \nu) = C(m, \nu) \quad \text{for } 2 \leq m \leq \nu.$$

This completes the proof of Theorem 2.3.

NOTE. We remark here that the first part of (2.9) can, alternatively, be proved by using (2.4). Since if $f_m(q) = \sum_{\nu=2}^{\infty} A(m, \nu)q^{\nu}$ then (2.4) yields

$$f_m(q) = q^m f_m(q) + q^{2m-1} f_m(q) - q^{3m-1} f_m(q) + q^m (1-q)^{-1}.$$

3. GENERATING FUNCTIONS

Our discussion in the preceding section leads us to the following

THEOREM 3.1.

$$\begin{aligned} \text{(i)} \quad & \sum_{\nu=1}^{\infty} A(1, \nu)q^{\nu} = \frac{q}{(1-q)^2} \\ \text{(ii)} \quad & \sum_{\nu=2}^{\infty} A(m, \nu)q^{\nu} = \frac{q^m}{(1-q)(1-q^m)(1-q^{2m-1})}, \quad m \geq 2 \\ \text{(iii)} \quad & \sum_{\nu=1}^{\infty} A(\nu)q^{\nu} = \frac{q}{(1-q)^2} + \sum_{m=2}^{\infty} \frac{q^m}{(1-q)(1-q^m)(1-q^{2m-1})}. \end{aligned}$$

4. AN EXPLICIT FORMULA

In this section we give an explicit formula for $A(m, \nu)$. We prove the following

THEOREM 4.1. Let r be the quotient of $\frac{\nu}{m}$ and $\lfloor \cdot \rfloor$ denotes the nearest integer function defined by (1.2) above. Then

$$A(m, \nu) = \begin{cases} 1 & \text{if } m \geq \frac{\nu+1}{2} \\ \sum_{j=0}^{r-1} \left\lfloor \frac{\nu-mj}{2m-1} \right\rfloor, & \text{if } m < \frac{\nu+1}{2}. \end{cases}$$

PROOF. If $m \geq \frac{\nu+1}{2}$, then clearly $A(m, \nu) = 1$ since the only relevant partition is

$$(\nu - m + 1)_{\nu-m+1} \underbrace{1_1 1_1 \dots 1_1}_{m-1 \text{ parts}}$$

so we consider the case when $m < \frac{\nu+1}{2}$. In (2.6) above we write

$$\nu = k(2m-1) + \ell.$$

Then

$$\begin{aligned} D(m, k(2m-1) + \ell) &= D(m, (k-1)(2m-1) + \ell) + 1 \\ &= D(m, (k-2)(2m-1) + \ell) + 2 \\ &\dots \\ &= D(m, \ell) + k \\ &= \begin{cases} k+1 & \text{if } \ell \geq m \\ k & \text{if } \ell < m \end{cases} \end{aligned}$$

Hence

$$D(m, \nu) = \left\lfloor \frac{\nu}{2m-1} \right\rfloor. \quad (4.1)$$

(2.5) in view of (4.1) can be written as

$$A(m, \nu) = \begin{cases} 0 & \text{if } \nu < m \\ A(m, \nu - m) + \left\lfloor \frac{\nu}{2m-1} \right\rfloor & \text{if } \nu \geq m \end{cases}.$$

In (4.2) we write $\nu = rm + s$, then

$$\begin{aligned}
A(m, rm + s) &= a(m, (r-1)m + s) + \left\lfloor \frac{\nu}{2m-1} \right\rfloor \\
&= A(m, (r-2)m + s) + \left\lfloor \frac{\nu-m}{2m-1} \right\rfloor + \left\lfloor \frac{\nu}{2m-1} \right\rfloor \\
&\dots \\
&= \left\lfloor \frac{\nu-(r-1)m}{2m-1} \right\rfloor + \left\lfloor \frac{\nu-(r-2)m}{2m-1} \right\rfloor + \dots + \left\lfloor \frac{\nu}{2m-1} \right\rfloor \\
&= \sum_{j=0}^{r-1} \left\lfloor \frac{\nu-mj}{2m-1} \right\rfloor.
\end{aligned}$$

This proves Theorem 4.1

5. A RELATIONSHIP WITH DIVISOR FUNCTION

In this section we shall prove a relationship between $A(\nu)$ and $d(\nu)$ where $d(\nu)$ denotes the number of positive divisors of ν

THEOREM 5.1. Let a sequence $\{b_\nu\}$ be defined by $b_1 = 1$, $b_\nu = d(2\nu - 1) - d\left(\frac{\nu-1}{2}\right) - 1$, $\nu \geq 2$. Let $\{B_\nu\}$ be the sequence of the partial sums of b_ν . Then

$$b_\nu = A(\nu) - 2A(\nu-1) + A(\nu-2), \quad \nu \geq 1 \quad (5.1)$$

hence,

$$B_\nu = A(\nu) - A(\nu-1), \quad \nu \geq 1. \quad (5.2)$$

PROOF. From Theorem 3.1 (iii), we have

$$\begin{aligned}
(1-q)^2 \sum A(\nu)q^\nu &= q + \sum_{m \geq 2} \frac{q^m(1-q)}{(1-q^m)(1-q^{2m-1})} \\
&= \sum_{m=1}^{\infty} \frac{q^m(1-q)}{(1-q^m)(1-q^{2m-1})} - \frac{q}{1-q} + q \\
&= \sum_{m=1}^{\infty} q^m \left\{ \frac{(1+q^m)(1-q^m) - q(1-q^{2m-1})}{(1-q^m)(1-q^{2m-1})} \right\} - \frac{q^2}{(1-q)} \\
&= \sum_{m=1}^{\infty} q^m \left\{ \frac{1+q^m}{1-q^{2m-1}} - \frac{q}{(1-q^m)} \right\} - \frac{q^2}{(1-q)} \\
&= \sum_{m=1}^{\infty} \frac{q^m}{1-q^{2m-1}} + \sum \frac{q^{2m}}{1-q^{2m-1}} - \sum \frac{q^{m+1}}{1-q^m} - \frac{q^2}{1-q} \\
&= a(q) + b(q) - c(q) - \frac{q^2}{1-q}, \quad \text{say}
\end{aligned}$$

Now writing the even m 's and odd m 's separately in $c(q)$, we get

$$\begin{aligned}
c(q) &= \sum_{\ell \geq 1} \frac{q^{2\ell+1}}{1-q^{2\ell}} + \sum_{\ell \geq 1} \frac{q^{2\ell}}{1-q^{2\ell-1}} \\
&= \sum_{\ell \geq 1} \frac{q^{2\ell+1}}{1-q^{2\ell}} + b(q) = e(q) + b(q), \quad \text{say} \\
q^{-1}e(q) &= \sum_{\ell \geq 1} \frac{q^{2\ell}}{1-q^{2\ell}}; \quad \text{Replacing } q \text{ by } \sqrt{q}, \text{ we get} \\
q^{-\frac{1}{2}}e(q^{\frac{1}{2}}) &= \sum_{\ell \geq 1} \frac{q^\ell}{1-q^\ell} = \sum d(n)q^n
\end{aligned}$$

Hence

$$e\left(q^{\frac{1}{2}}\right)=\sum d(n)q^{n+\frac{1}{2}}$$

$$e(q)=\sum_{n \geq 1} d(n)q^{2n+1}=\sum_{m \geq 1} d\left(\frac{m-1}{2}\right)q^m.$$

Thus

$$c(q)=\sum_{m \geq 1} d\left(\frac{m-1}{2}\right)q^m+b(q).$$

Now

$$a(q)=\sum_{m=1}^{\infty} \frac{q^m}{1-q^{2m-1}}=\sum_{m=1}^{\infty} q^m \sum_{k \geq 1} q^{(2m-1)(k-1)}$$

$$=\sum q^{2mk-m-k+1}$$

Hence

$$q^{-1}a(q^2)=\sum_{m,k} q^{4mk-2m-2k+1}=\sum_{m \geq 1, k \geq 1} q^{(2m-1)(2k-1)}$$

$$=\sum_{n \text{ odd}} d(n)q^n=\sum_{m \geq 1} d(2m-1)q^{2m-1}$$

$$a(q^2)=\sum d(2m-1)q^{2m}.$$

Hence

$$a(q)=\sum_{m \geq 1} d(2m-1)q^m.$$

Putting these results together, we get,

$$(1-q)^2 \sum_{\nu} A(\nu)q^{\nu}=\sum_{m \geq 1} d(2m-1)q^m-\sum d\left(\frac{m-1}{2}\right)q^m-\frac{q^2}{(1-q)}$$

$$\sum_{m \geq 1} \left(d(2m-1)-d\left(\frac{m-1}{2}\right)\right)q^m-\sum_{m \geq 2} q^m.$$

On comparing the coefficients of q^{ν} in the last equation, we are led to (5.1) (5.2) follows from (5.1) immediately once we note that the right-hand side of (5.2) is the n th partial sum of the right-hand side of (5.1). This completes the proof of Theorem 5.1.

REMARK. We remark here that from Theorem 5.1, we get an algorithm to compute the values of $A(\nu)$. The procedure is illustrated in the following table

ν	b_{ν}	B_{ν}	$A(\nu)$
1	1	1	1
2	1	2	3
3	0	2	5
4	1	3	8
5	0	3	11
6	1	4	15
7	-1	3	18
8	3	6	24
9	-2	4	28
10	1	5	33
11	1	6	39
12	1	7	46

From the table it is clear that each $A(\nu)$ equals the sum of the number immediately above it in the table and the number in the same row in the preceding column.

6. n -COLOR PARTITIONS WITH WEIGHTED DIFFERENCES -2 AND $k_k (k \geq 1)$ AS A PART.

It is easily seen that $D(m, \nu)$ defined by (2.5) above is the number of those partitions which are enumerated by $A(m, \nu)$ and have $k_k (k \geq 1)$ as a part. Let $D(\nu)$ equal $\sum_m D(m, \nu)$. Following the methods of preceding sections we can easily prove the following results:

THEOREM 6.1. Let $F(m, \nu)$ denote the number of ordinary partitions of all numbers $\leq \nu$ into parts where the lowest part is m which does not repeat and the differences between parts are 0 or $m-1$. Then

$$D(m, \nu) = F(m, \nu).$$

EXAMPLE. Consider the case when $m = 2$ and $\nu = 11$. We see that $D(2, 11) = 4$, since the relevant partitions are $10_1 1_1, 9_7 2_2, 8_4 3_3, 7_1 4_4$, also, $F(2, 11) = 4$, since in this case the relevant partitions are

$$2, 2 + 3, 2 + 3 + 3, 2 + 3 + 3 + 3.$$

THEOREM 6.2.

$$(i) \quad \sum_{\nu=1}^{\infty} D(1, \nu) q^{\nu} = \frac{q}{1-q}$$

$$(ii) \quad \sum_{\nu=2}^{\infty} D(m; \nu) q^{\nu} = \frac{q^m}{(1-q)(1-q^{2m-1})}$$

$$(iii) \quad \sum_{\nu=1}^{\infty} D(\nu) q^{\nu} = \frac{q}{(1-q)} + \sum_{m=2}^{\infty} \frac{q^m}{(1-q)(1-q^{2m-1})}.$$

THEOREM 6.3.

$$D(m, \nu) = \left\lfloor \frac{\nu}{2m-1} \right\rfloor.$$

THEOREM 6.4. Let k be the quotient of $\frac{\nu}{2m-1}$. Then

$$A(m, \nu) = \sum_{j=1}^{k-1} D(m, \nu - mj).$$

THEOREM 6.5.

$$D(\nu) - D(\nu - 1) = \begin{cases} 1 & \text{if } \nu = 1 \\ d(2\nu - 1) - 1 & \text{if } \nu > 1 \end{cases}.$$

Using Theorems 6.3 and 6.5, we shall now prove the following explicit expression for the sum of the divisors of odd numbers:

THEOREM 6.6.

$$\sum_{j=1}^{\nu} d(2j - 1) = \sum_{m=1}^{\nu} \left\lfloor \frac{\nu}{2m-1} \right\rfloor.$$

PROOF. Replacing ν by $1, 2, \dots, \nu$ in Theorem 6.5 and then adding all equations, we get

$$D(\nu) = \sum_{j=1}^{\nu} d(2j-1) - (\nu-1),$$

that is

$$1 + \sum_{m \geq 2}^{\nu} D(m, \nu) = \sum_{j=1}^{\nu} d(2j-1) - (\nu-1).$$

This proves Theorem 6.6 by Theorem 6.3

7. A COMBINATORIAL IDENTITY FOR SELF-CONJUGATE n -COLOR PARTITIONS

In this section we shall prove a combinatorial identity which involves self-conjugate n -color partitions defined in Section 1 above.

THEOREM 7.1. Let $F(\nu)$ denote the number of self-conjugate n -color partitions of ν and $G(\nu)$ denote those n -color partitions of ν where each pair of parts has weighted difference greater than 1 and even parts appear with even subscripts and odd with odd. Then $F(\nu) = G(\nu)$ for all ν .

EXAMPLE. $F(7) = 5$; the relevant partitions are

$$7_4, 5_3 + 1_1 + 1_1, 3_2 + 3_2 + 1_1, 3_2 + 1_1 + 1_1 + 1_1 + 1_1, 1_1 + 1_1 + 1_1 + 1_1 + 1_1 + 1_1 + 1_1$$

$$G(7) = 5, \text{ in this case the relevant partitions are } 7_1, 7_3, 7_5, 7_7, 6_2 + 1_1$$

PROOF. We observe that if an n -order partition is self-conjugate then in each part m_i of it, m must be odd. Because, $m_i = m_{m-i+1} \Rightarrow m = 2i - 1$. Thus if we ignore the subscripts of all parts in a self-conjugate n -color partition of ν , we get a unique ordinary partition of ν into odd parts. Conversely, if we consider an ordinary partition of ν into odd parts and replace each part $2a - 1$ by $(2a - 1)_a$. We get a unique self-conjugate n -color partition of ν . This bijection shows that the number of self-conjugate n -color partitions of ν equals the number of ordinary partitions of ν into odd parts. That is,

$$1 + \sum_{\nu=1}^{\infty} F(\nu)q^{\nu} = \prod_{n=1}^{\infty} \frac{1}{1 - q^{2n-1}}.$$

Now an appeal to the following theorem [1, Theorem 1.4, p. 301]

“Let $R(\nu)$ denote the number of n -color partitions of ν such that each pair of parts has weighted difference greater than 1 and even parts appear with even subscripts and odd with odd. Let $S(\nu)$ denote the number of ordinary partitions of ν into distinct parts. Then

$$R(\nu) = S(\nu)”$$

proves Theorem 7.1.

8. A TABLE FOR $A(m, \nu)$

In this section we give a brief table of $A(m, \nu)$ which was obtained on computer by using the recurrence relation (2.4) above. The following observations can serve as a check on the table

Observation 1	$A(1, \nu)$	$= \nu,$	$\forall \nu$
Observation 2	$A(m, \nu)$	$= 1,$	if $m \geq \frac{\nu+1}{2}$
Observation 3	$A(m, 2m)$	$= 2$	
Observation 4	$A(m, 2m+1)$	$= 2$	

We remark here that Observation 1 was proved in the proof of Theorem 2.3 while Observation 2 in the proof of Theorem 4.1. Observations 3 and 4 follow immediately once we note that the only partitions enumerated by $A(m, 2m)$ are

$$(m+1)_{m+1} \underbrace{1_1 1_1 \dots 1_1}_{m-1} \text{ and } \underbrace{2_1 2_1 \dots 2_1}_m,$$

and those enumerated by $A(m, 2m + 1)$ are

$$(m+2)_{m+2} \underbrace{1_1 1_1 \dots 1_1}_{m-1} \quad \text{and} \quad 3_2 \underbrace{2_1 2_1 \dots 2_1}_{m-1}.$$

ν/m	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	1														
2	2	1													
3	3	1	1												
4	4	2	1	1											
5	5	3	1	1	1										
6	6	4	2	1	1	1									
7	7	5	2	1	1	1	1								
8	8	7	3	2	1	1	1	1							
9	9	8	4	2	1	1	1	1	1						
10	10	10	4	2	2	1	1	1	1	1					
11	11	12	5	3	2	1	1	1	1	1	1				
12	12	14	6	4	2	2	1	1	1	1	1	1			
13	13	16	7	4	2	2	1	1	1	1	1	1	1		
14	14	19	8	4	3	2	2	1	1	1	1	1	1	1	
15	15	21	9	5	4	2	2	1	1	1	1	1	1	1	1

9. CONCLUSION

The most obvious questions arising from this work are:

1. Do the combinatorial identities of this paper have their analytic counterparts? Our main hope in pursuing the analytic aspects of these results is that we shall find q -identities resembling Euler's identities [4, p. 277] or Rogers-Ramanujan identities [4, p. 290] which will shed light on how to proceed with further study of the identities of this paper.
2. Is it possible to prove the identities of this paper graphically?
3. By using n -color partitions we have given an explicit expression for $\sum_{j=1}^{\infty} d(2j-1)$. Is it possible to find similar expressions for $\sum_{j=1}^{\infty} d(2j)$?

ACKNOWLEDGMENT. The authors would like to thank Dr. Venkatesh Raman for his computer assistance.

REFERENCES

- [1] AGARWAL, A.K., Rogers-Ramanujan identities for n -color partitions, *J. Number Theory* **28** (1988), 299-305.
- [2] AGARWAL, A.K. and ANDREWS, G.E., Rogers-Ramanujan identities for partitions with " N copies of N ", *J. Combin. Theory Ser. A* **45**, No. 1 (1987), 40-49.
- [3] AGARWAL, A.K. and MULLEN, G.L., Partitions with " $d(a)$ copies of a ", *J. Combin. Theory Ser. A*, **48** (1) (1988), 120-135.
- [4] HARDY, G.H. and WRIGHT, E.M., *An Introduction to the Theory of Numbers*, Oxford Univ. Press, 5th ed., London, 1979.