ON AN INVERSE TO THE HÖLDER INEQUALITY

J. PEČARIĆ

Faculty of Textile Technology
Zagreb University
Zagreb

C.E.M. PEARCE

Applied Mathematics Department
University of Adelaide
Adelaide 5005
South Australia

(Received October 5, 1994)

ABSTRACT. An extension is given for the inverse to Hölder's inequality obtained recently by Zhuang.

KEY WORDS AND PHRASES. Inverse Hölder inequality.

1991 AMS SUBJECT CLASSIFICATION CODE. 26D15.

Recently Zhuang [1] proved the following inverse of the arithmetico-geometric inequality.

THEOREM A. Let $0 < a \leq x \leq A$, $0 < b \leq y \leq B$, $1/p + 1/q = 1$, $p > 1$; then

$$\frac{x}{p} + \frac{y}{q} \leq \max \left\{ \frac{A/p + b/q}{A^{1/p}b^{1/q}}, \frac{a/p + B/q}{a^{1/p}B^{1/q}} \right\} x^{1/p}y^{1/q}, \quad (1)$$

or

$$x + y \leq \max \left\{ \frac{A + b}{A^{1/p}b^{1/q}}, \frac{a + B}{a^{1/p}B^{1/q}} \right\} x^{1/p}y^{1/q}, \quad (2)$$

the sign of equality in (1) and (2) holds if and only if either $(x, y) = (a, B)$ or $(x, y) = (A, b)$.

Moreover, if $a \geq B$, then

$$\frac{a/p + B/q}{a^{1/p}B^{1/q}} x^{1/p}y^{1/q} \leq \frac{x}{p} + \frac{y}{q} \leq \frac{A/p + b/q}{A^{1/p}b^{1/q}} x^{1/p}y^{1/q}, \quad (3)$$

the sign of equality on the right-hand side of (3) holds if and only if $(x, y) = (A, b)$, and the sign of equality on the left-hand side of (3) holds if and only if $(x, y) = (a, B)$. The sign of inequality in (3) is reversed if $b \geq A$.

This enables us to formulate the following theorem.

THEOREM 1. Suppose x, y, a, b, A, B, p, q are as in Theorem A and $\alpha, \beta > 0$. Then

$$\alpha x^p + \beta y^q \leq \max(C, D)xy, \quad (4)$$

where

$$C = (\alpha A^p + \beta b^q)/(Ab), \quad D = (\alpha a^p + \beta B^q)/(aB). \quad (5)$$

Equality occurs if and only if either $(x, y) = (a, B)$ or $(x, y) = (A, b)$. Moreover, if $\alpha pa^p \geq \beta qB^q$, then

$$Cxy \leq \alpha x^p + \beta y^q \leq Dxy, \quad (6)$$

with equality on the right-hand side if and only if $(x, y) = (A, b)$ and on the left if and only if $(x, y) = (a, B)$. The inequalities in (6) are reversed if $\alpha pa^p \leq \beta qB^q$.

PROOF. Inequalities (4) and (6) follow from (1) and (3) under the substitutions

$$x \rightarrow \alpha p x^p, \quad y \rightarrow \beta q y^q, \quad a \rightarrow \alpha p a^b, \quad b \rightarrow \beta q b^q, \quad A \rightarrow \alpha p A^p, \quad B \rightarrow \beta q B^q.$$

REMARK. Theorem 1 gives (1) and (2) together, (1) resulting from the substitutions $\alpha = 1/p$, $x \rightarrow x^{1/p}$, $A = A^{1/p}$, $a \rightarrow a^{1/p}$ and corresponding relations for β, y etc. with q in place of p , while (2) results from similar substitutions with $\alpha = 1 = \beta$.

The following result now gives an extension of the inverse to Hölder's inequality obtained in [1]. We suppose that all the integrals involved exist.

THEOREM 2. Let the functions f, g satisfy $0 < a \leq f(x) \leq A$, $0 < b \leq g(x) \leq B$ for almost all $x \in X$ with respect to a measure μ . Suppose $\alpha, \beta, p, q, C, D$ are as in Theorem 1. Then

$$\left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q} \leq (\alpha\beta)^{-1/p} (\beta q)^{-1/q} \max(C, D) \int_X fg d\mu \quad (7)$$

and equality holds if and only if

$$\mu(E_1 \cup F_1) = \mu(X)$$

and

$$\mu(E_1) = \frac{(\alpha p A^p - \beta q b^q) \mu(X)}{\alpha p (A^p - a^p) + \beta q (B^q - b^q)},$$

where

$$E_1 = \{x \in X : f(x) = a, g(x) = B\},$$

$$F_1 = \{x \in X : f(x) = A, g(x) = b\}.$$

Moreover, if $\alpha p a^p \geq \beta q B^q$, then

$$\left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q} \leq (\alpha p)^{-1/p} (\beta q)^{-1/q} D \int_X fg d\mu, \quad (8)$$

with equality only if $(f, g) = (a, B)$ a.e. on X and $\alpha p a^p = \beta q B^q$, and if $\alpha p A^p \leq \beta q b^q$, then

$$\left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q} \leq (\alpha p)^{-1/p} (\beta q)^{-1/q} C \int_X fg d\mu, \quad (9)$$

with equality only if $(f, g) = (A, b)$ a.e. on X and $\alpha p A^p = \beta q b^q$.

PROOF. The first statement was proved in [1]. A simple proof of the remainder of the theorem was given for the case $\alpha = 1/p$, $\beta = 1/q$ in [2]. We give a similar simple proof for the general case.

$$\begin{aligned} \max(C, D) \int_X fg d\mu &= \int_X \max(C, D) fg d\mu \\ &\geq \int_X (\alpha f^p + \beta g^q) d\mu \\ &= \frac{1}{p} (\alpha p) \int_X f^p d\mu + \frac{1}{q} (\beta q) \int_X g^q d\mu \\ &\geq (\alpha p)^{1/p} (pq)^{1/q} \left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q}, \end{aligned}$$

by the arithmetico-geometric inequality.

The equality conditions result from those in Theorem 1 and the arithmetico-geometric inequality.

Similarly we can prove (8). Using the second inequality in (6) we have

$$\begin{aligned} D \int_X fg \, d\mu &= \int_X Dfg \, d\mu \\ &\geq \int_X (\alpha f^p + \beta g^q) d\mu \\ &= \frac{1}{p} (\alpha p) \int_X f^p d\mu + \frac{1}{q} (\beta q) \int_X g^q d\mu \\ &\geq (\alpha p)^{1/p} (\beta q)^{1/q} \left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q}. \end{aligned}$$

Relation (9) follows similarly.

REMARK. The simplest cases of (8) and (9) occur for $\alpha = 1/p$, $\beta = 1/q$. Then we have that if $a^p \geq B^q$, then

$$\left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q} \leq D_1 \int_X fg \, d\mu$$

and if $A^p \leq b^q$, then

$$\left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q} \leq C_1 \int_X fg \, d\mu,$$

where

$$D_1 = \left(\frac{1}{p} a^p + \frac{1}{q} B^q \right) / (aB),$$

$$C_1 = \left(\frac{1}{p} A^p + \frac{1}{q} b^q \right) / (Ab).$$

REFERENCES

1. ZHUANG, YA-DONG. On inverses of the Hölder inequality, J. Math. Anal. Applic., **161** (1991), 566-575.
2. MOND, B. and PEČARIĆ, J.E. Remark on a recent converse of Hölder's inequality, J. Math. Anal. Applic., **181** (1994), 280-281.