LIMIT SETS IN PRODUCT OF SEMI-DYNAMICAL SYSTEMS

RAMJEE PRASAD BHAGAT

(Received 12 November 1996 and in revised form 9 March 1998)

ABSTRACT. Continuing the study of the properties of Poisson stability and distality [4], we mention the conditions under which $\Omega_x(x) = \Pi\Omega_\alpha(x_\alpha), \alpha \in I$ and thus, the product of Poisson stable motions remains Poisson stable in the product system.

Keywords and phrases. Semi-dynamical system, Lagrange stability, distality, limit sets.

1991 Mathematics Subject Classification. 47H10, 54H25.

1. Introduction. We deal mainly with the product of w -limit sets in the product space of semi-dynamical systems (s.d.s.). In [1], Prem Bajaj has shown that the product of semi-dynamical systems is a semi-dynamical system. He has also shown that $\Pi\Omega_\alpha(x_\alpha), \alpha \in I$ contains the w -limit set $\Omega_x(x)$ of x in the product system. In general, equality does not hold in the above. Indeed $\Omega_x(x)$ may be empty. He has given two theorems: one in which $\Omega_x(x)$ is nonempty and the other indicating a case of equality viz. Theorems 2.3 and 2.4.

In this paper, continuing the study of the properties of Poisson stability and distality [4], we mention the conditions under which $\Omega_x(x) = \Pi\Omega_\alpha(x_\alpha), \alpha \in I, x = \{x_\alpha\}$ and therefore, the product of Poisson stable motions, under these conditions, is Poisson stable.

2. Definitions and notations

DEFINITION 2.1. A continuous mapping $\pi : X \times \mathbb{R}^+ \rightarrow X$ on a topological space X is said to define a semi-dynamical system (X, π) if $\pi(x, 0) = x$ and $\pi(\pi(x, t), s) = \pi(x, t + s)$ for every $x \in X$ and $t, s \in \mathbb{R}^+$. ($\mathbb{R}^+$ denotes the set of nonnegative reals.)

DEFINITION 2.2. Let $(X_\alpha, \pi_\alpha), \alpha \in I$ be a family of dynamical systems. Let $X = \Pi X_\alpha$ be the product space. Let $x \in X$ and $x = \{x_\alpha\}$. Define a map π from $X \times \mathbb{R}$ into X by $\pi(x_\alpha t) = (x_\alpha t), \alpha \in I$, then (X, π) is a dynamical system. The dynamical system (X, π) , obtained above, is called the direct product or the product of the family $(X_\alpha, \pi_\alpha), \alpha \in I$.

We take the usual definitions of positive limit set Ω_x , positive distal, positive Poisson stable, and positive Lagrange stable motions. As usual, we drop the word positive and we use the notations of [1, 4].

3. Main results

PROPOSITION 3.1. Let $(X_\alpha, \pi_\alpha), \alpha \in I$, be a family of $\{\text{Lagrange stable}\} \{\text{distal}\}$ s.d.s.

and (X, π) the product s.d.s. Let $x \in X$ and $x = \{x_\alpha\}$, then (X, π) is {Lagrange stable} {distal}.

PROPOSITION 3.2. *If a Lagrange stable motion is Poisson stable and distal, then $\text{Cl}Y(x) = Y(x) = \Omega_x$.*

PROOF. The proof follows from [4, Thm. 2.1]. □

THEOREM 3.3. *Let (X_α, π_α) , $\alpha \in I$, be a family of dynamical systems and (X, π) the product of the dynamical systems. Let $x \in X$ and $x = \{x_\alpha\}$. Then $\Omega_x(x) \subseteq \Pi\Omega_\alpha(x_\alpha)$, where $\Omega_\alpha(x_\alpha)$ is the positive limit set of x_α in the dynamical systems (X_α, π_α) . (The two π 's have distinct meanings according to the context.)*

Since, in general, the equality does not hold and Ω_x may be empty, the Poisson stability in the constituent dynamical system may be lost from the product of the dynamical systems. Here, we find the conditions under which $\Omega_x(x) = \Pi\Omega_\alpha(x_\alpha)$, $\alpha \in I$ and thus, the product of Poisson stable motions remains Poisson stable in the product system.

THEOREM 3.4. *If a compact motion is Poisson stable and distal, then it is a compact recurrent motion.*

PROOF. Let the motion $\pi(x, t)$ be Poisson stable and distal, then its trajectory $Y(x)$ is closed. Therefore,

$$Y(x) = \text{Cl}Y(x) = \Omega_x. \quad (3.1)$$

As the motion is compact, each of the above sets is compact and minimal and thus, by Birkhoff recurrence theorem, $\pi(x, t)$ is compact and recurrent. □

THEOREM 3.5. *Let (X, π) be a semi-dynamical system. Let π be a Lagrange stable, then π is distal if and only if, for every net t_i in $\mathbb{R}^+$, the phase space*

$$X = \{z \in X : xt_j \rightarrow z \text{ for some } x \in X \text{ and some subnet } t_j \text{ of } t_i\} \quad (3.2)$$

[2, Thm. 2.6].

THEOREM 3.6. *Let (X, π) be Lagrange stable and distal s.d.s. then every net in the trajectory $Y(x)$ of the Poisson stable motion $\pi(x, t)$ is a Cauchy net.*

PROOF. Let $Y(x)$ be the trajectory of the Poisson stable motion $\pi(x, t)$ in s.d.s. (X, π) which is Lagrange stable and distal. Let xt_n be a net in $Y(x)$ which is compact (Proposition 3.2). Therefore, xt_n has a subnet, say xt_m with $xt_m \rightarrow z$, i.e., z is a cluster point of xt_n . Hence, xt_n is a Cauchy net. □

THEOREM 3.7. *Let (x_α, π_α) , $\alpha \in I$, be a family of Lagrange stable and distal s.d.s. and (X, π) be the product s.d.s. Let $x \in X$ and $x = \{x_\alpha\}$. A motion $\pi(x, t)$ is Poisson stable in (X, π) if and only if $\pi_\alpha(x_\alpha, t)$ is Poisson stable in (X_α, π_α) for each $\alpha \in I$.*

PROOF. Let (x_α, π_α) , $\alpha \in I$, be a Lagrange stable and distal s.d.s. Let $\pi(x_\alpha, t) = x_\alpha t$ be a Poisson stable motion in (X_α, π_α) , $\alpha \in I$, then its trajectory $Y_\alpha(x_\alpha)$ is compact and the net $x_\alpha t_n$, $\alpha \in I$, is a Cauchy net in $Y_\alpha(x_\alpha)$ (Theorem 3.6). Now, the Cauchy

nets $x_\alpha t_n, \alpha \in I$ yield the Cauchy net xt_n in $Y(x)$ in (X, π) [3, p. 194]. As the product of compact sets is a compact set, $Y(x)$ is compact and xt_n is a net in compact $Y(x)$. Thus, it has a subnet $xt_m \rightarrow z$, i.e., z is a cluster point of xt_n . Hence, xt_n is frequently in every neighborhood U of z . Given a neighborhood U of z for every $i \in A$, there is a $j \in A, i \geq J$ such that $xt_i \in U$ however $t_i \rightarrow +\infty$. Hence, $\pi(x, t)$ is Poisson stable. The converse follows from [3, Thm. 25, p. 194] which states that a net in the product is a Cauchy net if and only if its projection into each coordinate space is a Cauchy net. $\square$

THEOREM 3.8. *Let $(X_\alpha, \pi_\alpha), \alpha \in I$, be a family of Lagrange stable distal s.d.s. Let $x \in X, x = \{x_\alpha\}$, and (X, π) the product s.d.s. Let $Y_\alpha(x_\alpha), \alpha \in I$, be the product of trajectories. Then $\Pi Y_\alpha(x_\alpha) = Y(x)$. Moreover,*

$$\Pi \Omega_\alpha(x_\alpha) = \Omega_x(x). \quad (3.3)$$

PROOF. Since each $Y_\alpha(x_\alpha), \alpha \in I$, is closed and compact,

$$\text{Cl} \Pi Y_\alpha(x_\alpha) = \Pi \text{Cl} Y_\alpha(x_\alpha) = \text{Cl} Y(x), \quad (3.4)$$

$$\Pi Y_\alpha(x_\alpha) = Y(x). \quad (3.5)$$

Moreover,

$$\Pi \Omega_\alpha(x_\alpha) = \Omega_x(x). \quad (3.6)$$

$\square$

REFERENCES

- [1] P. N. Bajaj, *Products of semi-dynamical systems*, Recent advances in topological dynamics (Proc. Conf., Yale Univ., New Haven, Conn., 1972; in honor of Gustav Arnold Hedlund) (Berlin), Lecture Notes in Math, vol. 318, Springer, Verlag, 1973, pp. 23-29. MR 53 1549. Zbl 258.54042.
- [2] N. P. Bhatia and M. Nishihama, *Distal semidynamical systems*, Dynamical systems (Proc. Internat. Sympos., Brown Univ., Providence, R.I., 1974) (New York), vol. II, Academic Press, 1976, pp. 187-190. MR 58 31006. Zbl 359.54033.
- [3] J. L. Kelley, *General topology*, Springer-Verlag, New York, Berlin, 1975. MR 51 6681. Zbl 518.54001.
- [4] S. S. Prasad and A. Kumar, *Stable P and distal dynamical systems*, Internat. J. Math. Math. Sci. 7 (1984), no. 1, 181-185. MR 85e:34045. Zbl 562.54061.

BHAGAT: DEPARTMENT OF MATHEMATICS, A. S. COLLEGE BIKRAMGANJ, PIN 802212, ROHTAS, BIHAR, INDIA