NONLINEAR FUNCTIONAL INTEGRODIFFERENTIAL EQUATIONS IN HILBERT SPACE

J. Y. PARK, S. Y. LEE, and M. J. LEE

(Received 7 November 1996 and in revised form 3 November 1997)

ABSTRACT. Let X be a Hilbert space and let $\Omega \subset R^n$ be a bounded domain with smooth boundary $\partial\Omega$. We establish the existence and norm estimation of solutions for the parabolic partial functional integro-differential equation in X by using the fundamental solution.

Keywords and phrases. Functional integro-differential equation, elliptic differential operators, fundamental solution, Gårding's inequality, successive approximation, norm estimation.

1991 Mathematics Subject Classification. 35A05, 35J60, 45K05.

1. Introduction. Let X be a Hilbert space and let $\Omega \subset R^n$ be a bounded domain with smooth boundary $\partial\Omega$. We consider the following parabolic partial functional integro-differential equation.

$$\begin{aligned} \frac{\partial u}{\partial t} = & \mathcal{A}_0 u(t, x) + \mathcal{A}_1 u(t-h, x) + \int_{-h}^0 a(s) \mathcal{A}_2 u(t+s, x) ds \\ & + \int_0^t \{k(t, s) G(s, u(s-h), x) + H(t, s, u(s-h, x))\} ds \\ & + F(t, u(t-h, x)) + f(t, x), \quad 0 < t \leq T, x \in \Omega, \end{aligned} \quad (1.1)$$

where $\mathcal{A}_i (i = 0, 1, 2)$ are elliptic differential operators, f is a forcing function, $h > 0$ is a delay time, $a(s)$ is a real scalar function on $[-h, 0]$, G, H , and F are nonlinear functions, and k is a kernel. The boundary condition attached to (1.1) is, e.g., given by the Dirichlet boundary condition

$$u|_{\partial\Omega} = 0, \quad 0 < t \leq T, \quad (1.2)$$

and the initial condition is given by

$$u(\theta, x) = g(\theta, x), \quad \theta \in [-h, 0], x \in \Omega. \quad (1.3)$$

From [4], the above mixed problems (1.1), (1.2), and (1.3) can be formulated abstractly as

$$\begin{aligned} \frac{du(t)}{dt} = & A_0 u(t) + A_1 u(t-h) + \int_{-h}^0 a(s) A_2 u(t+s) ds \\ & + \int_0^t \{k(t, s) G(s, u_s) + H(t, s, u_s)\} ds \end{aligned} \quad (1.4)$$

$$\begin{aligned} & + F(t, u_t) + f(t), \quad 0 < t \leq T, \\ & u(\theta) = g(\theta), \quad \theta \in [-h, 0], \end{aligned} \quad (1.5)$$

where the state $u(x)$ of the system (1.5) lies in an appropriate Hilbert space and $A_i (i = 0, 1, 2)$ are unbounded operators associated with $\mathcal{A}_i (i = 0, 1, 2)$, respectively. Next, we explain the notation u_t in (1.5). Let $I = [-h, 0]$. If a function $u(t)$ is continuous from $I \cup [0, T]$ into a Hilbert space X , then u_t is an element in $C = C([-h, 0]; X)$, which has the point-wise definition

$$u_t(\theta) = u(t + \theta) \quad \text{for } \theta \in I. \quad (1.6)$$

Let $\Delta_T = \{(s, t); 0 \leq s \leq t \leq T\}$. We assume in (1.5) that $G : [0, T] \times C \rightarrow X$, $H : \Delta_T \times C \rightarrow X$, $F : [0, T] \times C \rightarrow X$ and the kernel $k : \Delta_T \rightarrow R$ (R denotes the set of real numbers) are continuous, $f : [0, T] \rightarrow V^*$ with some enlarged space $V^* \supset H$ and $g : [-h, 0] \rightarrow V$ with some dense subspace $V \subset H$. It is assumed that the inclusions $V \subset H \subset V^*$ are continuous and V^* is the dual space of V .

Many authors [2, 8] studied the following delay differential equation:

$$\begin{aligned} \frac{du(t)}{dt} &= A_0 u(t) + A_1 u(t-h) + \int_{-h}^0 a(s) A_2 u(t+s) ds + f(t), \quad \text{a.e. } t \geq 0, \\ u(\theta) &= g(\theta), \quad \theta \in [-h, 0]. \end{aligned} \quad (1.7)$$

The fundamental solution is constructed in Tanabe [8]. In this paper, we establish the existence and norm estimation of solutions for the equation (1.5) by using the fundamental solution.

2. Preliminaries. Let H be a pivot complex Hilbert space and V be a complex Hilbert space such that V is dense in H and the inclusion map $i : V \rightarrow H$ is continuous. The norms of H, V , and the inner product of H are denoted by $|\cdot|$, $\|\cdot\|$, and $\langle \cdot, \cdot \rangle$, respectively. Identifying the antidual of H with H , we may consider that $V \subset H \subset V^*$. The norm of the dual space V^* is denoted by $\|\cdot\|_*$. We consider the following linear functional differential equation on the Hilbert space H .

$$\begin{aligned} \frac{du(t)}{dt} &= A_0 u(t) + A_1 u(t-h) + \int_{-h}^0 a(s) A_2 u(t+s) ds + f(t), \quad \text{a.e. } t \geq 0, \\ u(0) &= g^0, \quad u(s) = g^1(s), \quad \text{a.e. } s \in [-h, 0]. \end{aligned} \quad (2.1)$$

Let $a(u, v)$ be a bounded sesquilinear form defined in $V \times V$ satisfying Gårding's inequality

$$\operatorname{Re} a(u, u) \geq c_0 \|u\|^2 - c_1 |u|^2, \quad (2.2)$$

where $c_0 > 0$ and $c_1 \geq 0$ are real constants. Let A_0 be the operator associated with this sesquilinear form

$$\langle v, A_0 u \rangle = -a(u, v), \quad u, v \in V, \quad (2.3)$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing between V and V^* . The operator A_0 is bounded linear from V into V^* . The realization of A_0 in H , which is the restriction of A_0 to the domain $D(A_0) = \{u \in V : A_0 u \in H\}$, is also denoted by A_0 . It is proved in Tanabe [6] that A_0 generates an analytic semigroup $e^{tA_0} = T(t)$ both in H and V^* and that $T(t) : V^* \rightarrow V$ for each $t > 0$. Throughout this paper, it is assumed that each $A_i (i = 1, 2)$ is bounded and linear from V to V^* (i.e., $A_i \in \mathcal{L}(V, V^*)$) such that A_i maps $D(A_0)$

endowed with the graph norm of A_0 to H continuously. The real valued scalar function $a(s)$ is assumed to be Hölder continuous on $[-h, 0]$. We introduce a Stieltjes measure η given by

$$\eta(s) = -\chi_{(-\infty, -h]}(s)A_1 - \int_s^0 a(\xi) d\xi A_2 : V \rightarrow V^*, \quad s \in [-h, 0], \quad (2.4)$$

where $\chi_{(-\infty, -h]}$ denotes the characteristic function of $(-\infty, -h]$. Then the delay term in (2.1) is written simply as $\int_{-h}^0 d\eta(s)u(t+s)$. The fundamental solution $W(t)$ of (2.1) is defined as a unique solution of

$$W(t) = \begin{cases} T(t) + \int_0^t T(t-s) \int_{-h}^0 d\eta(\xi) W(\xi+s) ds, & t \geq 0, \\ 0, & t < 0, \end{cases} \quad (2.5)$$

and $W(t)$ is constructed by Tanabe [7] under the Hölder continuity of $a(s)$.

THEOREM 2.1 [2]. *The fundamental solution $W(t)$ is strongly continuous in V, H , and V^* , and for each $t > 0$, $W(t) : V^* \rightarrow V$. Furthermore, $W(t)$ satisfies*

$$\frac{d}{dt} W(t) = A_0 W(t) + \int_{-h}^0 d\eta(s) W(t+s), \quad \text{a.e. } t > 0. \quad (2.6)$$

For each $t > 0$, we define the operator valued function $U_t(\cdot)$ by

$$U_t(s) = \int_{-h}^s W(t-s+\xi) d\eta(\xi) : V \rightarrow V, \quad \text{a.e. } s \in [-h, 0]. \quad (2.7)$$

Let $T > 0$ be fixed. Associated with $U_t(\cdot)$, we consider the operator $\mathcal{U} : L^2(-h, 0; V) \rightarrow L^2(0, T; V)$ defined by

$$(\mathcal{U}g^1)(t) = \int_{-h}^0 U_t(s)g^1(s) ds, \quad t \in [0, T] \quad (2.8)$$

for $g^1 \in L^2(-h, 0; V)$.

THEOREM 2.2 [8]. *Let $T > 0$ be fixed. Assume that $f \in L^2(0, T; V^*)$ and $g = (g^0, g^1) \in H \times L^2(-h, 0; V)$. Then there exists a unique solution $u(t) = u(t; f, g)$ of (2.1) on $[0, T]$ satisfying*

$$u \in L^2(0, T; V) \cap W^{1,2}(0, T; V^*) \subset C([0, T]; H). \quad (2.9)$$

Further, for each $T > 0$, there is a constant K_T such that

$$\int_0^T \|u(t)\|^2 dt + \int_0^T \left\| \frac{du(t)}{dt} \right\|_*^2 dt \leq K_T \left(\|g^0\|^2 + \int_{-h}^0 \|g^1(s)\|^2 ds + \int_0^T \|f(t)\|_*^2 dt \right). \quad (2.10)$$

This solution $u(t)$ is represented by

$$u(t; f, g) = W(t)g^0 + (\mathcal{U}g^1)(t) + \int_0^t W(t-s)f(s) ds. \quad (2.11)$$

In what follows, in order to consider the solutions in the state space $C = C([-h, 0]; H)$, we assume that $g = (g^0, g^1)$ is continuous in H , i.e.,

$$g(0) = g^0, \quad g(\cdot) = g^1(\cdot) \in C([-h, 0]; H). \quad (2.12)$$

Let

$$\hat{u}(t; f, g) = \begin{cases} u(t; f, g), & t \in [0, T], \\ g(t), & t \in [-h, 0]. \end{cases} \quad (2.13)$$

Then, by Theorem 2.2, we get

$$\hat{u}(\cdot; f, g) \in C([-h, T]; H) \quad (2.14)$$

if (2.12) is satisfied.

3. Existence and uniqueness of functional integro-differential equations. Using the fundamental solution $W(t)$ in Section 2, we consider the following abstract functional integral equation.

$$\begin{aligned} v(t) &= u(t; f, g) \\ &+ \int_0^t W(t-s) \left[\int_0^s \{k(s, \tau)G(\tau, v_\tau) \right. \\ &\quad \left. + H(s, \tau, v_\tau)\} d\tau + F(s, v_s) \right] ds, \quad 0 < t \leq T, \\ v(\theta) &= g(\theta), \quad \theta \in [-h, 0], \end{aligned} \quad (3.1)$$

where $u(t; f, g)$ is given by (2.11).

We list the following hypotheses.

(A₁) The nonlinear functions $G : [0, T] \times C \rightarrow H$, $H : \Delta_T \times C \rightarrow H$, $F : [0, T] \times C \rightarrow H$, and the kernel $k : \Delta_T \rightarrow R$ (R denotes the set of real numbers) are continuous.

(A₂) Let $b_1, b_3 : [0, T] \rightarrow R$, $b_2 : \Delta_T \rightarrow R^+$ be continuous functions such that

$$\begin{aligned} |G(t, \phi) - G(t, \bar{\phi})|_X &\leq b_1(t) \|\phi - \bar{\phi}\|_C; \\ |H(t, s, \phi) - H(t, s, \bar{\phi})|_X &\leq b_2(t, s) \|\phi - \bar{\phi}\|_C; \\ |F(t, \phi) - F(t, \bar{\phi})|_X &\leq b_3(t) \|\phi - \bar{\phi}\|_C \end{aligned} \quad (3.2)$$

for $t, s \in [0, T]$, $\phi, \bar{\phi} \in C$.

(A₃) The function $k(t, s)$ is Hölder continuous with exponent α , i.e., there exists a positive constant a such that

$$|k(t_1, s_1) - k(t_2, s_2)| \leq a(|t_1 - t_2|^\alpha + |s_1 - s_2|^\alpha) \quad (3.3)$$

for $t_1, t_2, s_1, s_2 \in [0, T]$, $0 < \alpha \leq 1$.

(A₄) For all $0 \leq s \leq t \leq T$,

$$G(t, 0) = 0, \quad H(t, s, 0) = 0, \quad F(t, 0) = 0. \quad (3.4)$$

THEOREM 3.1. Let $f \in L^2(0, T; V^*)$ and $g = (g(0), g(\cdot)) \in H \times L^2(-h, 0; V)$ satisfy (2.12). Assume that the hypotheses (A₁)–(A₄) hold. Then there exists a time $t_1 > 0$ such that the functional integral equation (3.1) admits a unique solution $v(t)$ on $[0, t_1]$.

PROOF. We prove this theorem by using the method of successive approximations.

Set $v^0(t) = u(t; f, g)$, $t \geq 0$. Let $\hat{v}^0(t)$ be the extension of $v^0(t)$ on $[-h, T]$ by (2.13). Then, by the assumptions on f and g , we have $\hat{v}^0(t) \in C([-h, T]; H)$. By hypotheses (A₁)–(A₄), we define $\{\hat{v}^n\}_{n=0}^\infty \subset C([-h, T]; H)$ successively by

$$\begin{aligned}\hat{v}^n(t) &= u(t; f, g) \\ &+ \int_0^t W(t-s) \left[\int_0^s \{k(s, \tau) G(\tau, \hat{v}_\tau^{n-1}) \right. \\ &\quad \left. + H(s, \tau, \hat{v}_\tau^{n-1})\} d\tau + F(s, \hat{v}_s^{n-1}) \right] ds, \quad 0 < t \leq T, \end{aligned} \quad (3.5)$$

$$\hat{v}^n(\theta) = g(\theta), \quad \theta \in [-h, 0]. \quad (3.6)$$

It is obvious that $M = \sup_{t \in [0, T]} \|W(t)\|_{L(H)}$ is finite and that

$$\hat{v}^{n+1}(\theta) - \hat{v}^n(\theta) = 0, \quad \theta \in [-h, 0]. \quad (3.7)$$

For $0 \leq t \leq T$, we have, by (A₁)–(A₄) and the strong continuity of $W(t)$ on $[0, T]$,

$$\begin{aligned} &|\hat{v}^{n+1}(t) - \hat{v}^n(t)| \\ &= \left| \int_0^t W(t-s) \left[\int_0^s \{k(s, \tau) G(\tau, \hat{v}_\tau^n) + H(s, \tau, \hat{v}_\tau^n)\} d\tau + F(s, \hat{v}_s^n) \right] ds \right. \\ &\quad \left. - \int_0^t W(t-s) \left[\int_0^s \{k(s, \tau) G(\tau, \hat{v}_\tau^{n-1}) + H(s, \tau, \hat{v}_\tau^{n-1})\} d\tau + F(s, \hat{v}_s^{n-1}) \right] ds \right| \\ &\leq M \int_0^t \left[\int_0^s |k(s, \tau)| |G(\tau, \hat{v}_\tau^n) - G(\tau, \hat{v}_\tau^{n-1})| d\tau ds \right. \\ &\quad \left. + \int_0^s |H(s, \tau, \hat{v}_\tau^n) - H(s, \tau, \hat{v}_\tau^{n-1})| d\tau \right] + M \int_0^t |F(s, \hat{v}_s^n) - F(s, \hat{v}_s^{n-1})| ds \\ &\leq M \int_0^t \left[\int_0^s \{ |k(s, \tau)| |b_1(\tau)| \|\hat{v}_\tau^n - \hat{v}_\tau^{n-1}\| + |b_2(s, \tau)| \|\hat{v}_\tau^n - \hat{v}_\tau^{n-1}\| \} d\tau \right] ds \\ &\quad + M \int_0^t |b_3(s)| \|\hat{v}_s^n - \hat{v}_s^{n-1}\| ds \\ &\leq M \int_0^t [KL_1 + L_2] \|\hat{v}_\tau^n - \hat{v}_\tau^{n-1}\| s ds + M \int_0^t L_3 \|\hat{v}_s^n - \hat{v}_s^{n-1}\| ds \\ &\leq [M(KL_1 + L_2) \frac{1}{2} t^2 + ML_3 t] \|\hat{v}^n - \hat{v}^{n-1}\|_{C([-h, T]; H)} \\ &= (c_1 t + c_2) t \|\hat{v}^n - \hat{v}^{n-1}\|_{C([-h, T]; H)}, \end{aligned} \quad (3.8)$$

where $c_1 = (1/2)M(KL_1 + L_2)$ and $c_2 = ML_3$. We now choose a sufficiently small constant $t_1 > 0$ such that

$$L = (c_1 t_1 + c_2) t_1 < 1. \quad (3.9)$$

Then by (3.6), (3.8), and (3.9), we get

$$\begin{aligned} \|\hat{v}^{n+1} - \hat{v}^n\|_{C([-h, T]; H)} &\leq L \|\hat{v}^n - \hat{v}^{n-1}\|_{C([-h, T]; H)} \\ &\dots\dots\dots \\ &\leq L^n \|\hat{v}^1 - \hat{v}^0\|_{C([-h, T]; H)}. \end{aligned} \quad (3.10)$$

This implies that $\{\hat{v}^n\}_{n=0}^\infty$ converges uniformly to some $\hat{v} \in C([-h, 0]; H)$. Therefore,

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, t_1]} \|\hat{v}_t^n - \hat{v}_t\|_{C([-h, 0]; H)} = 0. \quad (3.11)$$

Hence, by letting $n \rightarrow \infty$ in (3.5), in view of (A₁)–(A₄) and (3.11), we get

$$\begin{aligned}\hat{v}(t) &= u(t; f, g) \\ &+ \int_0^t \left[W(t-s) \left[\int_0^s \{k(s, \tau)G(\tau, \hat{v}_\tau) \right. \right. \\ &\quad \left. \left. + H(s, \tau, \hat{v}_\tau)\} d\tau \right] + F(s, \hat{v}_s) \right] ds, \quad 0 < t \leq t_1, \\ \hat{v}(\theta) &= g(\theta), \quad \theta \in [-h, 0].\end{aligned}\tag{3.12}$$

This shows the local existence of a solution $v(t) = \hat{v}(t)|_{[0, t_1]}$ of (3.1) on $[0, t_1]$. Let v_1 and v_2 be the solution of (3.1) on $[0, t_1]$. Then it is easy to see, similarly to the above, that

$$\|\hat{v}^1 - \hat{v}^2\|_{C([-h, t_1]; H)} \leq L \|\hat{v}^1 - \hat{v}^2\|_{C([-h, t_1]; H)},\tag{3.13}$$

so that by $L < 1$, $v_1(t) = v_2(t)$ on $[0, t_1]$. This proves the uniqueness. $\square$

Since $k(s, \tau)G(\tau, \hat{v}_\tau)$, $H(s, \tau, \hat{v}_\tau)$, $F(s, v_s) \in L^2(0, t_1; H) \subset L^2(0, t_1; V^*)$, by Theorem 2.1, we see that the solution $v(t)$ of (3.1) satisfies

$$\begin{aligned}\frac{dv(t)}{dt} &= A_0 v(t) + A_1 v(t-h) + \int_{-h}^0 a(s) A_2 v(t+s) ds \\ &+ \int_0^t \{k(t, s)G(s, v_s) + H(t, s, v_s)\} ds \\ &+ F(t, v_t) + f(t), \quad \text{a.e. } t \in [0, t_1], \\ v(\theta) &= g(\theta), \quad \theta \in [-h, 0],\end{aligned}\tag{3.14}$$

and $v \in L^2(0, t_1; V) \cap W^{1,2}(0, t_1; V^*)$. In this sense, we call this v a mild solution of (1.5) on $[0, t_1]$. We give a norm estimation of the mild solution of (1.5) and establish the global existence of solutions with the aid of norm estimations. It is well known (cf. Lions and Magenes [3, Prop. 2.1, Thm. 3.1]) that the inclusion $L^2(0, T; V) \cap W^{1,2}(0, T; V^*) \subset C([0, T]; H)$ is continuous, that is, there exists a constant c_0 such that

$$\|u\|_{C([0, T]; H)} \leq c_0 \left(\|u\|_{L^2(0, T; V)} + \left\| \frac{du}{dt} \right\|_{L^2(0, T; V^*)} \right)\tag{3.15}$$

for all $u \in L^2(0, T; V) \cap W^{1,2}(0, T; V^*)$.

LEMMA 3.1 [5]. *Let $a(t)$, $b(t)$, and $c(t)$ be real valued nonnegative continuous functions defined on R^+ , for which the inequality*

$$c(t) \leq c_0 + \int_0^t a(s)c(s) ds + \int_0^t a(s) \left[\int_0^s b(\tau)c(\tau) d\tau \right] ds\tag{3.16}$$

holds for all $t \in R^+$, where c_0 is a nonnegative constant. Then

$$c(t) \leq c_0 \left(1 + \int_0^t a(s) \exp \left[\int_0^s (a(\tau) + b(\tau)) d\tau \right] ds \right) \quad \text{for all } t \in R^+.\tag{3.17}$$

THEOREM 3.2. Assume that the conditions in Theorem 3.1 hold. Then for any solution $v(t) = v(t; f, g)$ of (3.1) on $[-h, T]$, we have the estimate

$$\|v_t(\cdot; f, g)\|_{C([0, T]; H)} \leq c \left(\|g(0)\| + \|g\|_{L^2(-h, 0; V)} + \|f\|_{L^2(0, T; V^*)} \right) e^{Kt}, \quad (3.18)$$

where c is a positive constant which does not depend on v .

PROOF. From hypotheses (A₁)–(A₄), we have

$$\begin{aligned} & |v(t + \theta; f, g)| \\ & \leq |u(t + \theta; f, g)| + \left| \int_0^{t+\theta} W(t + \theta - s) \right. \\ & \quad \times \left[\int_0^s \{k(s, \tau)G(\tau, v_\tau) + H(s, \tau, v_\tau)\} d\tau + F(s, v_s) \right] ds \Big| \\ & \leq \|u(\cdot; f, g)\|_{C([0, T]; H)} \\ & \quad + M \int_0^{t+\theta} \left[\int_0^s \{K|b_1(\tau)| \|v_\tau\| + |b_2(s, \tau)| \|v_\tau\|\} d\tau + |b_3(s)| \|v_s\| \right] ds. \end{aligned} \quad (3.19)$$

Hence, by (2.10) and (3.15),

$$\begin{aligned} \|v_t(\cdot; f, g)\| &= \sup_{\theta \in [-h, 0]} |v(t + \theta; f, g)| \\ &\leq K_T c_0 \left(\|g(0)\| + \|g\|_{L^2(-h, 0; V)} + \|f\|_{L^2(0, T; V^*)} \right) \\ &\quad + \int_0^t c_1 \|v_s(\cdot; f, g)\| ds + \int_0^t \int_0^s c_2 \|v_\tau(\cdot; f, g)\| d\tau ds \\ &\leq c' \left(\|g(0)\| + \|g\|_{L^2(-h, 0; V)} + \|f\|_{L^2(0, T; V^*)} \right) \\ &\quad + M \left(\int_0^t \|v_s(\cdot; f, g)\| ds + \int_0^t \int_0^s \|v_\tau(\cdot; f, g)\| d\tau ds \right). \end{aligned} \quad (3.20)$$

By using Lemma 3.1, we get

$$\begin{aligned} \|v_t(\cdot; f, g)\|_{C([0, T]; H)} &\leq c \left(\|g(0)\| + \|g\|_{L^2(-h, 0; V)} + \|f\|_{L^2(0, T; V^*)} \right) \\ &\quad \times \left(1 + M \int_0^t \exp \left(\int_0^s (M + 1) d\tau \right) ds \right) \\ &\leq c' \left(\|g(0)\| + \|g\|_{L^2(-h, 0; V)} + \|f\|_{L^2(0, T; V^*)} \right) \\ &\quad \times [1 + M \exp \{(M + 1)T\} t] \\ &\leq c \left(\|g(0)\| + \|g\|_{L^2(-h, 0; V)} + \|f\|_{L^2(0, T; V^*)} \right) e^{Kt} \end{aligned} \quad (3.21)$$

for some constants c and K . This completes the proof. $\square$

By using Theorems 3.1, 3.2, we get the following theorem:

THEOREM 3.3. Assume that the conditions in Theorem 3.1 hold. Then there exists a unique solution $v(t)$ on $[0, T]$ of (3.1) which satisfies the estimate

$$\|v(\cdot; f, g)\|_{C([0, T]; H)} \leq c \left(\|g(0)\| + \|g\|_{L^2(-h, 0; V)} + \|f\|_{L^2(0, T; V^*)} \right) e^{KT} \quad (3.22)$$

for some constants c and K .

ACKNOWLEDGEMENT. This work was supported by KOSEF, 1996.

REFERENCES

- [1] M. B. Dhakne and B. G. Pachpatte, *On some abstract functional integrodifferential equations*, Indian J. Pure Appl. Math. **22** (1991), no. 2, 109–134. MR 92e:47127. Zbl 728.45013.
- [2] J. M. Jeong, S. Nakagiri, and H. Tanabe, *Structural operators and semigroups associated with functional-differential equations in Hilbert spaces*, Osaka J. Math. **30** (1993), no. 3, 365–395. MR 94i:34155. Zbl 818.47039.
- [3] J. L. Lions and E. Magenes, *Non-homogeneous boundary value problems and applications, Vol. I*, Die Grundlehren der mathematischen Wissenschaften, vol. 181, Springer-Verlag, New York, Heidelberg, 1972, Translated from the French by P. Kenneth. MR 50 2670. Zbl 223.35039.
- [4] S. Nakagiri, *Existence, uniqueness and norm estimation for partial functional integrodifferential equations*, Preprint.
- [5] B. G. Pachpatte, *A note on Gronwall-Bellman inequality*, J. Math. Anal. Appl. **44** (1973), 758–762. MR 49 501. Zbl 274.45011.
- [6] H. Tanabe, *On fundamental solution of differential equation with time delay in Banach space*, Proc. Japan Acad. Ser. A Math. Sci. **64** (1988), no. 5, 131–134. MR 89j:34090. Zbl 658.34056.
- [7] ———, *Structural operators for linear delay-differential equations in Hilbert space*, Proc. Japan Acad. Ser. A Math. Sci. **64** (1988), no. 8, 265–266. MR 89k:47061. Zbl 668.45013.
- [8] ———, *Fundamental solutions for linear retarded functional-differential equations in Banach space*, Funkcial. Ekvac. **35** (1992), no. 1, 149–177. MR 93f:34136. Zbl 771.34060.

PARK, Y. LEE, AND J. LEE: DEPARTMENT OF MATHEMATICS, PUSAN NATIONAL UNIVERSITY, PUSAN 609-735, KOREA