RESEARCH NOTES

ON A DENSITY PROBLEM OF ERDÖS

SAFWAN AKBİK

(Received 13 April 1998 and in revised form 10 June 1998)

ABSTRACT. For a positive integer n , let $P(n)$ denotes the largest prime divisor of n and define the set: $\mathcal{S}(x) = \mathcal{S} = \{n \leq x : n \text{ does not divide } P(n)!\}$. Paul Erdős has proposed that $|\mathcal{S}| = o(x)$ as $x \rightarrow \infty$, where $|\mathcal{S}|$ is the number of $n \in \mathcal{S}$. This was proved by Ilias Kastanas. In this paper we will show the stronger result that $|\mathcal{S}| = O(xe^{-1/4}\sqrt{\log x})$.

Keywords and phrases. Number theory, primes, factorial, density, divisibility.

1991 Mathematics Subject Classification. 11B05, 11N25.

Introduction. For a positive integer n , let $P(n)$ denote the largest prime divisor of n and define the set

$$\mathcal{S}(x) = \mathcal{S} = \{n \leq x : n \text{ does not divide } P(n)!\}. \quad (1)$$

Paul Erdős [1] proposed that $|\mathcal{S}| = o(x)$ as $x \rightarrow \infty$, where $|\mathcal{S}|$ is the number of $n \in \mathcal{S}$. A solution [3] was provided by Ilias Kastanas. There was also [3] a claim of proving that $|\mathcal{S}(x)| = O(x/\log x)$. In this paper, we show the stronger result.

THEOREM. *For some constant $a > 0$, we have*

$$|\mathcal{S}| = O\left(xe^{-a\sqrt{\log x}}\right). \quad (2)$$

In fact, $a = 1/4$ suffices.

LEMMA 1. *Let $\nu(n)$ be the number of distinct prime divisors of n . Define*

$$S_1 = \{n \leq x : \nu(n) > 4k \log \log x, k \geq 1\}. \quad (3)$$

Then

$$|S_1| = O\left(\frac{x}{(\log x)^k}\right) \quad (4)$$

uniformly in k .

PROOF. It is well known [2] that if $d(m)$ is the number of divisors of m , then $\sum_{m \leq x} d(m) = O(x \log x)$. Since $d(m) \geq 2^{\nu(m)}$,

$$O(x \log x) \geq \sum_{m \leq x} 2^{\nu(m)} \geq \sum_{m \in S_1} 2^{\nu(m)} \geq \sum_{m \in S_1} (\log x)^{(4 \log 2)k} \geq |S_1| (\log x)^{k+1}, \quad (5)$$

and the lemma follows. □

LEMMA 2. Let $C(x) = C = (\log x)^k$, where $k = k(x)$ will be chosen later. Define

$$S_2 = \{n \leq x : p^2 \mid n \text{ for some prime } p > C\}. \quad (6)$$

Then

$$|S_2| = O\left(\frac{x}{(\log x)^k}\right). \quad (7)$$

PROOF. Since $n \in S_2$ if and only if $n = tp^2$ for some $C < p \leq \sqrt{x}$ and some $t \leq x/p^2$,

$$|S_2| = \sum_{C < p \leq \sqrt{x}} \left\lfloor \frac{x}{p^2} \right\rfloor \leq \sum_{C < p \leq \sqrt{x}} \frac{x}{p^2} = O\left(\frac{x}{C}\right) = O\left(\frac{x}{(\log x)^k}\right). \quad (8)$$

The first big O in (8) follows since $\sum_{p > C} 1/p^2 \leq \int_{[C]}^{\infty} du/u^2 = 1/[C] = O(1/C)$, for $C \geq 1$. $\square$

LEMMA 3. Let

$$S_3 = \{n \leq x : p^\alpha \mid n \text{ for some } \alpha \geq T \text{ and some prime } P \leq C\}, \quad (9)$$

where $T = 2 \log C$. Then

$$|S_3| = O\left(\frac{x}{(\log x)^k}\right). \quad (10)$$

PROOF.

$$\begin{aligned} |S_3| &= \sum_{\substack{p \leq C \\ \alpha \geq T}} \left\lfloor \frac{x}{p^\alpha} \right\rfloor \leq 2 \sum_{p \leq C} \frac{x}{p^T} \leq 2 \left(\frac{x}{2^T}\right) \sum_p \frac{4}{p^2} = O\left(\frac{x}{2^T}\right) \\ &= O\left(\frac{x}{C^{2 \log 2}}\right) \leq O\left(\frac{x}{C}\right) = O\left(\frac{x}{(\log x)^k}\right). \end{aligned} \quad (11)$$

The first inequality in (11) is valid because $\sum_{\alpha \geq T} 1/p^\alpha \leq 1/p^T + 1/p^{T+1} + \dots \leq 2/p^T$ and the second is valid because $p^T = 2^T (p/2)^T \geq 2^T (p/2)^2$ for $T \geq 2$. $\square$

LEMMA 4. Let $S'(x) = S - (S_1 \cup S_2 \cup S_3)$, then, for any $n \in S'$, we have

$$P(n) \leq 2CT. \quad (12)$$

PROOF. Let $n \in S'$. Then $n \in S$ and so n does not divide $P(n)!$. There exists a prime p_0 dividing n such that

$$v_{p_0}(n) > v_{p_0}(P(n)!), \quad (13)$$

where $v_p(m)$ denotes the largest integer t such that p^t divides m . Since $v_{p_0}(P(n)!) \geq 1$, (13) implies that $v_{p_0}(n) \geq 2$. Since $n \notin S_2$, it follows that $p_0 \leq C$. Also $n \notin S_3$, so that $T \geq v_{p_0}(n)$. Hence,

$$T \geq v_{p_0}(n) > v_{p_0}(P(n)!) \geq \frac{P(n)}{2p_0} \geq \frac{P(n)}{2C} \quad (14)$$

which implies (12). Note that the third inequality in (14) is true because $v_p(m!) \geq m/2p$ for any m and any $p \mid m$. This is because

$$v_p(m!) \geq \left\lfloor \frac{m}{p} \right\rfloor > \frac{m}{p} - 1 \geq \frac{m}{2p}, \quad p \neq m, \quad (15)$$

and $[m/p] = 1 \geq m/2p$ if $p = m$. $\square$

PROOF OF THE THEOREM. For $n \in S'$, we have $n \notin S_1 \cup S_2 \cup S_3$. Thus, $v_p(n) \leq 4k \log \log x$. Also, if p^α is any prime power dividing n , then one of the following two possibilities must occur:

- (a) $p \leq C$ and $\alpha \leq T$,
- (b) $p > C$ and $\alpha = 0$ or 1 .

Case (a) generates at most $C(T+1) \leq 2CT$ prime powers. For Case (b), the number of prime powers p^α with $p > C$ and $\alpha \leq 1$ is at most $P(n)$. By Lemma 4, this is at most $2CT$. Hence, the number of possible prime powers p^α that divide an $n \in S'$ is at most $4CT$. But such an n can be the product of at most $4k \log \log x$ distinct prime powers. Therefore,

$$\begin{aligned} |S'| &\leq (4CT)^{4k \log \log x} = (8C \log C)^{4k \log \log x} \\ &= e^{4k \log \log x (\log 8 + k \log \log x + \log k + \log \log \log x)} \leq e^{8k^2 (\log \log x)^2} \end{aligned} \quad (16)$$

since $\log 8 + \log(k \log \log x) \leq k \log \log x$, for $k \log \log x \geq 4$.

Choosing

$$k = \frac{1}{4} \cdot \frac{\sqrt{\log x}}{\log \log x}, \quad (17)$$

(16) gives $|S'| \leq e^{(1/2) \log x} = x^{1/2}$. Hence,

$$S' = O\left(x e^{-(1/4) \sqrt{\log x}}\right). \quad (18)$$

From (17), we have $x/(\log x)^k = x e^{-1/4 \sqrt{\log x}}$. Lemmas 1, 2, and 3 imply that

$$|S_i| = O\left(x e^{-1/4 \sqrt{\log x}}\right), \quad i = 1, 2, 3. \quad (19)$$

Finally, $S = S' \cup [S \cap (S_1 \cup S_2 \cup S_3)]$. Hence, (18) and (19) yield

$$|S| \leq |S'| + |S_1| + |S_2| + |S_3| = O\left(x e^{-(1/4) \sqrt{\log x}}\right), \quad (20)$$

and (2) follows with $a = 1/4$. $\square$

REMARK. If $\pi(x)$ is the number of prime integers that are less than or equal to x , an early version of the prime numbers theorem asserts that

$$\pi(x) = \int_2^x \frac{du}{\log u} + O\left(x e^{-a \sqrt{\log x}}\right), \quad (21)$$

for some constant a . Although the big O terms in (19) and (2) are similar, there is no apparent relationship between the PNT and (2).

REFERENCES

- [1] P. Erdős, *A Proposed Problem*, Amer. Math. Monthly **98** (1991), 965.
- [2] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 5th ed., The Clarendon Press, Oxford University Press, New York, 1979. MR 81i:10002. Zbl 423.10001.
- [3] I. Kostas, *The smallest factorial that is a multiple of n* , Amer. Math. Monthly **101** (1994), 179.

AKBİK: DEPARTMENT OF MATHEMATICS, HOFSTRA UNIVERSITY, HEMPSTEAD, NY 11550, USA