ON A FOUR-GENERATOR COXETER GROUP

MUHAMMAD A. ALBAR

(Received 4 December 1999)

ABSTRACT. We study one of the 4-generator Coxeter groups and show that it is SQ-universal (SQU). We also study some other properties of the group.

Keywords and phrases. Group presentation, Reidemeister-Schreier process, Coxeter group, SQ-universality.

2000 Mathematics Subject Classification. Primary 20F05.

1. Introduction. We consider the Coxeter group P given by the presentation

$$\begin{aligned} P = \langle x_1, x_2, x_3, x_4 \mid x_1^2 = x_2^2 = x_3^2 = x_4^2 = (x_1 x_2)^3 \\ = (x_2 x_3)^3 = (x_1 x_3)^3 = (x_1, x_4)^3 = (x_3 x_4)^3 = (x_2 x_3)^3 = e \rangle. \end{aligned} \quad (1.1)$$

The Coxeter graph of this group is clearly just a combinatorial tetrahedron:

We observe that each face is the graph of the Euclidean triangle group $\Delta(3, 3, 3)$ which is an affine Weyl group and this contains a nilpotent subgroup of finite index. The group P is infinite and it will be interesting to see its largeness by answering whether it is SQ-universal or not.

2. SQ-universality. We let S_3 be the symmetric group of degree 3. Thus

$$S_3 = \langle y_1, y_2 \mid y_1^2 = y_2^2 = (y_1 y_2)^3 = e \rangle. \quad (2.1)$$

We consider the map $\theta : P \rightarrow S_3$ defined by

$$\theta(x_1) = y_1, \quad \theta(x_2) = y_2, \quad \theta(x_3) = \theta(x_4) = y_1 y_2 y_1. \quad (2.2)$$

It is easy to see that θ is an epimorphism and $P/\ker \theta \cong S_3$. A Schreier transversal for S_3 in P is $\{e, x_1, x_2, x_1 x_2, x_2 x_1, x_1 x_2 x_1\}$. A straightforward application of the

Reidemeister-Schreier process gives the following presentation for $\ker \theta$:

$$\ker \theta = \langle a, b, c, d \mid (ad)^3 = (bc)^3 = (abcd)^3 = e \rangle. \quad (2.3)$$

Letting $a = d^{-1}$ and $b = c^{-1}$, we see that $\ker \theta$ is mapped homomorphically onto the free group of rank 2, F_2 . Hence $\ker \theta$ is SQU. Since the index of $\ker \theta$ in P is finite (6), we get that P is also SQU [4].

3. The growth series. Let (P, X) be a Coxeter system and let $Y \subseteq X$. We denote the subgroup of P , generated by Y , by P_Y . Then (W_Y, Y) is also a Coxeter system. In Bourbaki [2, Section 1 of Chapter 4], Exercise 26 gives the following formula for computing the growth series of P (word growth in the sense of Milner and Gromov):

$$\sum_{Y \subseteq X} \frac{(-1)^{|Y|}}{P_Y(t)} = \begin{cases} \frac{t^m}{P(t)} & \text{if } P \text{ is finite,} \\ 0 & \text{if } P \text{ is infinite.} \end{cases} \quad (3.1)$$

In the formula, $G(t)$ is the growth series of G , m is the length of the unique element of P of maximal length.

We use (3.1) to compute $P(t)$. We compute $P(t)$ in steps corresponding to the cardinality of Y :

$|Y| = 0$ is the trivial subgroup with growth series $y_0 = 1$.

$|Y| = 1$ four cyclic subgroups of order 2 with growth series $y_1 = 1 + t$.

$|Y| = 2$ six dihedral subgroups of order 6 with growth series $y_2 = (1 + t)(1 + t + t^2)$.

$|Y| = 3$ four affine subgroups with growth series given by $1/y_0 - 3/y_1 + 3/y_2 - 1/y_3 = 0$, that is, $y_3 = (1 + t + t^2)/(1 - t)^2$.

$|Y| = 4$ the whole group with growth $y_4(t) = P(t)$ given by $1/y_0 - 4/y_1 + 6/y_2 - 4/y_3 + 1/y_4 = 0$, that is, $y_4 = (1 + t)(1 + t + t^2)/(1 - t)(1 - t - 3t^2)$.

The growth coefficients $\{c_n\}$ are given by the linear recurrence $c_0 = 1$, $c_1 = 4$, $c_2 = 12$, $c_3 = 30$, $c_n = 2c_{n-1} + 2c_{n-2} - 3c_{n-3}$, $n \geq 4$ (see [3]). We observe from the growth series y_4 that zeros of the denominator are not on the unit circle. This implies that P has no nilpotent subgroup of finite index—in accordance with the fact that P is SQU.

It is possible to show that the group P and the Geisking group $G = \langle x, y \mid x^2 y^2 = xy \rangle$ are isometric and hence y_4 is also the growth series of G (see [3]). In [1], it appears that the two Coxeter groups T_n and S_n are also isometric and so have the same growth series.

4. The commutator subgroup. Using the Reidemeister-Schreier process, we get the following presentation for P' :

$$P' = \langle x, y, z \mid x^3 = y^3 = z^3 = (xy)^3 = (xz)^3 = (yz)^3 = e \rangle. \quad (4.1)$$

We use P' to show that P is SQU in a different method. Let K be the normal closure of the elements $xy^{-1}, xz^{-1}, yz^{-1}$ in P' . The group K has index 3 in P' . Using the Reidemeister-Schreier process, we get the following presentation for K :

$$\begin{aligned} K &= \langle u_1, u_2, u_3, v_1, v_2, v_3 \mid v_1^2 = v_2^2 = v_3^2 = u_1 u_2 u_3 \\ &= u_1 u_3 u_2 = v_1 v_2 v_3 = u_1 v_2 u_3 v_1 u_2 u_3 = e \rangle. \end{aligned} \quad (4.2)$$

Letting $u_3 = v_3 = e$, we see that K is mapped homomorphically onto $Z * Z_3$. Since $Z * Z_3$ is SQU (see [4]), therefore K is SQU. Since K is of finite index in P' and P' is of finite index in P , we get that P is SQU.

ACKNOWLEDGEMENT. The author thanks King Fahd University of Petroleum and Minerals for supporting him in his research.

REFERENCES

- [1] M. A. Albar, M. A. Al-Hamed, and N. A. Al-Saleh, *The growth of Coxeter groups*, Math. Japon. **47** (1998), no. 3, 417–428. MR 99f:20066. Zbl 912.20032.
- [2] N. Bourbaki, *Éléments de Mathématique. Fasc. XXXIV. Groupes et Algèbres de Lie. Chapitre IV: Groupes de Coxeter et Systèmes de Tits. Chapitre V: Groupes Engendrés par des Réflexions. Chapitre VI: Systèmes de Racines*, Actualités Scientifiques et Industrielles, no. 1337, Hermann, Paris, 1968, 288 pp. MR 39#1590. Zbl 186.33001.
- [3] D. L. Johnson and H.-J. Song, *The growth series of the Giesecking group*, Discrete Groups and Geometry (Birmingham, 1991), pp. 120–124, London Math. Soc. Lecture Note Ser., no. 173, Cambridge Univ. Press, Cambridge, 1992. MR 93m:20049. Zbl 769.57003.
- [4] P. M. Neumann, *The SQ-universality of some finitely presented groups*. Collection of articles dedicated to the memory of Hanna Neumann, I., J. Austral. Math. Soc. **16** (1973), 1–6. MR 48#11342. Zbl 267.20026.

MUHAMMAD A. ALBAR: DEPARTMENT OF MATHEMATICAL SCIENCES, KING FAHD UNIVERSITY OF PETROLEUM AND MINERALS, DHAHRAN 31261, SAUDI ARABIA