A NOTE ON (gDF) -SPACES

RENATA R. DEL-VECCHIO, DINAMÉRICO P. POMBO, JR., and
CYBELE T. M. VINAGRE

(Received 20 March 1998 and in revised form 24 July 1998)

ABSTRACT. Certain locally convex spaces of scalar-valued mappings are shown to be finite-dimensional.

Keywords and phrases. (gDF) -spaces, (DF) -spaces, finite-dimensional spaces.

2000 Mathematics Subject Classification. Primary 46A04.

1. Introduction. Radenović [6], generalizing a result of Iyahen [2], has shown that if E is a Banach space and $(E, \sigma(E, E'))$ (or $(E', \sigma(E', E))$) is a (DF) -space [1], then E is finite-dimensional. His result has been extended to arbitrary locally convex spaces by Krassowska and Šliwa [3].

In [4, 5], (DF) -spaces have been generalized as follows: a locally convex space (E, τ) is a (gDF) -space if

(a) (E, τ) has a fundamental sequence $(B_n)_{n \in \mathbb{N}}$ of bounded sets, and

(b) τ is the finest locally convex topology on E that agrees with τ on each B_n .

In this note, we prove that if an arbitrary vector space of scalar-valued mappings is a (gDF) -space under the locally convex topology of pointwise convergence, then it is finite-dimensional. As a consequence, the above-mentioned theorem of Krassowska and Šliwa readily follows.

2. The result. Throughout this note, all vector spaces under consideration are vector spaces over a field $\mathbb{K}$ which is either $\mathbb{R}$ or $\mathbb{C}$. In our result, E denotes an arbitrary set and H denotes a subspace of the vector space of all mappings from E into $\mathbb{K}$. We consider on H the separated locally convex topology of pointwise convergence and represent by H' the topological dual of H .

THEOREM 2.1. *The following conditions are equivalent:*

- (a) H is a finite-dimensional vector space;
- (b) H is a (DF) -space;
- (c) H is a (gDF) -space.

PROOF. It is clear that (a) implies (b) and (b) implies (c) (every (DF) -space is a (gDF) -space).

Suppose that condition (c) holds. If H is infinite-dimensional, there exists a countable linearly independent subset $\{\varphi_n; n \in \mathbb{N}\}$ of H' . Let $(B_n)_{n \in \mathbb{N}}$ be an increasing fundamental sequence of bounded subsets of H . Then, $(B_n^0)_{n \in \mathbb{N}}$ is a decreasing sequence of neighborhoods of zero in $(H', \beta(H', H))$ forming a fundamental system

of neighborhoods of zero in $(H', \beta(H', H))$. For each $n \in \mathbb{N}$, fix an $\alpha_n > 0$ such that $\alpha_n \varphi_n \in B_n^0$; then $(\alpha_n \varphi_n)_{n \in \mathbb{N}}$ converges to zero in $(H', \beta(H', H))$. By [5, Theorem 1.1.7], the set $\Gamma = \{\alpha_n \varphi_n; n \in \mathbb{N}\}$ is equicontinuous. Hence, there exist $x_1, \dots, x_m \in E$ and there exists an $\alpha > 0$ such that the relations

$$f \in H, \quad |f(x_1)| \leq \alpha, \dots, |f(x_m)| \leq \alpha, \quad \varphi \in \Gamma \quad (2.1)$$

imply

$$|\varphi(f)| \leq 1. \quad (2.2)$$

For each $i = 1, \dots, m$, let $\delta_i \in H'$ be given by $\delta_i(f) = f(x_i)$ for $f \in H$, and put $F = \{\delta_1, \dots, \delta_m\}$. We claim that $\Gamma \subset [F]$, where $[F]$ is the finite-dimensional vector space generated by F . Indeed, let $\varphi \in \Gamma$ and take an $f \in H$ such that $\delta_1(f) = \dots = \delta_m(f) = 0$. Then, for all $\lambda \in \mathbb{K}$,

$$|(\lambda f)(x_1)| = |\delta_1(\lambda f)| = 0 \leq \alpha, \dots, |(\lambda f)(x_m)| = |\delta_m(\lambda f)| = 0 \leq \alpha. \quad (2.3)$$

Consequently, $|\varphi(\lambda f)| = |\lambda| |\varphi(f)| \leq 1$. By the arbitrariness of λ , $\varphi(f) = 0$. By [7, Lemma 5, Chapter II], $\varphi \in [F]$. Therefore the vector space generated by the set $\{\varphi_n; n \in \mathbb{N}\}$ is finite-dimensional, which contradicts the choice of $(\varphi_n)_{n \in \mathbb{N}}$. This completes the proof of the theorem. $\square$

REMARK 2.2. The theorem of Krassowska and Šliwa mentioned at the beginning of this note follows from Theorem 2.1. In fact, let E be a separated locally convex space. If $(E', \sigma(E', E))$ is a (DF) -space, then E' is finite-dimensional by Theorem 2.1, and so E is finite-dimensional. Hence, E is finite-dimensional if $(E, \sigma(E, E'))$ is a (DF) -space.

REFERENCES

- [1] A. Grothendieck, *Sur les espaces (F) et (DF)*, Summa Brasil. Math. **3** (1954), 57–123. MR 17,765b. Zbl 058.09803.
- [2] S. O. Iyahen, *Some remarks on countably barrelled and countably quasibarrelled spaces*, Proc. Edinburgh Math. Soc. (2) **15** (1966/1967), 295–296. MR 37#1947. Zbl 157.20203.
- [3] D. Krassowska and W. Šliwa, *When $(E, \sigma(E, E'))$ is a DF -space?*, Comment. Math. Univ. Carolin. **33** (1992), no. 1, 43–44. MR 93f:46001. Zbl 782.46006.
- [4] K. Noureddine, *Espaces du type D_b* , C. R. Acad. Sci. Paris Sér. A-B **276** (1973), A1301–A1303. MR 51#11048. Zbl 257.46003.
- [5] ———, *Nouvelles classes d'espaces localement convexes*, Publ. Dép. Math. (Lyon) **10** (1973), no. 3, 259–277. MR 51#3847. Zbl 292.46002.
- [6] S. Radenović, *Some remarks on the weak topology of locally convex spaces*, Publ. Inst. Math. (Beograd) (N.S.) **44(58)** (1988), 155–157. MR 90d:46006. Zbl 684.46003.
- [7] A. P. Robertson and W. Robertson, *Topological Vector Spaces*, Cambridge University Press, London, 1973. MR 50#2854. Zbl 251.46002.

DEL-VECCHIO, POMBO, AND VINAGRE: INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL FLUMINENSE, RUA MÁRIO SANTOS BRAGA, S/Nº, 24020-140 NITERÓI, RJ, BRASIL