CONVEX ISOMETRIC FOLDING

E. M. ELKHOLY

(Received 27 September 1993 and in revised form 5 June 1996)

ABSTRACT. We introduce a new type of isometric folding called “convex isometric folding.” We prove that the infimum of the ratio $\text{Vol} N / \text{Vol} \varphi(N)$ over all convex isometric foldings $\varphi : N \rightarrow N$, where N is a compact 2-manifold (orientable or not), is $1/4$.

Keywords and phrases. Isometric and convex foldings, regular and universal covering spaces, manifolds, group of covering transformations, fundamental regions.

2000 Mathematics Subject Classification. Primary 51H10, 57N10.

1. Introduction. A map $\varphi : M \rightarrow N$, where M and N are C^∞ Riemannian manifolds of dimensions m and n , respectively, is said to be an isometric folding of M into N if and only if for any piecewise geodesic path $\gamma : J \rightarrow M$, the induced path $\varphi \circ \gamma : J \rightarrow N$ is a piecewise geodesic and of the same length. The definition is given by Robertson [4]. The set of all isometric foldings $\varphi : M \rightarrow N$ is denoted by $\mathcal{F}(M, N)$.

Let $p : M \rightarrow N$ be a regular locally isometric covering and let G be the group of covering transformations of p . An isometric folding $\varphi \in \mathcal{F}(M)$ is said to be p -invariant if and only if for all $g \in G$ and all $x \in X$, $p(\varphi(x)) = p(\varphi(g, x))$. See Robertson and Elkholy [5]. The set of p -invariant isometric foldings is denoted by $\mathcal{F}_i(M, p)$.

DEFINITION 1.1. Let $\varphi \in \mathcal{F}(M, N)$, where M and N are C^∞ Riemannian manifolds of dimensions m and n , respectively. We say that φ is a convex isometric folding if and only if $\varphi(M)$ can be embedded as a convex set in $\mathbb{R}^n$.

We denote the set of all convex isometric foldings of M into N by $C(M, N)$, and if $C(M, N) \neq \emptyset$, then it forms a subsemigroup of $\mathcal{F}(M, N)$.

DEFINITION 1.2. We say that $\varphi \in \mathcal{F}_i(M, p)$ is a p -invariant convex isometric folding if and only if $\varphi(M)$ can be embedded as a convex set in $\mathbb{R}^m$.

We denote the set of p -invariant convex isometric foldings of M by $C_i(M, p)$. If $C_i(M, p) \neq \emptyset$, then for any covering map, $p : M \rightarrow N$, $C_i(M, p)$ is a subsemigroup of $C(M)$.

To solve our main problem we need the following:

(1) Robertson and Elkholy [5] proved that if N is an n -smooth Riemannian manifold, $p : M \rightarrow N$ is its universal covering, and G is the group of covering transformations of p , then $\mathcal{F}(N)$ is isomorphic as a semigroup to $\mathcal{F}_i(M, p)/G$.

(2) Elkholy [1] proved that if N is an n -smooth Riemannian manifold, $p : M \rightarrow N$ is its universal covering, and $\varphi \in \mathcal{F}(N)$ such that $\varphi_* : \pi_1(N) \rightarrow \pi_1(N)$ is trivial, then the

corresponding folding $\psi \in \mathcal{F}_i(M, p)$ maps each fiber of p to a single point.

(3) Elkholy and Al-Ahmady [3] proved that under the same conditions of (2), if N is a compact 2-manifold, then

$$\frac{\text{Vol } N}{\text{Vol } \varphi(N)} = \frac{\text{Vol } F}{\text{Vol } \psi(F)}, \quad (1.1)$$

where F is a fundamental region of G in M .

2. Convex isometric folding and covering spaces. The next theorem establishes the relation between the set of convex isometric folding of a manifold, $C(N)$, and the set of p -invariant convex isometric folding of its universal covering space, $C_i(M, p)$.

THEOREM 2.1. *Let N be a manifold and $p : M \rightarrow N$ its universal covering. Let G be the group of covering transformations of p . If $C(N) \neq \emptyset$, then $C(N)$ is isometric as a semigroup to $C_i(M, p)/G$.*

PROOF. Let $C(N) \neq \emptyset$. Then by using (1), there exists an isomorphism f from $\mathcal{F}_i(M, p)/G$ into $\mathcal{F}(N)$. Since $C_i(M, p)$ is a subsemigroup of $\mathcal{F}_i(M, p)$, $C_i(M, p)/G$ is a subsemigroup of $\mathcal{F}_i(M, p)/G$.

Let $h = f \mid (C_i(M, p)/G)$. Since $C_i(M, p)/G$ is a semigroup, h is a homeomorphism and also it is one-one. To show that h is an onto map, we suppose that $\varphi \in C(N)$. Hence, $\varphi \in \mathcal{F}(N)$ and, consequently, there exists $\psi \in \mathcal{F}_i(M, p)/G$. Since $\varphi \in C(N)$, φ_* is trivial and hence for all $x \in M$, $\psi(G \cdot x) = \psi(x)$, and therefore $\psi \in C_i(M, p)/G$. $\square$

THEOREM 2.2. *Let N be a compact orientable 2-manifold and consider the universal covering space $(\mathbb{R}^2, P)$ of N . Let $\varphi \in C(N)$ and $\psi \in C_i(\mathbb{R}^2, p)$. Then for all $x, y \in \mathbb{R}^2$, $d(\psi(x), \psi(y)) \leq \Delta$, where Δ is the radius of a fundamental region for the covering space.*

PROOF. Elkholy [1] proved the truth of the theorem for $N = S^2$. So, we have to prove it for the connected sum of n -tori. First, let $N = T$ be a torus homomorphic to the quotient space obtained by identifying opposite sides of a square of length “ a ” as shown in Figure 1(a)

FIGURE 1.

Suppose that $\varphi : T \rightarrow T$ is a convex isometric folding. Then $\varphi_*(\pi_1(T))$ is trivial. By Theorem 2.1, there exists a convex isometric folding $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that for all $x, y \in \mathbb{R}^2$ and for all $g \in G$, $p(\psi(x)) = p(\psi(g \cdot x))$. Equivalently, for all $(P, Q) \in \mathbb{R}^2$ and for all $g \in \mathbb{Z} \times \mathbb{Z}$, there exists a unique $h \in \mathbb{Z} \times \mathbb{Z}$ such that $h \circ \psi(P, Q) = \psi(g(P, Q))$, i.e.,

$$\psi(P, Q) + (\sqrt{2}\Delta m, \sqrt{2}\Delta n') = \psi(P + \sqrt{2}\Delta m, Q + \sqrt{2}\Delta n), \quad \text{where } m, n, m', n' \in \mathbb{Z}. \quad (2.1)$$

Consider any fundamental region F of the covering space $(\mathbb{R}^2, p)$ of T , i.e., a closed square of length “ a ” with sides identified as shown in Figure 1(b). Since φ_* is trivial, by (2), for all $x \in \mathbb{R}^2$, $\psi(G \cdot x) = \psi(x)$. Now, let x and y be distinct points of $\mathbb{R}^2$ such that $x = g \cdot y$ for all $g \in G$ and let $d(x, y) = \alpha_1$. Then there exists a point $x^* = g \cdot x$ such that

$$d(y, x^*) = \min(\alpha_i), \quad \alpha_i = d(y, g_i \cdot x), \quad i = 1, \dots, 4. \quad (2.2)$$

Thus, there are always four equivalent points $g_i \cdot x$, $i = 1, \dots, 4$ which form the vertices of a square of length “ a ” and such that $d(g_i \cdot x, y) \leq 2\Delta$. From Figure 1(b), it is clear that $\max d(x^*, y) \leq \Delta$ and since ψ is an isometric folding, by Robertson [4], $d(\psi(x), \psi(y)) \leq d(x, y)$, i.e.,

$$d(\psi(x), \psi(y)) = d(\psi(g_i \cdot x), \psi(y)) \leq d(g_i \cdot x, y) = d(x, y) \leq \Delta, \quad (2.3)$$

and this proves the theorem for $N = T$.

Now, consider the connected sum of two tori, obtained as a quotient space of an octagon with sides identified as shown in Figure 2(a). The group of covering transformations G is isometric to $\mathbb{Z} \times \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}$. Using the same previous technique, we can

FIGURE 2.

obtain four equivalent points as the vertices of a square of diameter 2Δ such that $\max d(y, x^*) \leq \Delta$, and the result follows. This theorem, by using the above method, is true for the connected sum of n -tori. $\square$

THEOREM 2.3. *Let N be a compact nonorientable 2-manifold and consider the universal covering space (M, p) of N . Let $\phi \in C(N)$ and $\psi \in C_i(M, p)$. Then for all $x, y \in M$, $d(\psi(x), \psi(y)) \leq \Delta$, where Δ is the radius of a fundamental region for the covering space.*

PROOF. By Elkholy [2], the theorem is true for $N = p^2$ and $M = S^2$. Now, consider the connected sum of two projective planes, the Klein bottle K , homeomorphic to the quotient space obtained by identifying the opposite sides of a square as shown in Figure 3(a).

FIGURE 3.

Suppose that $\varphi : K \rightarrow K$ is a convex isometric folding. Then there exists a convex isometric folding $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that for all $x \in \mathbb{R}^2$ and $g \in G$, $p(\psi(x)) = p(\psi(g \cdot x))$. Equivalently, for all $(P, Q) \in \mathbb{R}^2$ and for all $g \in \mathbb{Z} \times \mathbb{Z}_2$, there exists a unique $h \in \mathbb{Z} \times \mathbb{Z}_2$ such that $h \circ \psi(P, Q) = \psi(g(P, Q))$, i.e.,

$$\begin{aligned} \psi(P, Q) + (\sqrt{2}\Delta m', \sqrt{2}\Delta n') \\ = \psi(P + \sqrt{2}\Delta m, \sqrt{2}\Delta n + (-)^m Q), \quad \text{where } m, n, m', n' \in \mathbb{Z}. \end{aligned} \quad (2.4)$$

Any fundamental region F of the covering space $(\mathbb{R}^2, p)$ of K is a closed square of diameter 2Δ with the boundary identified as shown in Figure 3(b). Since φ_* is trivial, for all $x \in \mathbb{R}^2$, $\psi(G \cdot x) = \psi(x)$.

Now, let x and y be distinct points of $\mathbb{R}^2$ such that $y \neq g \cdot x$ for all $g \in G$, and let $d(x, y) = \alpha_1$. Thus, there exists a point $x^* = g \cdot x$ such that

$$d(y, x^*) = \min(\alpha_i), \quad \alpha_i = d(y, g_i \cdot x), \quad i = 1, \dots, 4. \quad (2.5)$$

Thus, there are always four equivalent points $g_i \cdot x$ which form the vertices of a parallelogram such that the shortest diameter is of length less than 2Δ .

Now, the point y is either inside or on the boundary of a triangle of vertices $g_1 \cdot x = x, g_2 \cdot x, g_3 \cdot x$. Let y' be a point equidistant from the vertices of this triangle, i.e.,

$$d(y', x) = d(y', g_2 \cdot x) = d(y', g_3 \cdot x). \quad (2.6)$$

From Figure 3(b), it is clear that $d(y', x) < \Delta$ and, hence, $d(x^*, y) < \Delta$. Therefore,

$$d(\psi(x), \psi(y)) = d(\psi(g_i \cdot x), \psi(y)) \leq d(g \cdot x_i, y) = d(x^*, y) < \Delta \quad (2.7)$$

and the result follows.

Now, let N be the connected sum of three projective planes obtained as the quotient space of a hexagon with the sides identified in pairs as indicated in Figure 4(a). In this case, $(\mathbb{R}^2, p)$ is the universal cover of N and $G \simeq \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}_2$. Using the same method as that used above, we can always have equivalent points $g_i \cdot x$, $i = 1, \dots, 4$ which form the vertices of a parallelogram whose shortest diameter is of length less than 2Δ . From Figure 4(b), we can see that $\max d(y, x^*) < \Delta$ and the theorem is proved.

In general and by using the same technique, the theorem is also true for the connected sum of n -projective planes. $\square$

FIGURE 4.

3. Volume and convex folding. The following theorem succeeds in estimating the maximum volume we may have if we convexly folded a compact 2-manifold into itself.

THEOREM 3.1. *The infimum of the ratio*

$$e_N = \frac{\text{Vol } N}{\text{Vol } \varphi(N)}, \quad (3.1)$$

where N is a compact 2-manifold over all convex isometric foldings $\varphi \in C(N)$ of degree zero, is 4.

PROOF. Robertson [4] has shown that if N is a compact 2-manifold, and $\varphi : N \rightarrow N$ is a convex isometric folding, any convex isometric folding is an isometric folding, then $\deg \varphi$ is ± 1 or 0. We consider only the case for which $\deg \varphi$ is zero otherwise $\varphi(N)$ cannot be embedded as a convex subset of $\mathbb{R}^2$ unless N is. In this case, the set of singularities of φ decomposes N into an even number of strata, say k , each of which is homeomorphic to $\varphi(N)$ and, hence,

$$\text{Vol } N = k \text{Vol } \varphi(N), \quad (3.2)$$

that is, e_N should be an even number. To calculate the exact value of e_N , consider first an orientable 2-compact manifold N . By using (1.1)

$$e_N = \frac{\text{Vol } F}{\text{Vol } \varphi(F)} \quad (3.3)$$

and this means that e_N can be calculated by calculating the volume of F and of its image $\varphi(F)$, but F is a closed square of diameter 2Δ and $\varphi(F)$ is a closed subset of F such that the distance $d(x, x')$ between any two points $x, x' \in \varphi(F)$ is at most Δ . The supremum of 2-dimensional volume of such set is $\phi(\Delta/2)^2$ and, hence, $2 < e_N$. But e_N is an even number. Hence, $e_N = 4$.

Now, let N be a nonorientable 2-compact manifold, i.e., a connected sum of n -projective planes. Elkholy [2] proved the theorem for $n = 1$.

The fundamental region in this case is a square or a rectangle of diameter 2Δ according to whether n is even or odd. If n is an even number, then

$$\text{Vol } F = 2\Delta^2 \quad (3.4)$$

and the result follows. Now, let n be an odd number. Then F is a rectangle of lengths $((n+1)/2)a$, $((n-1)/2)a$ and hence

$$\text{Vol } F = 4\Delta^2 \sin \theta \cos \theta = 4\Delta^2 \frac{a(n+1)/2}{a\sqrt{(n^2+1)}/2} \frac{a(n-1)/2}{a\sqrt{(n^2+1)}/2} = \frac{n^2-1}{n^2+1} 2\Delta^2. \quad (3.5)$$

Therefore, $e_N > 2$ for all $n > 1$. Since e_N is an even number, $e_N = 4$. $\square$

REFERENCES

- [1] E. M. Elkholy, *Isometric and topological folding of manifolds*, Ph.D. thesis, Southampton University, England, 1982.
- [2] ———, *Maximum isometric folding of the real projective plane*, J. Natur. Sci. Math. **25** (1985), no. 1, 25–30. MR 87d:53081. Zbl 589.53052.
- [3] E. M. Elkholy and A. E. Al-Ahmady, *Maximum volume of isometric foldings of compact 2-Riemannian manifolds*, Proc. Conf. Oper. Res. and Math. Methods Bull. Fac. Sc. Alex. **26 A** (1977), no. 1, 54–66.
- [4] S. A. Robertson, *Isometric folding of Riemannian manifolds*, Proc. Roy. Soc. Edinburgh Sect. A **79** (1977), no. 3–4, 275–284. MR 58 7486. Zbl 418.53016.
- [5] S. A. Robertson and E. M. Elkholy, *Invariant and equivariant isometric folding*, Delta J. Sci. **8** (1984), no. 2, 374–385.

ELKHOLY: DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, TANTA UNIVERSITY, TANTA, EGYPT