A CHARACTERIZATION OF MÖBIUS TRANSFORMATIONS

PIYAPONG NIAMSUP

(Received 9 March 2000 and in revised form 5 June 2000)

ABSTRACT. We give a new invariant characteristic property of Möbius transformations.

Keywords and phrases. Möbius transformations, Schwarzian derivative, Newton derivative.

2000 Mathematics Subject Classification. Primary 30C35.

1. Introduction. Throughout this paper, we let $w = f(z)$ be a nonconstant meromorphic function in $\mathbb{C}$ unless otherwise stated.

We consider the following properties.

PROPERTY 1.1. $w = f(z)$ transforms circles in the z -plane onto circles in the w -plane, including straight lines among circles.

PROPERTY 1.2. Suppose that $w = f(z)$ is analytic and univalent in a nonempty simply connected domain $\mathbb{R}$ on the z -plane. Let $ABCD$ be an arbitrary quadrilateral (not self-intersecting) contained in $\mathbb{R}$. If we set $A' = f(A)$, $B' = f(B)$, $C' = f(C)$, $D' = f(D)$ and if $A'B'C'D'$ is a quadrilateral on the w -plane which is not self-intersecting, then the following hold

$$\angle A + \angle C = \angle A' + \angle C', \quad \angle B + \angle D = \angle B' + \angle D'. \quad (1.1)$$

The following is a well-known principle of circle transformation of Möbius transformations.

THEOREM 1.3. $w = f(z)$ satisfies Property 1.1 if and only if $w = f(z)$ is a Möbius transformations.

In [1], it is shown that Property 1.1 implies Property 1.2 and a new invariant characteristic property of Möbius transformations is given as follows.

THEOREM 1.4. Let α be an arbitrary fixed real number such that $0 < \alpha < 2\pi$. Suppose that $w = f(z)$ is analytic and univalent in a nonempty simply connected domain $\mathbb{R}$ on the z -plane. Let $ABCD$ be an arbitrary quadrilateral (not self-intersecting) contained in $\mathbb{R}$ satisfying

$$\angle A + \angle C = \alpha. \quad (1.2)$$

If $A' = f(A)$, $B' = f(B)$, $C' = f(C)$, $D' = f(D)$ is a quadrilateral on the w -plane which is not self-intersecting, then the only function which satisfies

$$\angle A' + \angle C' = \alpha \quad (1.3)$$

is a Möbius transformation.

Theorem 1.4 gives an alternative proof of “the only if part” of Theorem 1.3. Motivated by the above results, we consider the following property.

PROPERTY 1.5. Let k be an arbitrary positive real number. For three arbitrary distinct points a, b , and c in $\mathbb{R}$ satisfying

$$\left| \frac{a-b}{c-b} \right| = k, \quad (1.4)$$

we have

$$\left| \frac{f(a)-f(b)}{f(c)-f(b)} \cdot \frac{f(c)}{f(a)} \right| = k. \quad (1.5)$$

In Section 3, we prove the following result concerning the mapping property of an analytic and univalent function on a connected domain.

THEOREM 1.6. Let k be an arbitrary positive real number. Let $w = f(z)$ be analytic and univalent in a nonempty connected domain $\mathbb{R}$ on the z -plane such that $f(z) \neq 0$ for all $z \in \mathbb{R}$. Then f satisfies Property 1.5 if and only if f is a Möbius transformation of the form $u/(z+v)$, $u \neq 0$.

2. Lemmas

DEFINITION 2.1. Let f be a complex-valued function. The Schwarzian derivative of f is defined as follows:

$$S_f(z) = \frac{f'''(z)}{f'(z)} - \frac{3}{2} \left(\frac{f''(z)}{f'(z)} \right)^2. \quad (2.1)$$

Similar to Schwarzian derivative, we have the following.

DEFINITION 2.2. Let f be a complex-valued function. We define the Newton derivative of f as follows:

$$N_f(z) = \left(z - \frac{f(z)}{f'(z)} \right)' = \frac{f(z)f''(z)}{(f'(z))^2}. \quad (2.2)$$

REMARK 2.3. Note that $N_f(z)$ is the first derivative of Newton’s method of f .

REMARK 2.4. Let f be a complex-valued function. It is well known that $S_f(z) = 0$ if and only if f is a Möbius transformation.

From Remark 2.4, we have observed that a similar result holds true when we replace Schwarzian derivative by the Newton derivative.

LEMMA 2.5. Let f be a complex-valued function. Then $N_f(z) = 2$ if and only if f is a Möbius transformation of the form $u/(z+v)$, $u \neq 0$.

PROOF. Let f be a Möbius transformation of the form $u/(z+v)$, $u \neq 0$, then it is easily checked that $N_f(z) = 2$. Let f be a complex-valued function such that $N_f(z) = 2$. It follows that

$$\left(z - \frac{f(z)}{f'(z)} \right)' = 2 \quad (2.3)$$

which implies that

$$z - \frac{f(z)}{f'(z)} = 2z - c_1, \quad (2.4)$$

where c_1 is a complex constant, thus

$$\frac{f(z)}{f'(z)} = -z + c_1 \quad (2.5)$$

or

$$\frac{1}{f(z)} \frac{df(z)}{dz} = \frac{1}{-z + c_1}. \quad (2.6)$$

From which it follows by a simple calculation that f is a Möbius transformation of the form $u/(z + v)$, $u \neq 0$. $\square$

3. Main result. In this section, we assume that $w = f(z)$ is analytic and univalent on a nonempty connected domain $\mathbb{R}$ on the z -plane such that $f(z) \neq 0$ for all $z \in \mathbb{R}$.

PROOF OF THEOREM 1.6. Let $f(z)$ be a Möbius transformation of the form $u/(z + v)$, $u \neq 0$. Let a, b , and c be arbitrary three distinct points in $\mathbb{R}$ such that

$$\left| \frac{a-b}{c-b} \right| = k. \quad (3.1)$$

We observe that

$$\frac{a-b}{c-b} \quad (3.2)$$

is the cross-ratio of a, b, c , and d , where d is the point at infinity. Since $f(z) = u/(z + v)$, $u \neq 0$, we have $f(d) = 0$. Since Möbius transformations preserve the cross-ratio, we obtain

$$\frac{f(a) - f(b)}{f(c) - f(b)} \cdot \frac{f(c)}{f(a)} = \frac{a-b}{c-b} \quad (3.3)$$

which implies that

$$\left| \frac{f(a) - f(b)}{f(c) - f(b)} \cdot \frac{f(c)}{f(a)} \right| = \left| \frac{a-b}{c-b} \right| = k. \quad (3.4)$$

Therefore, any Möbius transformation of the form $u/(z + v)$, $u \neq 0$ satisfies Property 1.5.

Conversely, let x be an arbitrary fixed point in $\mathbb{R}$. Then there exists a positive real number r such that the r circular neighborhood $N_r(x)$ of x is contained in $\mathbb{R}$.

Throughout the proof let $A = x + k\gamma$, $B = x$, $C = x - \gamma$. Since $\mathbb{R}$ is a nonempty connected domain on the z -plane, there exists a positive real number s such that if

$$0 < |\gamma| < s, \quad (3.5)$$

then A, B , and C are contained in $N_r(x)$.

Since $w = f(z)$ is univalent in $\mathbb{R}$, $f(A) = f(x + ky)$, $f(B) = f(x)$, and $f(C) = f(x - y)$ are distinct points. By assumption, we have

$$\left| \frac{f(x + ky) - f(x)}{f(x - y) - f(x)} \cdot \frac{f(x - y)}{f(x + ky)} \right| = k \quad (3.6)$$

for all y such that $0 < |y| < s$.

Let

$$h(y) = \frac{f(x + ky) - f(x)}{f(x - y) - f(x)} \cdot \frac{f(x - y)}{f(x + ky)}. \quad (3.7)$$

Then

$$|h(y)| = k \quad (3.8)$$

for all y such that $0 < |y| < s$. The function $h(y)$ extends analytically at zero by $h(0) = -k$. Hence, by the maximum modulus principle, we have $h(y) = -k$ for all y with $|y| < s$. In other words, we have

$$\frac{f(x + ky) - f(x)}{f(x - y) - f(x)} \cdot \frac{f(x - y)}{f(x + ky)} = -k \quad (3.9)$$

in $|y| < s$. This equality implies that

$$(f(x + ky) - f(x))f(x - y) = -k(f(x - y) - f(x))f(x + ky). \quad (3.10)$$

Differentiate this equality twice with respect to y and then set $y = 0$, we obtain

$$-k(k+1)(2(f'(x))^2 - f(x)f''(x)) = 0 \quad (3.11)$$

which implies that

$$2(f'(x))^2 - f(x)f''(x) = 0 \quad (3.12)$$

or

$$\frac{f(x)f''(x)}{(f'(x))^2} = 2. \quad (3.13)$$

By the identity theorem and Lemma 2.5, we conclude that f is a Möbius transformation of the form $u/(z + v)$, $u \neq 0$. $\square$

ACKNOWLEDGEMENT. I would like to thank the referees for valuable comments and suggestions. This work is supported by the Thailand Research Fund.

REFERENCES

- [1] H. Haruki and T. M. Rassias, *A new invariant characteristic property of Möbius transformations from the standpoint of conformal mapping*, J. Math. Anal. Appl. **181** (1994), no. 2, 320–327. MR 94m:30018. Zbl 796.39008.

PIYAPONG NIAMSUP: DEPARTMENT OF MATHEMATICS, CHIANGMAI UNIVERSITY, CHIANGMAI, 50200, THAILAND

E-mail address: scipnmsp@chiangmai.ac.th