FIXED POINTS VIA A GENERALIZED LOCAL COMMUTATIVITY

GERALD F. JUNGCK

(Received 24 May 2000)

ABSTRACT. Let $g : X \rightarrow X$. The concept of a semigroup of maps which is “nearly commutative at g ” is introduced. We thereby obtain new fixed point theorems for functions with bounded orbit(s) which generalize a recent theorem by Huang and Hong, and results by Jachymski, Jungck, Ohta, and Nikaido, Rhoades and Watson, and others.

2000 Mathematics Subject Classification. Primary 47H10, 54H25.

1. Introduction. By a *semi-group of maps* we mean a family H of self maps of a set X which is closed with respect to composition of maps ($f \circ g = fg$) and includes the identity map $i_d(x) = x$, for $x \in X$. We often associate with a function $g : X \rightarrow X$ following semi-groups:

$$O_g = \{g^n \mid n \in \mathbb{N} \cup \{0\}\}, \quad (1.1)$$

where $\mathbb{N}$ is the set of positive integers and $g^0 = i_d$, and

$$C_g = \{f : X \rightarrow X \mid fg = gf\}. \quad (1.2)$$

A quick check confirms that C_g is a semi-group.

If H is a semi-group of self maps of a set X and $a \in X$, $H(a) = \{h(a) \mid a \in H\}$. In particular, if $H = O_g$, $O_g(a) = \{g^n(a) \mid n \in \mathbb{N} \cup \{0\}\}$ and is called the orbit of g at a .

In general, [Lemma 3.2](#) and some theorems in [Section 3](#) will be stated in the context of semi-metric spaces. A *semi-metric* on a set X is a function $d : X \times X \rightarrow [0, \infty)$ such that $d(x, y) = d(y, x)$ for $x, y \in X$ and $d(x, y) = 0$ if and only if $x = y$. A *semi-metric space* is a pair $(X; d)$, where X is a topological space and d is a semi-metric on X . The topology $t(d)$ on X is generated by the sets $S(p, \epsilon) = \{x \mid d(x, p) < \epsilon\}$ with the requirement that p is an interior point of $S(p, \epsilon)$. A sequence $\{x_n\}$ in X converges in $t(d)$ to $p \in X$ (denoted as $x_n \rightarrow p$) if and only if $d(x_n, p) \rightarrow 0$. We let $t(d)$ be T_2 (Hausdorff) to ensure unique limits. Thus, a metric space (X, d) is a semi-metric space having the triangle inequality. For further details on semi-metric spaces, see, for example, [\[1, 4, 6\]](#).

If $g : X \rightarrow X$, a semi-metric space $(X; d)$ is *complete (g-orbitally complete)* if and only if every Cauchy sequence (in the usual sense) in X ($O_g(x)$) converges to a point of X . g is *continuous* at $p \in X$ if and only if whenever $\{x_n\}$ is a sequence in X and $x_n \rightarrow p$, then $f(x_n) \rightarrow f(p)$. And if S is a bounded subset of X , $\delta(S) = \sup\{d(x, y) \mid x, y \in S\}$.

We are now ready to focus on the intent of this paper, namely, to introduce a generalized “local commutativity” and to demonstrate the concept’s usefulness.

2. Nearly commutative semi-groups. In [2], a semi-group H of maps is said to be *near-commutative* if and only if for each pair $f, g \in H$, there exists $h \in H$ such that $fg = gh$. We generalize as follows.

DEFINITION 2.1. A semi-group H of self maps of a set X is *nearly commutative (n.c.) at $g : X \rightarrow X$* if and only if $(f \in H)$ implies that there exists $h \in H$ such that $fg = gh$.

Of course, O_g and C_g are n.c. at g . Observe also that a *near-commutative semi-group* H of self maps of a set X is n.c. at *each* $g \in H$. The following provides for each $a \in (0, \infty)$ an example of a semi-group $H = S_a$ of self maps which is not near-commutative but is n.c. at a particular $g : X \rightarrow X$.

EXAMPLE 2.2. Let $X = [0, \infty)$ and $a \in (0, \infty)$. Let $g(x) = ax$ and define

$$S_a = \{a^m x^n \mid x \in [0, \infty), n \in \mathbb{N}, m \in \mathbb{N} \cup \{0\}\}, \quad (2.1)$$

where S_a is *nearly commutative (n.c.) at g* . For if $f(x) = a^m x^n$ is a representative element of S_a , then $fg(x) = f(g(x)) = a^m (ax)^n = a^{m+n} x^n$. We want $h(x) = a^r x^s \in S_a$ such that $fg = gh$. Now, $g(h(x)) = a(a^r x^s) = a^{r+1} x^s$, so we can let $s = n$ and $r + 1 = m + n$; that is, $r = m + (n - 1)$. Since $n \in \mathbb{N}$ and $(n - 1), m \in \mathbb{N} \cup \{0\}$, s and r so designated imply $h \in S_a$. Thus, $(f \in H = S_a)$ implies that there exists $h \in H$ such that $fg = gh$. Since $i_d \in S_a$, S_a is clearly a semi-group, and we are finished. On the other hand, S_a is not a *near-commutative* semi-group. For example, let $f(x) = a^2 x^2$ and $h(x) = a^2 x^3$. We want $t(x) = a^r x^s$ such that $fh = ht$. So we must have $3s = 6$ and $(2 + 3r) = 6$. But then $r = 4/3$, and $r \notin \mathbb{N} \cup \{0\}$.

Now, let $\mathcal{M}_n$ and $\mathcal{N}_n$ denote the set of all $n \times n$ real matrices and the set of all nonsingular $n \times n$ real matrices, respectively. Then, both sets $\mathcal{M}_n$ and $\mathcal{N}_n$ are semi-groups of linear transformations $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ relative to composition of maps (matrix multiplication).

EXAMPLE 2.3. $\mathcal{N}_n$ is n.c. For if $A, B \in \mathcal{N}_n$, there exists $C = B^{-1}(AB) \in \mathcal{N}_n$ such that $AB = BC$.

EXAMPLE 2.4. $\mathcal{M}_n$ is n.c. at any $B \in \mathcal{N}_n$, by [Example 2.3](#). But $\mathcal{M}_n$ is not near commutative. For instance, if $n = 2$, $B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, and $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, there exists no 2×2 matrix C such that $AB = BC$.

Now, let $g : X \rightarrow X$. Since any semi-group of self maps which commute with g is a subset of C_g , we might hope that $H_g = \{f : X \rightarrow X \mid fg = gh \text{ for some } h : X \rightarrow X\}$ would be a maximal semi-group which is n.c. at g . However, H_g so defined need not be n.c. at g ! For example, let $X = [0, \infty)$, $g(x) = 1/(x + 1)$, and $f(x) = x/2$. Then $h(x) = 2x + 1$ satisfies $f(g(x)) = g(h(x))$ for $x \in [0, \infty)$. However, there exists no $k \in H_g$ such that $h(g(x)) = g(k(x))$; that is, $2(x + 1)^{-1} + 1 = (k(x) + 1)^{-1}$ (note that $x, k(x) \geq 0$).

Note that the map $g(x) = 1/(x + 1)$ was not surjective. So consider the following example.

EXAMPLE 2.5. Let X be any set and let $g : X \rightarrow X$ be surjective. Then the family of all self mappings of X , $\mathcal{F} = \{f \mid f : X \rightarrow X\}$, is n.c. at g . For suppose $f \in \mathcal{F}$; we need $h \in \mathcal{F}$ such that $fg(x) = gh(x)$ for all $x \in X$. So let $a \in X$. Since g is onto, we can choose $x_a \in X$ such that $g(x_a) = f(g(a))$. Choose such an x_a for each $a \in X$ and define $h(a) = x_a$. Then $h : X \rightarrow X$ and $g(h(a)) = g(x_a) = f(g(a))$ for $a \in X$; that is, $fg = gh$.

PROPOSITION 2.6. Suppose that H is a semigroup of maps which is n.c. at $g : X \rightarrow X$. If $f \in H$ and $n \in \mathbb{N}$, there exists $h_n \in H$ such that $fg^n = g^n h_n$ (i.e., H is n.c. at g^n).

PROOF. Let $f \in H$. Since, H is n.c. at g , there exists $h_1 \in H$ such that $fg = gh_1$. So suppose that $k \in \mathbb{N}$ such that $fg^k = g^k h_k$ for some $h_k \in H$. Then

$$fg^{k+1} = (fg^k)g = (g^k h_k)g = g^k(h_k g). \quad (2.2)$$

Since $h_k \in H$, there exists $h_{k+1} \in H$ such that $h_k g = gh_{k+1}$, and therefore (2.2) implies $fg^{k+1} = g^k(gh_{k+1}) = g^{k+1}h_{k+1}$, as desired. $\square$

Throughout this paper, P denotes a function $P : [0, \infty) \rightarrow [0, \infty)$ which is non-decreasing, and satisfies $\lim_{n \rightarrow \infty} P^n(t) = 0$ for $t \in [0, \infty)$. (For example, we could let $P(t) = \alpha t$ for some $\alpha \in (0, 1)$, or $t/(t+1)$.) And throughout this paper, we appeal to the following lemma.

LEMMA 2.7. Let H be a semi-group of self maps of a set X and suppose that H is nearly commutative at $g : X \rightarrow X$. Let $d : X \times X \rightarrow [0, \infty)$. Suppose that for each pair $x, y \in X$ there exists a choice $r = r(\{x, y\})$, $s = s(\{x, y\}) \in H$, and $u, v \in \{x, y\}$ for which

$$d(gx, gy) \leq P(d(ru, sv)). \quad (2.3)$$

Then, if $n \in \mathbb{N}$, for each pair $x, y \in X$ there exist $r_n, s_n \in H$ and $u_n, v_n \in \{x, y\}$ such that

$$d(g^n x, g^n y) \leq P^n(d(r_n u_n, s_n v_n)). \quad (2.4)$$

PROOF. By (2.3), inequality (2.4) holds for $n = 1$, so suppose that $n \in \mathbb{N}$ for which (2.4) is true. Then, if $x, y \in X$,

$$d(g^{n+1}x, g^{n+1}y) = d(g(g^n x), g(g^n y)) \leq P(d(ru, sv)), \quad (2.5)$$

where $r, s \in H$ and $u, v \in \{g^n x, g^n y\}$, by (2.3). Specifically, $u = g^n c$, $v = g^n d$, where $c, d \in \{x, y\}$. And since $r, s \in H$, there exist $r', s' \in H$ such that $rg^n = g^n r'$ and $sg^n = g^n s'$, by Proposition 2.6. So (2.4) implies that

$$d(ru, sv) = d(rg^n(c), sg^n(d)) = d(g^n(r'c), g^n(s'd)) \leq P^n(d(r_n u_n, s_n v_n)), \quad (2.6)$$

where $r_n, s_n \in H$ and $u_n, v_n \in \{r'c, s'd\}$. Thus, $r_n u_n \in \{(r_n r')c, (r_n s')d\}$, where $r_n r'$ and $r_n s'$ are elements of H , since H is a semi-group. So $r_n u_n = r_{n+1} u_{n+1}$, where $r_{n+1} \in \{r_n r', r_n s'\}$ (i.e., $r_{n+1} \in H$) and $u_{n+1} \in \{c, d\} \subset \{x, y\}$. Similarly, $s_n v_n = s_{n+1} v_{n+1}$, where $s_{n+1} \in H$ and $v_{n+1} \in \{x, y\}$. Thus, (2.6) implies that

$$d(ru, sv) \leq P^n(d(r_{n+1} u_{n+1}, s_{n+1} v_{n+1})), \quad r_{n+1}, s_{n+1} \in H, \quad u_{n+1}, v_{n+1} \in \{x, y\}. \quad (2.7)$$

But P is nondecreasing, and therefore (2.7) and (2.5) yield

$$\begin{aligned} d(g^{n+1}x, g^{n+1}y) &\leq P(P^n(d(r_{n+1}u_{n+1}, s_{n+1}v_{n+1}))) \\ &= P^{n+1}(d(r_{n+1}u_{n+1}, s_{n+1}v_{n+1})), \end{aligned} \quad (2.8)$$

with $r_{n+1}, s_{n+1} \in H$ and $u_{n+1}, v_{n+1} \in \{x, y\}$. So, (2.4) is true for all n by induction. $\square$

3. Fixed point theorems

DEFINITION 3.1. Let $(X; d)$ be a semi-metric space and let H be a semi-group of self maps of X . A map $g : X \rightarrow X$ is *P-contractive relative to H* if and only if (2.3) holds. (We will also say, “ g is a *P-contraction relative to H* .”)

LEMMA 3.2. Let $(X; d)$ be a T_2 semi-metric space and let H be a semi-group of self maps of X n.c. at $g : X \rightarrow X$. Suppose that g is *P-contractive relative to H* and that $M \subset X$ such that $B = \cup \{H(c) \mid c \in M\}$ is bounded. Then $d(g^n(x), g^n(y)) \rightarrow 0$ uniformly on B as $n \rightarrow \infty$. Specifically, if $\epsilon > 0$, there exists $k \in \mathbb{N}$ such that

$$(n \geq k) \implies (d(g^n(x), g^n(y)) < \epsilon \quad \forall x, y \in B). \quad (3.1)$$

PROOF. By hypothesis $\delta(B) < \infty$, $P^n(\delta(B)) \rightarrow 0$ as $n \rightarrow \infty$. Let $\epsilon > 0$. We can choose $k \in \mathbb{N}$ such that

$$P^n(\delta(B)) < \epsilon \quad \text{for } n \geq k. \quad (3.2)$$

Let $x, y \in B$. If $n \in \mathbb{N}$, since g is *P-contractive relative to H* , Lemma 2.7 yields $r_n, s_n \in H$ and $u_n, v_n \in \{x, y\} \subset B$ such that

$$d(g^n(x), g^n(y)) \leq P^n(d(r_n u_n, s_n v_n)). \quad (3.3)$$

Since $u_n \in B$, there exist $h \in H$ and $c \in M$ such that $u_n = h(c)$. But $r_n, h \in H$, so $r_n h \in H$. Therefore, $r_n u_n = (r_n h)(c) \in H(c) \subset B$. Likewise, $s_n v_n \in B$. But then $d(r_n u_n, s_n v_n) \leq \delta(B)$ and therefore,

$$P^n(d(r_n u_n, s_n v_n)) \leq P^n(\delta(B)) \quad \text{for } n \in \mathbb{N}, \quad (3.4)$$

since P is nondecreasing and n is arbitrary. Formulae (3.2), (3.3), and (3.4) imply

$$d(g^n(x), g^n(y)) < \epsilon \quad \text{for } n \geq k. \quad (3.5)$$

Since the choice of k in (3.2) was independent of x and y , (3.5) holds for all $x, y \in B$. $\square$

THEOREM 3.3. Let $(X; d)$ be a T_2 semi-metric space, and let H be a semi-group of self maps of X which is n.c. at $g \in H$. Suppose that $H(a)$ is bounded for some $a \in X$ and X is *g-orbitally complete*. If g is a *P-contraction relative to H* , then $g^n(a) \rightarrow c$ for some $c \in X$. If g is continuous at c , $g(c) = c$.

PROOF. Since X is *g-orbitally complete*, to show that $g^n(a) \rightarrow c$ for some $c \in X$ it suffices to show that $\{g^n(a)\}$ is a Cauchy sequence.

To this end, let $\epsilon > 0$. Since, $H(a)$ is bounded, [Lemma 3.2](#) with $B = H(a)$ implies that there exists $k \in \mathbb{N}$ such that

$$n \geq k \Rightarrow d(g^n(x), g^n(y)) < \epsilon \quad \forall x, y \in H(a). \quad (3.6)$$

Therefore, if $m > n \geq k$, $m = n + r$ for some $r \in \mathbb{N}$, and

$$d(g^n(a), g^m(a)) = d(g^n(a), g^n(g^r(a))) < \epsilon, \quad (3.7)$$

since $a, g^r(a) \in H(a)$. We conclude that $\{g^n(a)\}$ is Cauchy, and there exists $c \in X$ such that $g^n(a) \rightarrow c$.

Now, if g is continuous at c , $\lim_{n \rightarrow \infty} g(g^n(a)) = g(c)$, since $g^n(a) \rightarrow c$. But then $g^{n+1}(a) \rightarrow c$ also, so $g(c) = c$ since $(X; d)$ is a T_2 semi-metric space. $\square$

DEFINITION 3.4. Let X and Y be topological spaces. A map $g : X \rightarrow Y$ is *closed* if and only if $g(M)$ is closed in Y whenever M is a closed subset of X .

Note that the conclusion of [Lemma 3.2](#) asserts that $d(g^k(x_k), g^k(y_k)) \rightarrow 0$ for any sequences $\{x_k\}$ and $\{y_k\}$ in B .

THEOREM 3.5. Let $(X; d)$ be a bounded and complete T_2 semi-metric space, and let H be a semi-group of maps n.c. at $g \in H$. If g is closed and P -contractive relative to H ,

- (i) there exists $p \in X$ such that $\{p\} = \cap \{g^n(X) \mid n \in \mathbb{N}\}$,
- (ii) p is the unique fixed point of g ,
- (iii) $g^n(x) \rightarrow p$ for all $x \in X$.

PROOF. Let $x \in X$. By [Theorem 3.3](#), $\{g^n(x)\}$ converges to p for some $p \in X$. Moreover, $p \in \cap \{g^n(X) \mid n \in \mathbb{N}\}$. Otherwise, there exists $k \in \mathbb{N}$ such that $p \notin g^k(X)$. Since $g^k(X)$ is closed, there exists $\epsilon > 0$ such that $S(p, \epsilon) \cap g^k(X) = \emptyset$. Thus, $d(g^n(x), p) \geq \epsilon$ for $n \geq k$ since $g^n(X)$ is a subset of $g^k(X)$ for $n \geq k$. This contradicts the fact that $g^n(x) \rightarrow p$.

In fact, $\{p\} = \cap \{g^n(X) \mid n \in \mathbb{N}\}$. For if $q \in \cap \{g^n(X) \mid n \in \mathbb{N}\}$, for each $k \in \mathbb{N}$ we can choose $x_k, y_k \in X$ such that $g^k(x_k) = p$ and $g^k(y_k) = q$. So

$$d(p, q) = d(g^k(x_k), g^k(y_k)) \rightarrow 0, \quad (3.8)$$

by [Lemma 3.2](#) with $M = X$.

Clearly, (i) implies that p is a fixed point of g , since $g(\{p\}) \subset \{p\}$. Thus, if $x \in X$, $d(g^n(x), p) = d(g^n(x), g^n(p)) \rightarrow 0$ as $n \rightarrow \infty$, so (iii) holds. Similarly, if q is a fixed point of g , then $d(p, q) = d(g^n(p), g^n(q)) \rightarrow 0$, so that $q = p$. Thus, p is the only fixed point of g . $\square$

In the following we need the triangle inequality, so we require the underlying space to be a metric space.

THEOREM 3.6. Let (X, d) be a metric space and let H be a semi-group of self maps of X n.c. at some $g \in H$. Suppose that X is g -orbitally complete and there exists $k \in \mathbb{N}$ such that for each pair $x, y \in X$, there exist $r, s \in H$ and $u, v \in \{x, y\}$ for which

$$d(g^k x, g^k y) \leq P(d(ru, sv)). \quad (3.9)$$

(i) If there exists $a \in X$ such that $H(a)$ is bounded, then there exists $c \in X$ such that $\lim_{n \rightarrow \infty} g^n(a) = c$. If h is continuous for some $h \in H$, then $h(c) = c$. (Specifically, $g(c) = c$ if g is continuous at c .)

(ii) If $H(x)$ is bounded for each $x \in X$, there exists a unique $c \in X$ such that $g^n(x) \rightarrow c$ for all $x \in X$. If g is continuous at c , c is a unique common fixed point for all $h \in H$.

PROOF. Suppose that $H(a)$ is bounded. Since H is n.c. at g , [Proposition 2.6](#) says that H is n.c. at g^k . And X is g^k -orbitally complete since X is g -orbitally complete. Therefore, (3.9) and [Theorem 3.3](#) imply that

$$\lim_{m \rightarrow \infty} (g^k)^m(a) = c \quad \text{for some } c \in X. \quad (3.10)$$

To see that $\lim_{n \rightarrow \infty} g^n(a) = c$, let $\epsilon > 0$. Then (3.10) and [Lemma 3.2](#) (with $B = H(a)$) imply that there exists $p \in \mathbb{N}$ such that $d((g^k)^p(a), c) < \epsilon/2$ and $d(g^{kp}(x), g^{kp}(y)) < \epsilon/2$ for $x, y \in B$; that is,

$$d(g^{kp}(a), c) < \frac{\epsilon}{2}, \quad d(g^{kp}(g^i(a)), g^{kp}(a)) < \frac{\epsilon}{2} \quad \forall i \in \mathbb{N}, \quad (3.11)$$

since $g \in H \Rightarrow g^i(a) \in H(a)$. So, if $n > kp$, $n = kp + i$ for some $i \in \mathbb{N}$, and

$$d(g^n(a), c) \leq d(g^n(a), g^{kp}(a)) + d(g^{kp}(a), c), \quad (3.12)$$

or

$$d(g^n(a), c) \leq d(g^{kp}(g^i(a)), g^{kp}(a)) + d(g^{kp}(a), c) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon, \quad (3.13)$$

by (3.11). Consequently, $g^n(a) \rightarrow c$.

Now, let $h \in H$ and suppose that h is continuous at c . Then, $\lim_{n \rightarrow \infty} h(g^n(a)) = h(c)$ and

$$d(h(c), c) = \lim_{n \rightarrow \infty} d(hg^n(a), g^n(a)) = \lim_{n \rightarrow \infty} d(h(g^k)^n(a), (g^k)^n(a)). \quad (3.14)$$

But H is n.c. at g^k , so for $n \in \mathbb{N}$ there exists $h_n \in H$ such that $hg^{kn} = g^{kn}h_n$. Then, by (3.14),

$$d(h(c), c) = \lim_{n \rightarrow \infty} d((g^k)^n(h_n(a)), (g^k)^n(a)) = 0, \quad (3.15)$$

since $a, h_n(a) \in H(a)$ and [Lemma 3.2](#) holds for g^k . Thus, (i) holds.

To prove (ii), suppose that $H(x)$ is bounded for each $x \in X$. If $a, b \in X$, $g^n(a) \rightarrow c_a$ and $g^n(b) \rightarrow c_b$ for some $c_a, c_b \in X$ by (i). But $c_a = c_b$, since $H(a) \cup H(b)$ is bounded, and therefore, [Lemma 3.2](#) applied to g^k implies that $d((g^k)^n(a), (g^k)^n(b)) = 0$.

Thus, there exists a unique $c \in X$ such that $g^n(x) \rightarrow c$ for all $x \in X$. We know that $g(c) = c$ by part (i), if g is continuous at c . Since $g^n(d) = d$ for all n if d is a fixed point of g , and therefore $g^n(d) \rightarrow d$, c must be the only fixed point of g . Moreover, $h(c) = c$ for all $h \in H$ (even though h may not be continuous). This follows, since [Proposition 2.6](#) applied to g^k implies that for each $n \in \mathbb{N}$,

$$d(c, h(c)) = d((g^k)^n(c), h(g^k)^n(c)) = d((g^k)^n(c), (g^k)^n(h_n(c))) \quad (3.16)$$

for some $h_n \in H$. But $H(c)$ is bounded, so [Lemma 3.2](#) applied to g^k implies that the right member of (3.16) converges to zero as $n \rightarrow \infty$, and thus, $c = h(c)$. $\square$

REMARK 3.7. [Theorem 3.3](#) appreciably generalizes Theorem 2.1 in [5] and [Theorem 3.6](#) generalizes Corollary 2.3 in [5]—and hence Theorem 2 in [3] and the theorems of Rhoades and Watson [9]. Note that in [Theorem 3.6\(ii\)](#), the mappings $h \in H$ ($h \neq g$) need not be continuous. Remember also that C_g and O_g are special instances of H .

The following example suggests that the requirement in [Theorem 3.6\(ii\)](#), that $H(x)$ be bounded for each $x \in X$, is not as restrictive as may first appear.

EXAMPLE 3.8. Let $S = \{\text{continuous functions } f : [0, \infty) \rightarrow [0, \infty) \mid \text{there exists } a_f \in (0, \infty) \text{ such that } f(x) < x \text{ for } x > a_f\}$. (So, e.g., $\{f \mid f(x) = mx + b, m \in [0, 1) \text{ and } b \geq 0\} \subset S$, and $\ln(x+b) \in S$ for $b \geq 1$.) Then (1) $S \cup \{i_d\}$ is a semi-group under composition of functions, and (2) $O_f(x)$ is bounded for $f \in S$ and $x \in [0, \infty)$.

First note that, we can let M_f denote the maximum value of f on $[0, a_f]$ for each $f \in S$ since each f is continuous. To see that (1) is true, let $f, g \in S$. We need only to show that $g \circ f = gf \in S$. Clearly, gf is a continuous self map of $[0, \infty)$. So let $a_{gf} = \max\{a_f, M_g\}$ and suppose that $x > a_{gf}$. We want $gf(x) < x$. Now, $x > a_{gf}$ implies that $x > a_f$ so that (i) $f(x) < x$. If $f(x) > a_g$, then $g(f(x)) < f(x) < x$ by (i) and the definition of a_g . If $f(x) \leq a_g$, $g(f(x)) \leq M_g \leq a_{gf} < x$. So, in any event, $(g \circ f)(x) < x$ if $x > a_{gf}$, and thus, $g \circ f \in S$. (2) follows easily by using induction to show that $(f \in S)$ implies that (if $x \in [0, \infty)$, $f^n(x) \leq \max\{x, M_f\}$ for $n \in \mathbb{N}$). We omit the details.

If we let $P(t) = \alpha t$ for fixed $\alpha \in (0, 1)$ and $t \in [0, \infty)$, we have the following corollary.

COROLLARY 3.9. Let (X, d) be a bounded complete metric space and let $g : X \rightarrow X$ be continuous. Suppose that H is a semi-group of self maps of X n.c. at g and $g \in H$. If there exists $\alpha \in (0, 1)$ such that for any pair $x, y \in X$ there exist $r, s \in H$ and $u, v \in \{x, y\}$ for which

$$d(gx, gy) \leq \alpha d(ru, sv), \quad (3.17)$$

then there exists a unique $c \in X$ such that $g^n(x) \rightarrow c$ for $x \in X$, and $c = gc = hc$ for all $h \in H$.

4. Some consequences

DEFINITION 4.1. A gauge function is an upper semicontinuous (u.s.c.) function $\phi : [0, \infty) \rightarrow [0, \infty)$ such that $\phi(0) = 0$ and $\phi(t) < t$ for all $t > 0$.

LEMMA 4.2. Let (X, d) be a metric space and let H be a semi-group of self maps of X which is n.c. at $g \in H$. Suppose that $H(x, y) = H(x) \cup H(y)$ is bounded for $x, y \in X$ and there exists a gauge function ϕ such that

$$d(gx, gy) \leq \phi(\delta(H(x, y))) \quad \text{for } x, y \in X. \quad (4.1)$$

Then, there exists a nondecreasing continuous function $P : [0, \infty) \rightarrow [0, \infty)$ such that $P^n(t) \rightarrow 0$ for all $t > 0$ and which satisfies the following condition: for any pair $x, y \in X$ there exist $r = r(x, y)$, $s = s(x, y) \in H$, and $u, v \in \{x, y\}$ such that

$$d(gx, gy) \leq P(d(ru, sv)). \quad (4.2)$$

PROOF. Let $x, y \in X$ and suppose that (4.1) holds. Since, ϕ is a gauge function, as is well known [2], there exists a nondecreasing continuous function $P : [0, \infty) \rightarrow [0, \infty)$ such that $P^n(t) \rightarrow 0$ for $t \geq 0$, and

$$\phi(t) < P(t), \quad P(t) < t \quad \forall t \in (0, \infty). \quad (4.3)$$

Since P is continuous, (4.3) implies that for any $t > 0$, there exists $\epsilon_t \in (0, t)$ such that

$$t' \in (t - \epsilon_t, t + \epsilon_t) \Rightarrow \phi(t) < P(t'). \quad (4.4)$$

And since $H(x, y)$ is bounded, the definition of δ implies that there exist $r, s \in H$ and $u, v \in \{x, y\}$ such that, with $t = \delta(H(x, y))$,

$$t = \delta(H(x, y)) \geq d(ru, sv) > \delta(H(x, y)) - \epsilon_t. \quad (4.5)$$

So, with $t' = d(ru, sv)$, (4.4) and (4.5) imply that

$$\phi(\delta(H(x, y))) < P(d(ru, sv)). \quad (4.6)$$

Therefore, (4.1) implies that $d(gx, gy) \leq P(d(ru, sv))$. $\square$

The following theorem provides a generalization of Theorem 2.1 in [2].

THEOREM 4.3. *Let (X, d) be a complete metric space and let H be a semi-group of self maps of X which is n.c. at some $g \in H$. Suppose that the following conditions are satisfied:*

- (i) $H(x)$ is bounded for all $x \in X$, g is continuous,
- (ii) there exists a gauge function ϕ and $k \in \mathbb{N}$ such that

$$d(g^k x, g^k y) \leq \phi(\delta(H(x, y))) \quad \text{for } x, y \in X. \quad (4.7)$$

Then

- (a) H has a unique common fixed point c and $g^n(x) \rightarrow c$ for $x \in X$.
- (b) If for each $h \in H - \{i_d\}$ there exists $k = k_h \in \mathbb{N}$ such that (4.7) holds with $g = h$, then

$$h^n(x) \rightarrow c \quad \forall x \in X, h \in H - \{i_d\}. \quad (4.8)$$

PROOF. Now, (i) implies that $H(x, y) = H(x) \cup H(y)$ is bounded for $x, y \in X$. To see that (a) is true, note that H is n.c. at g^k by Proposition 2.6 and substitute g^k for g in Lemma 4.2 to conclude that (3.9) holds. Consequently, we can appeal to Theorem 3.6(ii) to obtain a $c \in X$ such that $g^n(x) \rightarrow c$ for $x \in X$. And since g is continuous, c is the unique fixed point of g and a fixed point for each $h \in H$. Thus, c is the unique common fixed point of H (remember, $g \in H$) and therefore (a) holds.

To prove (b) note that, by part (a), if $h \in H - \{i_d\}$, $h \neq g$, $h^n(c) = g(c) = c$ for $n \in \mathbb{N}$. But Theorem 3.6 applied to h yields a unique $c_1 \in X$ such that $h^n(x) \rightarrow c_1$ for all $x \in X$. Since $h^n(c) = c$ for all n , $c_1 = c$. $\square$

REMARK 4.4. Theorem 4.3 generalizes Theorem 2.1 in [2] in the following ways:

- (i) The semi-group H is not required to be near-commutative (i.e., n.c. at each $h \in H$), but n.c. only at g ,

- (ii) g is the only member of H required to be continuous,
- (iii) in (b), (4.7) is required to hold only for $k = k_h$, not for all $k \geq k_h$.

Theorem 4.3 yields the following corollary, which generalizes the theorem of Ohta and Nikaido [8] by requiring only that the orbits of f —but not all of X —be bounded.

COROLLARY 4.5. *Let f be a continuous self mapping of a metric space (X, d) having bounded orbits $O_f(x)$ for all $x \in X$. If there exist $c \in (0, 1)$ and $k \in \mathbb{N}$ such that*

$$d(f^k x, f^k y) \leq c \delta(\{f^i t \mid t \in \{x, y\}, i \in \mathbb{N} \cup \{0\}\}) \quad (4.9)$$

for all $x, y \in X$, then f has a unique fixed point.

Observe that **Lemma 3.2** does not require that $g \in H$, whereas the theorems in **Section 3** do. The requirement that $g \in H$ was convenient in the proof, but the following proposition says that it is not necessary when $O_g(a)$ is bounded. Moreover, this result is needed for the proof of **Theorem 4.7**.

PROPOSITION 4.6. *If H is a semi-group of self maps n.c. at g and $g \notin H$, then $H_g = \{g^n h \mid n \in \mathbb{N} \cup \{0\} \text{ and } h \in H\}$ is a semi-group which is n.c. at g . Moreover, $g \in H_g$ and $H \subset H_g$.*

PROOF. H_g is a semi-group. For if $g^n h_1, g^m h_2 \in H_g$, since H is n.c. at g , we have $g^n h_1 g^m h_2 = g^n (h_1 g^m) h_2 = g^n (g^m h_3) h_2 = g^{n+m} (h_4)$, where $h_4 = h_3 h_2 \in H$.

H_g is n.c. at g , since (H n.c. at g) implies that there exists $h_2 \in H$ such that $(g^n h)g = g^n (hg) = g^n (gh_2) = g(g^n h_2)$. $\square$

It is clear that if $g : X \rightarrow X$ is a P -contraction relative to H , then it is certainly a P -contraction relative to H_g since $H \subset H_g$. We use this fact in the proof of **Theorem 4.7**.

THEOREM 4.7. *Let C be a compact subset of a normed linear space X which is star-shaped with respect to $q \in C$. Let $T : C \rightarrow C$ be continuous and let H be a semi-group of affine maps $I : C \rightarrow C$ n.c. at T such that $I(q) = q$. If for each pair $x, y \in C$ there exist $I, J \in H$ and $u, v \in \{x, y\}$ for which*

$$\|Tx - Ty\| \leq \|Iu - Jv\|, \quad (4.10)$$

then there exists $a \in C$ such that $a = Ta$ and $a = Ia$ for all continuous $I \in H$.

PROOF. Choose a sequence $\{k_n\}$ in $(0, 1)$ such that $k_n \rightarrow 1$, and for each $n \in \mathbb{N}$, let

$$T_n(x) = k_n Tx + (1 - k_n)q. \quad (4.11)$$

Since C is star-shaped with respect to q , $T_n : C \rightarrow C$ for $n \in \mathbb{N}$. Moreover, if $I \in H$, there exists $J \in H$ such that

$$\begin{aligned} IT_n x &= I(k_n Tx + (1 - k_n)q) = k_n I(Tx) + (1 - k_n)Iq \\ &= k_n T(Jx) + (1 - k_n)q = T_n Jx, \end{aligned} \quad (4.12)$$

since I is affine, H is n.c. at T , and $Iq = q$. Thus, for each $n \in \mathbb{N}$, H is a semi-group of affine maps which is n.c. at T_n . Then, by **Proposition 4.6**, H_{T_n} is a semi-group of self maps of C which is n.c. at T_n , $T_n \in H_{T_n}$, and $H \subset H_{T_n}$ for $n \in \mathbb{N}$.

Now fix n . By hypothesis, for each pair $x, y \in C$ there exist $I, J \in H(\subset H_{T_n})$ and $u, v \in \{x, y\}$ such that

$$\|Tx - Ty\| \leq \|Iu - Jv\|, \quad (4.13)$$

so

$$\|T_n x - T_n y\| \leq k_n \|Iu - Jv\|, \quad (4.14)$$

by (4.11). Therefore, since T_n is continuous and $k_n \in (0, 1)$, Corollary 3.9 applied to T_n and H_{T_n} (C compact implies that C is bounded and complete) implies that there exists a unique $x_n \in C$ such that

$$x_n = T_n(x_n) = I(x_n) \quad \forall I \in H_{T_n}. \quad (4.15)$$

Thus we have a sequence $\{x_n\}$ in C which satisfies (4.15). Since C is compact, $\{x_n\}$ has a subsequence $\{x_{i_n}\}$ which converges to some $a \in C$. Equations (4.11) and (4.15) thus imply that

$$a = \lim_{n \rightarrow \infty} x_{i_n} = \lim_{n \rightarrow \infty} k_{i_n} T x_{i_n} + \lim_{n \rightarrow \infty} (1 - k_{i_n}) q = \lim_{n \rightarrow \infty} I x_{i_n}. \quad (4.16)$$

But T is continuous, so (4.16) implies that $a = Ta$, and $a = Ia$ for all continuous I . $\square$

REMARK 4.8. We see that Theorem 4.7 does indeed extend Theorem 3 in [7] if we observe that the family $\mathcal{F}$ in Theorem 3 [7], is a family of sets which is a subset of C_g . We can let

$$H = \{\text{maps } h : C \rightarrow C \mid h \text{ is affine, } h \in C_g\}. \quad (4.17)$$

Then H is a semi-group and $\mathcal{F} \subset H$.

5. Conclusion. We conclude with further evidence of the generality and applicability of the concept of being nearly commutative at a function g . The theorem below generalizes Theorem 4.2 in [5] by replacing the semi-group C_{gf} with a more general semi-group H .

THEOREM 5.1. *Let f and g be commuting self maps of a compact metric space (X, d) such that gf is continuous. If H is a semi-group of self maps of X which is n.c. at gf , and*

$$fx \neq gy \Rightarrow d(fx, gy) < \delta(H(x, y)), \quad (5.1)$$

then there exists a unique point $a \in X$ such that $a = fa = ga = ha$ for all $h \in H$.

We leave the proof of Theorem 5.1 to the interested reader.

REFERENCES

- [1] T. L. Hicks and B. E. Rhoades, *Fixed points for pairs of mappings in d -complete topological spaces*, Int. J. Math. Math. Sci. **16** (1993), no. 2, 259–266. [MR 93k:54076](#). [Zbl 796.54049](#).
- [2] Y.-Y. Huang and C.-C. Hong, *Common fixed point theorems for semigroups on metric spaces*, Int. J. Math. Math. Sci. **22** (1999), no. 2, 377–386. [MR 2000b:54053](#). [Zbl 944.47034](#).
- [3] J. Jachymski, *A generalization of the theorem by Rhoades and Watson for contractive type mappings*, Math. Japon. **38** (1993), no. 6, 1095–1102. [MR 94k:54076](#). [Zbl 795.54054](#).

- [4] J. Jachymski, J. Matkowski, and T. Świątkowski, *Nonlinear contractions on semimetric spaces*, J. Appl. Anal. **1** (1995), no. 2, 125–134. [CMP 1 395 268](#).
- [5] G. Jungck, *Common fixed points for commuting and compatible maps on compacta*, Proc. Amer. Math. Soc. **103** (1988), no. 3, 977–983. [MR 89h:54030](#). [Zbl 661.54043](#).
- [6] ———, *Fixed point theorems for semi-groups of self maps of semi-metric spaces*, Int. J. Math. Math. Sci. **21** (1998), no. 1, 125–132. [CMP 1 486 967](#). [Zbl 892.54026](#).
- [7] G. Jungck and S. Sessa, *Fixed point theorems in best approximation theory*, Math. Japon. **42** (1995), no. 2, 249–252. [MR 96f:47112](#). [Zbl 834.54026](#).
- [8] M. Ohta and G. Nikaido, *Remarks on fixed point theorems in complete metric spaces*, Math. Japon. **39** (1994), no. 2, 287–290. [MR 94m:54099](#). [Zbl 802.47055](#).
- [9] B. E. Rhoades and B. Watson, *Generalized contractions and fixed points in metric spaces*, Math. Japon. **34** (1989), no. 6, 975–982. [MR 91c:54057](#). [Zbl 697.54023](#).

GERALD F. JUNGCK: DEPARTMENT OF MATHEMATICS, BRADLEY UNIVERSITY, PEORIA, IL 61625, USA

E-mail address: [gfj@hilltop.bradley.edu](mailto:gfej@hilltop.bradley.edu)