ON THE DIOPHANTINE EQUATION $Ax^2 + 2^{2m} = y^n$

FADWA S. ABU MURIEFAH

(Received 6 May 1999 and in revised form 20 February 2000)

ABSTRACT. Let h denote the class number of the quadratic field $\mathbb{Q}(\sqrt{-A})$ for a square free odd integer $A > 1$, and suppose that $n > 2$ is an odd integer with $(n, h) = 1$ and $m > 1$. In this paper, it is proved that the equation of the title has no solution in positive integers x and y if n has any prime factor congruent to 1 modulo 4. If n has no such factor it is proved that there exists at most one solution with x and y odd. The case $n = 3$ is solved completely. A result of E. Brown for $A = 3$ is improved and generalized to the case where A is a prime $\not\equiv 7 \pmod{8}$.

2000 Mathematics Subject Classification. Primary 11D41.

1. Introduction. Let A, m, n denote positive integers where n is odd > 1 and A square free odd integer. Let $K = \mathbb{Q}(\sqrt{-A})$, where $\mathbb{Q}$ is the field of rational numbers, let further h denote the number of classes of ideals in K and suppose $(h, n) = 1$. In this paper, we consider the Diophantine equation $Ax^2 = 2^{2m} = y^n$, where x and y are integers. The case $A = 1$ was studied in [1] so we will assume that $A > 1$. The first result regarding this equation is due to Nagell [5] who proved that when $m = 0, 1$, this equation has no solutions in integers x and y under the above assumptions about A and n so we will suppose that $m > 0$. Since n is odd, there is no loss of generality in considering only odd primes p and x, y positive integers, so we will assume this in what follows.

We start by proving the main result of this paper.

THEOREM 1.1. *Let $A > 1$ be a square free odd integer, p an odd prime with $(h, p) = 1$ and $m \geq 1$. Then the Diophantine equation*

$$Ax^2 + 2^{2m} = y^p \tag{1.1}$$

has no solution with x odd in any of the following cases:

- (i) if $A = 3$;
- (ii) if $p \equiv 1 \pmod{4}$;
- (iii) if $A \equiv 3 \pmod{4}$ and $p > 3$.

For $p = 3$, such a solution exists if and only if A is the square-free part of either $(1/3)(1 + 2^{m+3})$ with m even or of $(1/3)(2^{2m} - 1)$, although in these cases, there might be other solutions if $3 \mid h$.

PROOF. We factorize (1.1) in the field K ,

$$(2^m + x\sqrt{-A})(2^m - x\sqrt{-A}) = y^p. \tag{1.2}$$

Now the principal ideal $[2^m + x\sqrt{-A}]$ and its conjugate ideal are coprime, so $[2^m + x\sqrt{-A}] = \pi^p$ for some ideal π in K . It follows that π^p is a principal ideal and since $(h, p) = 1$, therefore π is a principal ideal, say $\pi = [\xi]$ for some element ξ in K . So we get the equation

$$[2^m + x\sqrt{-A}] = [\xi]^p, \quad (1.3)$$

and, consequently,

$$(2^m + x\sqrt{-A}) = \varepsilon \xi^p, \quad (1.4)$$

for some unit ε in K . Therefore we have the following three cases:

$$\begin{aligned} x\sqrt{-3} + 2^m &= \left(\frac{1 \pm \sqrt{-3}}{2}\right) \left(\frac{a + b\sqrt{-3}}{2}\right)^3, & a \equiv b \pmod{2}, \\ x\sqrt{-A} + 2^m &= \left(\frac{a + b\sqrt{-A}}{2}\right)^p, & a \equiv b \equiv 1 \pmod{2}, \\ x\sqrt{-A} + 2^m &= (a + b\sqrt{-A})^p, \end{aligned} \quad (1.5)$$

for some rational integers a and b .

Equating the imaginary parts in the first case we get

$$16x = \pm(a^3 - 9ab^2) + (3a^2b - 3b^3), \quad (1.6)$$

and we can absorb the lower sign into a . Then

$$16x = (a + b)^3 - 12ab^2 - 4b^3. \quad (1.7)$$

Since a and b have the same parity, we write $2c = a + b$, and obtain

$$2x = c^3 - 3cb^2 + b^3. \quad (1.8)$$

Equation (1.8) is impossible, since the right-hand side is odd unless both b and c are even, and then this side is divisible by 8 if they are which is not possible since x is odd. So this case does not arise.

The second case arises only if $A \equiv 3 \pmod{4}$, and we will prove that in this case $p = 3$ and $A \neq 3$.

Observe that $((a + b\sqrt{-A})/2)^p \in Z[\sqrt{-A}]$ only if $A \equiv 3 \pmod{8}$ and $p = 3$ and then equating the real parts in this case, we get

$$2^{m+3} = a(a^2 - 3Ab^2). \quad (1.9)$$

Since a is odd we get $a = \pm 1$ and then

$$\pm 2^{m+3} = 1 - 3Ab^2. \quad (1.10)$$

Now $A > 1$, so only the negative sign holds and then

$$Ab^2 = \frac{1 + 2^{m+3}}{3}. \quad (1.11)$$

Considering this equation modulo 3 we deduce that m should be even. If $A = 3$, then we get

$$-2^{m+3} = (1-3b)(1+3b). \quad (1.12)$$

So

$$2^t = 1+3b, \quad -2^k = 1-3b, \quad (1.13)$$

where $t+k = m+3$. By adding these two equations we get $t = 2$, $k = 1$, which is impossible since $m \geq 1$.

Finally the third case can occur for all A , and we will prove that there is no solution when either $p \equiv 1 \pmod{4}$ or $A \equiv 3 \pmod{4}$.

Since x is odd it follows that $y = a^2 + Ab^2$ is odd, so a and b have opposite parity. On equating the real parts we get

$$2^m = a \sum_{r=0}^{(p-1)/2} \binom{p}{2r} a^{p-2r-1} (-Ab^2)^r. \quad (1.14)$$

Here $\sum$ is odd, since the first and the last terms have opposite parity and the rest are all even. So $a = \pm 2^m$, b is odd and from (1.14) we get

$$\pm 1 = \sum_{r=0}^{(p-1)/2} \binom{p}{2r} 2^{m(p-2r-1)} (-Ab^2)^r. \quad (1.15)$$

Then $\pm 1 \equiv 2^{m(p-1)} \pmod{p}$ and so the lower sign is impossible. That is,

$$1 = \sum_{r=0}^{(p-1)/2} \binom{p}{2r} 2^{m(p-2r-1)} (-Ab^2)^r, \quad (1.16)$$

and $a = 2^m$. So $y = 2^{2m} + Ab^2$.

Now suppose that $p \equiv 1 \pmod{4}$, say $p = 1 + 2^k u$, where $(u, 2) = 1$ and $k \geq 2$. Since both b and A are odd,

$$b^{p-1} = (b^u)^{2^k} \equiv 1 \pmod{2^{k+2}}, \quad (-A)^{(p-1)/2} = (A^u)^{2^{k-1}} \equiv 1 \pmod{2^{k+1}}. \quad (1.17)$$

Then from (1.16) we get

$$\begin{aligned} 1 &\equiv \binom{p}{3} 2^{2m} (-Ab^2)^{(p-3)/2} + p b^{p-1} (-A)^{(p-1)/2} \pmod{2^{k+1}} \\ &\equiv \frac{p(p-2)}{3} \cdot 2^{k+2m-1} u (-Ab^2)^{(p-1)/3} + p \pmod{2^{k+1}}, \end{aligned} \quad (1.18)$$

since $m \geq 1$, therefore $k+2m-1 \geq k+1$, so from (1.18) we get $3 \equiv 3p \pmod{2^{k+1}}$. Hence $p \equiv 1 \pmod{2^{k+1}}$ which is not possible. We conclude that there is no solution when $p \equiv 1 \pmod{4}$. If $A \equiv 3 \pmod{4}$, then considering (1.16) modulo 4 we get $p \equiv 1 \pmod{4}$ hence there is no solution when $A \equiv 3 \pmod{4}$.

Now let $p = 3$ in (1.16) then

$$1 = 2^{2m} - 3Ab^2 \quad (1.19)$$

or $Ab^2 = (2^{2m} - 1)/3$. This completes the proof. $\square$

REMARK 1.2. From [Theorem 1.1](#), we note that to solve (1.1) it is sufficient to consider (1.16) where b is odd, $p \equiv 3 \pmod{4}$, and $A \equiv 1 \pmod{4}$. If there is a solution then $y = 2^{2m} + Ab^2$.

Now we prove the following theorem which gives us the number of solutions of our equation.

THEOREM 1.3. *For a given A , if (1.1) has a solution in x odd where $(h, p) = 1$, then it is unique.*

PROOF. If $A \equiv 3 \pmod{4}$, we have proved that there is a solution only if $p = 3$, and we have found this unique solution. If $A \equiv 1 \pmod{4}$, then from the last proof it is sufficient to consider (1.16), where b is odd and $p \equiv 3 \pmod{4}$. Suppose $b_1 > b > 0$ is another solution, then from (1.16) we obtain

$$1 = \sum_{r=0}^{(p-1)/2} \binom{p}{2r} 2^{m(p-2r-1)} (-Ab_1^2)^r. \quad (1.20)$$

Subtracting (1.20) from (1.16) and dividing by $b_1^2 - b^2$, we get

$$\begin{aligned} 0 &= \sum_{r=0}^{(p-1)/2} \binom{p}{2r} \frac{b_1^{2r} - b^{2r}}{b_1^2 - b^2} \cdot 2^{m(p-2r-1)} (-A)^r \\ &\equiv p \cdot \frac{b_1^{p-1} - b^{p-1}}{b_1^2 - b^2} \pmod{2}. \end{aligned} \quad (1.21)$$

Since $p \equiv 3 \pmod{4}$, the number $(b_1^{p-1} - b^{p-1}) / (b_1^2 - b^2)$ is odd, so (1.21) is impossible and the solution is unique as required. $\square$

Now we prove that to solve (1.1) it is sufficient to consider only x odd. First we need the following lemma.

LEMMA 1.4 [4]. *The Diophantine equations*

$$Ax^2 + 1 = 2y^n, \quad A \equiv 1 \pmod{4}, \quad Ax^2 + 1 = 4y^n, \quad A \equiv 3 \pmod{4}, \quad (1.22)$$

have no solutions in positive integers with $y > 1$, $n > 2$, $2 \nmid ny$ and $(n, h) = 1$.

THEOREM 1.5. *If $A = 3$, equation (1.1) has a solution with x even only if $m \equiv -1 \pmod{p}$, and this solution is given by $x = 2^m$; for all other $A \not\equiv 7 \pmod{8}$ with $(h, p) = 1$ there exists a solution with x even of the form $x = 2^u X$ with X odd, if and only if there is a solution of the equation $AX^2 + 2^{2(m-u)} = Y^p$.*

PROOF. If x is even then y is even, so let $x = 2^u X$, $y = 2^v \cdot Y$, where $u > 0$, $v > 0$, $(2, X) = (2, Y) = 1$. Then (1.1) becomes

$$A(2^u X)^2 + 2^{2m} = 2^{vp} Y^p. \quad (1.23)$$

We have three cases:

(1) $vp > 2u = 2m$. Then cancelling 2^{2m} in (1.23) we get

$$AX^2 + 1 = 2^{vp-2m} Y^p, \quad (1.24)$$

where X is odd. Now $A \not\equiv 7 \pmod{8}$, so $vp - 2m = 1$ or 2 .

If $A \equiv 1 \pmod{4}$ then $\nu p - 2m = 1$ and so $AX^2 + 1 = 2Y^p$. This equation has no solution from [Lemma 1.4](#). If $A \equiv 3 \pmod{8}$, then $\nu p - 2m = 2$, so $AX^2 + 1 = 4Y^p$, and again from [Lemma 1.4](#) this equation has no solution in integers with $Y > 1$. Let $Y = 1$, then $AX^2 + 1 = 4Y^p$ implies that $A = 3$, $X = 1$ and hence $x = 2^m$, also $\nu p = 2m + 2$ implies that $m \equiv -1 \pmod{p}$,

(2) $2u > 2m = \nu p$. Then cancelling 2^{2m} in (1.23) we get $A(2^{u-m}X)^2 + 1 = Y^p$. This equation has no solution [[5](#), Theorem 25].

(3) $2m > 2u = \nu p$. Then

$$AX^2 + 2^{2(m-u)} = Y^p, \quad (1.25)$$

and this is (1.1) with x odd and smaller m . $\square$

REMARK 1.6. From the proof of the last theorem we deduce that to solve (1.1) in even integers when $A \neq 3$ and $A \not\equiv 7 \pmod{8}$, it is sufficient to consider the equation

$$AX^2 + 2^{2(m-u)} = Y^p, \quad (1.26)$$

where $x = 2^u X$, $y = 2^\nu \cdot Y$, $m > u > 0$, $\nu > 0$, $(2, X) = (2, Y) = 1$, and $2u = \nu p$.

Summarizing the above we give the following theorem.

THEOREM 1.7. *The Diophantine equation (1.1) where $A \not\equiv 7 \pmod{8}$ and $(h, p) = 1$ has no integer solution if $p \equiv 1 \pmod{4}$. In particular, the equation $px^2 + 2^{2m} = y^p$ has no solution for all $p > 3$ and $p \not\equiv 7 \pmod{8}$.*

PROOF. If x is odd, then from [Theorem 1.1](#), equation (1.1) has no solution when $p \equiv 1 \pmod{4}$. Now let x be even then from [Theorem 1.5](#) it is sufficient to consider the equation

$$AX^2 + 2^{2(m-u)} = Y^p, \quad (1.27)$$

where X is odd and $0 < u < m$. Since $p \equiv 1 \pmod{4}$ then again [Theorem 1.1](#) implies that there is no solution.

Now the class number of the field $\mathbb{Q}(\sqrt{-p})$ is less than p , so as above the equation $px^2 + 2^{2m} = y^p$ has no solution if $p \equiv 1 \pmod{4}$. Let $p \equiv 3 \pmod{4}$, since $p > 3$, therefore the equation has no solution in odd integers from [Theorem 1.1\(iii\)](#). If x is even then we have

$$pX^2 + 2^{2(m-u)} = Y^p, \quad (1.28)$$

where X is odd. Equation (1.28) has no solution in odd integers from the first part. $\square$

Brown [[2](#), Theorem 3] considered the Diophantine equation (1.1) when $A = 3$, but he did not solve it completely. In the following we give the complete solution.

THEOREM 1.8. *The Diophantine equation $3x^2 + 2^{2m} = y^p$ has a solution only if $m \equiv -1 \pmod{p}$, and this solution is given by $x = 2^m$, $y = 2^{(2m+2)/p}$.*

PROOF. Now $A = 3$ and the field $\mathbb{Q}(\sqrt{-3})$ is a unique prime factorization domain, so from [Theorem 1.1](#) this equation has no solution for all p if x is odd. If x is even then from [Theorem 1.5](#) we have $x = 2^m$, $y = 2^{(2m+2)/p}$. Also the equation

$$3X^2 + 2^{2(m-u)} = Y^p, \quad (1.29)$$

where X is odd, has no solution from the first part of this proof. $\square$

Combining the two last theorems we can generalize Brown's result [2] for any odd prime p as follows.

THEOREM 1.9. *The Diophantine equation $px^2 + 2^{2m} = y^p$, where $p \not\equiv 7 \pmod{8}$, has a solution only if $p = 3$ and $m \equiv 2 \pmod{3}$ and this solution is given by $x = 2^m$, $y = 2^{(2m+2)/3}$.*

Considering (1.16) modulo 8 it is easy to prove the following.

COROLLARY 1.10. *For a given A , in (1.16) where $m \geq 2$, if $A \equiv 1 \pmod{8}$ then $p \equiv 7 \pmod{8}$ and if $A \equiv 5 \pmod{8}$ then $p \equiv 3 \pmod{8}$.*

As a special case we consider $A = q$ an odd prime and prove the following theorem.

THEOREM 1.11. *The Diophantine equation $qx^2 + 2^{2m} = y^3$, where $q \equiv 1 \pmod{4}$ is a prime integer and $(3, h) = 1$, has a solution only if $q = 5$ and $m = 2 + 3M$, and the unique solution is given by $x = 43 \cdot 2^{3M}$ and $y = 21 \cdot 2^{2M}$.*

PROOF. First suppose that x is odd, since $q \equiv 1 \pmod{4}$ and $p = 3$, therefore it is sufficient to consider (1.16), then $y = 2^{2m} + qb^2$ and

$$3qb^2 = 2^{2m} - 1 = (2^m - 1)(2^m + 1). \quad (1.30)$$

From [5] it is sufficient to consider $m \geq 2$ and from Corollary 1.10 we have $q \equiv 5 \pmod{8}$. Now $(2^m - 1, 2^m + 1) = 1$, let $b = cd$, where $(c, d) = 1$ and both c and d are odd, then from (1.30) we have only the following possibilities:

(1) $2^m - 1 = 3qc^2$, $2^m + 1 = d^2$, subtracting these two equations, we get $2 = d^2 - 3qc^2$ which is not possible modulo 3.

(2) $2^m - 1 = 3c^2$, $2^m + 1 = qd^2$, considering the first equation modulo 8, we get $m = 2$ and hence $q = 5$. Therefore $y = 2^{2m} + qb^2 = 2^4 + 5(1) = 21$ and so $x = 43$.

(3) $2^m - 1 = d^2$, $2^m + 1 = 3qc^2$, again considering the first equation modulo 8, we get $m = 1$ and then $q = 1$ which is not our case.

(4) $2^m - 1 = qc^2$, $2^m + 1 = 3d^2$, considering the first equation modulo 8, we get a contradiction.

Now suppose that x is even, then we have only the following equation:

$$qX^2 + 2^{2(m-u)} = Y^3, \quad (1.31)$$

where $x = 2^u X$, $y = 2^v \cdot Y$, $m > u > 0$, $v > 0$, $(2, X) = (2, Y) = 1$, and $2u = 3v$. From the first part of this proof, equation (1.31) has a unique solution given by $q = 5$, $m - u = 2$, $X = 43$, and $Y = 21$. Since $2u = 3v$ we get $3 \mid u$, let $u = 3M$ then $m = 2 + 3M$ and $v = 2M$. Hence $x = 43 \cdot 2^{3M}$ and $y = 21 \cdot 2^{2M}$. $\square$

We are unable to solve (1.1) completely when $A \equiv 1 \pmod{4}$ but we are able to solve it for many particular values of A for all p as we will show in the following example. But before this we give a corollary which will help us.

COROLLARY 1.12. *If m is odd then the Diophantine equation (1.1) has no solution in x odd when $5 \mid A$.*

PROOF. Since m is odd, therefore from the proof of [Theorem 1.1](#), it is sufficient to consider (1.16), where $p \equiv 3 \pmod{4}$. If $5 \mid A$ in (1.16), then we get $1 \equiv 2^{m(p-1)} \pmod{5}$ which implies that $4 \mid m(p-1)$ and this is not possible. $\square$

EXAMPLE 1.13. Consider the Diophantine equation $5x^2 + 2^{10} = y^p$.

Here $m = 5$, $A = 5$, $h = 2$, so from [Corollary 1.12](#), this equation has no solution in x odd for all p . If x is even then it is sufficient to consider the equation

$$5X^2 + 2^{2(5-u)} = Y^p, \quad (1.32)$$

where $x = 2^u X$, $y = 2^v \cdot Y$, $5 > u > 0$, $v > 0$, $(2, X) = (2, Y) = 1$, and $2u = pv$. Since p is an odd prime, the only possibility is $u = 3$, $p = 3$, and (1.32) becomes $5X^2 + 2^4 = Y^3$, which has a unique solution from [Theorem 1.11](#), given by $X = 43$ and $Y = 21$, so the given equation has a unique solution, $x = 8.43$, $y = 4.21$, and $p = 3$.

By using the method similar to [3, Lemma 3] we can prove the following lemma.

LEMMA 1.14. *If q is any odd prime which divides the integer b defined in (1.16), then*

$$2^{m(q-1)} \equiv 1 \pmod{q^2}. \quad (1.33)$$

Considering (1.16) modulo 3 we are able to prove the following theorem.

THEOREM 1.15. *If $3 \mid b$ in (1.16), then $m = 3^k \cdot m'$, where $k \geq 1$, $(3, m') = 1$ and either*

- (1) $3 \nmid A$ and then there is no solution if k even, or
- (2) $3 \mid A$ and then there is no solution if k odd.

PROOF. Let $3 \mid b$ then from [Lemma 1.14](#), $2^{2m} \equiv 1 \pmod{9}$ which implies that $3 \mid m$. Let $m = 3^k \cdot m'$, where $(3, m') = 1$, $k \geq 1$. Since $p \equiv 3 \pmod{4}$, put $p-1 = 2 \cdot 3^t \cdot p'$, where $(2, p') = (3, p') = 1$, $t \geq 0$ and put $b = 3^s \cdot b'$, where $(3, b') = 1$, $s \geq 1$. Rewrite (1.16) as

$$1 - 2^{m(p-1)} = \sum_{r=1}^{(p-1)/2} \binom{p}{2r} 2^{m(p-2r-1)} (-Ab^2)^r. \quad (1.34)$$

The general term in the right-hand side is

$$\binom{p}{2r} 2^{m(p-2r-1)} (-Ab^2)^r = \binom{p-2}{2r-2} 2^{m(p-2r-1)} \times \frac{pb^{2r-2}}{r(2r-1)} \cdot b^2 (-A)^r. \quad (1.35)$$

Since $3^{2r-2} \geq r(2r-1)$ for $r \geq 1$, this right-hand side is divisible at least by 3^{2s+t} if $(3, A) = 1$, so from (1.34) we get

$$2^{m(p-1)} \equiv 1 \pmod{3^{2s+t}}. \quad (1.36)$$

Since 2 is a primitive root of 3^{2s+t} , therefore $\phi(3^{2s+t}) \mid m(p-1)$ which implies that $3^{2s+t-1} \mid 2 \cdot 3^k m' \cdot 3^t p'$, hence $3^{2s-k-1} \mid m' p'$. But $(3, m') = (3, p') = 1$, so $2s-k-1 = 0$ which implies that k is odd.

Now if $3 \mid A$, then the right-hand side in (1.34) is divisible at least by 3^{2s+t+1} and as above we get $k = 2s$, implying k even. $\square$

We are unable to solve (1.1) completely when $p = 7$, but as a special case we prove the following theorem.

THEOREM 1.16. *The Diophantine equation (1.1), where $(7, h) = 1$, has no solution in x odd when $p = 7$, $A \equiv 1 \pmod{12}$, and $m = 3^{2k} \cdot m'$, where $k \geq 1$, $(3, m') = 1$.*

PROOF. Here $p = 7$, so from Theorem 1.1(iii) we get $A \equiv 1 \pmod{4}$. Put $p = 7$ in (1.16), then

$$1 = 2^{6m} - 21Ab^22^{4m} + 35A^2b^42^{2m} - 7A^3b^6. \quad (1.37)$$

If $3 \mid b$ in (1.37) then from Theorem 1.15(1), this equation has no solution. So $(3, b) = 1$, and then considering (1.37) modulo 3 we get

$$2A^2 - A^3 \equiv 0 \pmod{3} \quad (1.38)$$

which is not true since $A \equiv 1 \pmod{3}$. $\square$

THEOREM 1.17. *If $(3, m) = 1$, then the Diophantine equation (1.1), where $(p, h) = 1$ has no solution in x odd when $A \equiv 1 \pmod{24}$.*

PROOF. The case $m = 1$, the Diophantine equation (1.1) has no solution [5]. Let $m \geq 2$, since $A \equiv 1 \pmod{8}$ then from Corollary 1.10, $p = 7 + 8H$. Since $(3, m) = 1$, then from Theorem 1.15, $(3, b) = 1$ so $b^2 \equiv 1 \pmod{3}$. Considering (1.16) modulo 3, where $A \equiv 1 \pmod{3}$ we get

$$1 \equiv \sum_{r=0}^{(p-1)/2} \binom{p}{2r} (-1)^r \pmod{3} \equiv \frac{(1+i)^p + (1-i)^p}{2} \pmod{3}. \quad (1.39)$$

But $(1 \pm i)^8 \equiv 1 \pmod{3}$, so (1.39) implies that

$$\begin{aligned} 1 &\equiv \frac{\{(1+i)^{8(1+H)}(1-i) + (1-i)^{8(1+H)}(1+i)\}}{2(1+i)(1-i)} \pmod{3} \\ &\equiv 4^{1+H} \times \frac{1}{2} \pmod{3} \\ &\equiv 2 \pmod{3} \end{aligned} \quad (1.40)$$

which is a contradiction. $\square$

EXAMPLE 1.18. Consider the Diophantine equation $73x^2 + 2^{14} = y^p$.

Here $m = 7$, $A = 73$, $h = 4$ so from Theorem 1.17, this equation has no solution in x odd. If x is even then it is sufficient to consider the equation

$$73X^2 + 2^{2(7-u)} = Y^p, \quad (1.41)$$

where $x = 2^u X$, $y = 2^v \cdot Y$, $7 > u > 0$, $v > 0$, $(2, X) = (2, Y) = 1$, and $2u = pv$. If $(7-u, 3) = 1$, then (1.41) has no solution from Theorem 1.17. If $3 \mid 7-u$ then $u = 1$ or 4 , which is not possible since $2u = pv$. So the given equation has no solution in integers.

ACKNOWLEDGEMENT. The author would like to thank the referee for his valuable suggestions.

REFERENCES

- [1] S. A. Arif and F. S. Abu Muriefah, *On the Diophantine equation $x^2 + 2^k = y^n$* , Int. J. Math. Math. Sci. **20** (1997), no. 2, 299–304. [MR 98b:11032](#). [Zbl 881.11038](#).
- [2] E. Brown, *Diophantine equations of the form $ax^2 + Db^2 = y^p$* , J. Reine Angew. Math. **291** (1977), 118–127. [MR 55#12631](#). [Zbl 338.10018](#).
- [3] J. H. E. Cohn, *The Diophantine equation $x^2 + C = y^n$* , Acta Arith. **65** (1993), no. 4, 367–381. [MR 94k:11037](#). [Zbl 795.11016](#).
- [4] W. Ljunggren, *Über die Gleichungen $1 + Dx^2 = 2y^n$ und $1 + Dx^2 = 4y^n$* , Norske Vid. Selsk. Forh. (Trondhjem) **15** (1943), no. 30, 115–118 (German). [MR 8,442g](#). [Zbl 028.34604](#).
- [5] T. Nagell, *Contributions to the theory of a category of Diophantine equations of the second degree with two unknowns*, Nova Acta Soc. Sci. Upsal. (4) **16** (1954), no. 2, 38 pp. [MR 17,13b](#). [Zbl 057.28304](#).

FADWA S. ABU MURIEFAH: GIRLS COLLEGE OF EDUCATION, SCIENCE SECTIONS (MATHEMATICS),
SITTEN STREET, AL MALAZ, P.O. BOX 27104, RIYADH, SAUDI ARABIA
E-mail address: abumuriefah@yahoo.com