ON CERTAIN ANALYTIC UNIVALENT FUNCTIONS

B. A. FRASIN and M. DARUS

(Received 10 March 2000)

ABSTRACT. We consider the class of analytic functions $B(\alpha)$ to investigate some properties for this class. The angular estimates of functions in the class $B(\alpha)$ are obtained. Finally, we derive some interesting conditions for the class of strongly starlike and strongly convex of order α in the open unit disk.

2000 Mathematics Subject Classification. Primary 30C45.

1. Introduction. Let $\mathbb{A}$ denote the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1.1)$$

which are analytic in the open unit disk $U = \{z : |z| < 1\}$. A function $f(z)$ belonging to $\mathbb{A}$ is said to be starlike of order α if it satisfies

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > \alpha \quad (z \in U) \quad (1.2)$$

for some α ($0 \leq \alpha < 1$). We denote by S_{α}^* the subclass of $\mathbb{A}$ consisting of functions which are starlike of order α in U . Also, a function $f(z)$ belonging to $\mathbb{A}$ is said to be convex of order α if it satisfies

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > \alpha \quad (z \in U) \quad (1.3)$$

for some α ($0 \leq \alpha < 1$). We denote by C_{α} the subclass of $\mathbb{A}$ consisting of functions which are convex of order α in U .

If $f(z) \in \mathbb{A}$ satisfies

$$\left| \arg \left(\frac{zf'(z)}{f(z)} \right) \right| < \frac{\pi}{2} \alpha \quad (z \in U) \quad (1.4)$$

for some α ($0 \leq \alpha < 1$), then $f(z)$ said to be strongly starlike of order α in U , and this class denoted by $\tilde{S}_{\alpha}^*$.

If $f(z) \in \mathbb{A}$ satisfies

$$\left| \arg \left(1 + \frac{zf''(z)}{f'(z)} \right) \right| < \frac{\pi}{2} \alpha \quad (z \in U) \quad (1.5)$$

for some α ($0 \leq \alpha < 1$), then we say that $f(z)$ is strongly convex of order α in U , and we denote by $\tilde{C}_{\alpha}$ the class of all such functions.

The object of the present paper is to investigate various properties of the following class of analytic functions defined as follows.

DEFINITION 1.1. A function $f(z) \in \mathbb{A}$ is said to be a member of the class $B(\alpha)$ if and only if

$$\left| \frac{z^2 f'(z)}{f^2(z)} - 1 \right| < 1 - \alpha \quad (1.6)$$

for some α ($0 \leq \alpha < 1$) and for all $z \in U$.

Note that condition (1.6) implies

$$\operatorname{Re} \left(\frac{z^2 f'(z)}{f^2(z)} \right) > \alpha. \quad (1.7)$$

2. Main results. In order to derive our main results, we have to recall here the following lemmas.

LEMMA 2.1 (see [2]). *Let $f(z) \in \mathbb{A}$ satisfy the condition*

$$\left| \frac{z^2 f'(z)}{f^2(z)} - 1 \right| < 1 \quad (z \in U), \quad (2.1)$$

then f is univalent in U .

LEMMA 2.2 (see [1]). *Let $w(z)$ be analytic in U and such that $w(0) = 0$. Then if $|w(z)|$ attains its maximum value on circle $|z| = r < 1$ at a point $z_0 \in U$, we have*

$$z_0 w'(z_0) = k w(z_0), \quad (2.2)$$

where $k \geq 1$ is a real number.

LEMMA 2.3 (see [3]). *Let a function $p(z)$ be analytic in U , $p(0) = 1$, and $p(z) \neq 0$ ($z \in U$). If there exists a point $z_0 \in U$ such that*

$$|\arg(p(z))| < \frac{\pi}{2} \alpha, \quad \text{for } |z| < |z_0|, \quad |\arg(p(z_0))| = \frac{\pi}{2} \alpha, \quad (2.3)$$

with $0 < \alpha \leq 1$, then we have

$$\frac{z_0 p'(z_0)}{p(z_0)} = i k \alpha, \quad (2.4)$$

where

$$\begin{aligned} k &\geq \frac{1}{2} \left(a + \frac{1}{a} \right) \geq 1 \quad \text{when } \arg(p(z_0)) = \frac{\pi}{2} \alpha, \\ k &\leq -\frac{1}{2} \left(a + \frac{1}{a} \right) \leq -1 \quad \text{when } \arg(p(z_0)) = -\frac{\pi}{2} \alpha, \\ p(z_0)^{1/\alpha} &= \pm a i, \quad (a > 0). \end{aligned} \quad (2.5)$$

We begin with the statement and the proof of the following result.

THEOREM 2.4. *If $f(z) \in \mathbb{A}$ satisfies*

$$\left| \frac{(zf(z))''}{f'(z)} - \frac{2zf'(z)}{f(z)} \right| < \frac{1-\alpha}{2-\alpha} \quad (z \in U), \quad (2.6)$$

for some α ($0 \leq \alpha < 1$), then $f(z) \in B(\alpha)$.

PROOF. We define the function $w(z)$ by

$$\frac{z^2 f'(z)}{f^2(z)} = 1 + (1-\alpha)w(z). \quad (2.7)$$

Then $w(z)$ is analytic in U and $w(0) = 0$. By the logarithmic differentiations, we get from (2.7) that

$$\frac{(zf(z))''}{f'(z)} - \frac{2zf'(z)}{f(z)} = \frac{(1-\alpha)zw'(z)}{1+(1-\alpha)w(z)}. \quad (2.8)$$

Suppose there exists $z_0 \in U$ such that

$$\max_{|z| < |z_0|} |w(z)| = |w(z_0)| = 1, \quad (2.9)$$

then from Lemma 2.2, we have (2.2). □

Letting $w(z_0) = e^{i\theta}$, from (2.8), we have

$$\left| \frac{(z_0 f(z_0))''}{f'(z_0)} - \frac{2z_0 f'(z_0)}{f(z_0)} \right| = \left| \frac{(1-\alpha)ke^{i\theta}}{1+(1-\alpha)e^{i\theta}} \right| \geq \frac{1-\alpha}{2-\alpha}, \quad (2.10)$$

which contradicts our assumption (2.6). Therefore $|w(z)| < 1$ holds for all $z \in U$. We finally have

$$\left| \frac{z^2 f'(z)}{f^2(z)} - 1 \right| = (1-\alpha) |w(z)| < 1-\alpha \quad (z \in U), \quad (2.11)$$

that is, $f(z) \in B(\alpha)$.

Taking $\alpha = 0$ in Theorem 2.4 and using Lemma 2.1 we have the following corollary.

COROLLARY 2.5. *If $f(z) \in \mathbb{A}$ satisfies*

$$\left| \frac{(zf(z))''}{f'(z)} - \frac{2zf'(z)}{f(z)} \right| < \frac{1}{2} \quad (z \in U), \quad (2.12)$$

then f is univalent in U .

Next, we prove the following theorem.

THEOREM 2.6. *Let $f(z) \in \mathbb{A}$. If $f(z) \in B(\alpha)$, then*

$$\left| \arg \left(\frac{f(z)}{z} \right) \right| < \frac{\pi}{2} \alpha \quad (z \in U), \quad (2.13)$$

for some α ($0 < \alpha < 1$) and $(2/\pi) \tan^{-1} \alpha - \alpha = 1$.

PROOF. We define the function $p(z)$ by

$$\frac{f(z)}{z} = p(z) = 1 + \sum_{n=2}^{\infty} a_n z^{n-1}. \quad (2.14)$$

Then we see that $p(z)$ is analytic in U , $p(0) = 1$, and $p(z) \neq 0$ ($z \in U$). It follows from (2.14) that

$$\frac{z^2 f'(z)}{f^2(z)} = \frac{1}{p(z)} \left(1 + \frac{z p'(z)}{p(z)} \right). \quad (2.15)$$

Suppose there exists a point $z_0 \in U$ such that

$$|\arg(p(z))| < \frac{\pi}{2} \alpha, \quad \text{for } |z| < |z_0|, \quad |\arg(p(z_0))| = \frac{\pi}{2} \alpha. \quad (2.16)$$

Then, applying Lemma 2.3, we can write that

$$\frac{z_0 p'(z_0)}{p(z_0)} = i k \alpha, \quad (2.17)$$

where

$$\begin{aligned} k &\geq \frac{1}{2} \left(a + \frac{1}{a} \right) \geq 1 \quad \text{when } \arg(p(z_0)) = \frac{\pi}{2} \alpha, \\ k &\leq -\frac{1}{2} \left(a + \frac{1}{a} \right) \leq -1 \quad \text{when } \arg(p(z_0)) = -\frac{\pi}{2} \alpha, \\ p(z_0)^{1/\alpha} &= \pm a i, \quad (a > 0). \end{aligned} \quad (2.18)$$

Therefore, if $\arg(p(z_0)) = \pi \alpha / 2$, then

$$\frac{z_0^2 f'(z_0)}{f^2(z_0)} = \frac{1}{p(z_0)} \left(1 + \frac{z_0 p'(z_0)}{p(z_0)} \right) = a^{-\alpha} e^{-i\pi \alpha / 2} (1 + i k \alpha). \quad (2.19)$$

This implies that

$$\begin{aligned} \arg \left(\frac{z_0^2 f'(z_0)}{f^2(z_0)} \right) &= \arg \left(\frac{1}{p(z_0)} \left(1 + \frac{z_0 p'(z_0)}{p(z_0)} \right) \right) \\ &= -\frac{\pi}{2} \alpha + \arg(1 + i k \alpha) \geq -\frac{\pi}{2} \alpha + \tan^{-1} \alpha \\ &= \frac{\pi}{2} \left(\frac{2}{\pi} \tan^{-1} \alpha - \alpha \right) = \frac{\pi}{2} \end{aligned} \quad (2.20)$$

if

$$\frac{2}{\pi} \tan^{-1} \alpha - \alpha = 1. \quad (2.21)$$

Also, if $\arg(p(z_0)) = -\pi \alpha / 2$, we have

$$\arg \left(\frac{z_0^2 f'(z_0)}{f^2(z_0)} \right) \leq -\frac{\pi}{2} \quad (2.22)$$

if

$$\frac{2}{\pi} \tan^{-1} \alpha - \alpha = 1. \quad (2.23)$$

These contradict the assumption of the theorem.

Thus, the function $p(z)$ has to satisfy

$$|\arg(p(z))| < \frac{\pi}{2}\alpha \quad (z \in U) \quad (2.24)$$

or

$$\left| \arg\left(\frac{f(z)}{z}\right) \right| < \frac{\pi}{2}\alpha \quad (z \in U). \quad (2.25)$$

This completes the proof. $\square$

Now, we prove the following theorem.

THEOREM 2.7. *Let $p(z)$ be analytic in U , $p(z) \neq 0$ in U and suppose that*

$$\left| \arg\left(p(z) + \frac{z^3 f'(z)}{f^2(z)} p'(z)\right) \right| < \frac{\pi}{2}\alpha \quad (z \in U), \quad (2.26)$$

where $0 < \alpha < 1$ and $f(z) \in B(\alpha)$, then we have

$$|\arg(p(z))| < \frac{\pi}{2}\alpha \quad (z \in U). \quad (2.27)$$

PROOF. Suppose there exists a point $z_0 \in U$ such that

$$|\arg(p(z))| < \frac{\pi}{2}\alpha, \quad \text{for } |z| < |z_0|, \quad |\arg(p(z_0))| = \frac{\pi}{2}\alpha. \quad (2.28)$$

Then, applying Lemma 2.3, we can write that

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik\alpha, \quad (2.29)$$

where

$$\begin{aligned} k &\geq \frac{1}{2}\left(a + \frac{1}{a}\right) \quad \text{when } \arg(p(z_0)) = \frac{\pi}{2}\alpha, \\ k &\leq -\frac{1}{2}\left(a + \frac{1}{a}\right) \quad \text{when } \arg(p(z_0)) = -\frac{\pi}{2}\alpha, \\ p(z_0)^{1/\alpha} &= \pm ai, \quad (a > 0). \end{aligned} \quad (2.30)$$

Then it follows that

$$\begin{aligned} \arg\left(p(z_0) + \frac{z_0^3 f'(z_0)}{f^2(z_0)} p'(z_0)\right) &= \arg\left(p(z_0) \left(1 + \frac{z_0^2 f'(z_0)}{f^2(z_0)} \frac{z p'(z_0)}{p(z_0)}\right)\right) \\ &= \arg\left(p(z_0) \left(1 + i \frac{z_0^2 f'(z_0)}{f^2(z_0)} \alpha k\right)\right). \end{aligned} \quad (2.31)$$

When $\arg(p(z_0)) = \pi\alpha/2$, we have

$$\arg\left(p(z_0) + \frac{z_0^3 f'(z_0)}{f^2(z_0)} p'(z_0)\right) = \arg(p(z_0)) + \arg\left(1 + i \frac{z_0^2 f'(z_0)}{f^2(z_0)} \alpha k\right) > \frac{\pi}{2}\alpha, \quad (2.32)$$

because

$$\operatorname{Re} \frac{z_0^2 f'(z_0)}{f^2(z_0)} \alpha k > 0 \quad \text{and therefore } \arg\left(1 + i \frac{z_0^2 f'(z_0)}{f^2(z_0)} \alpha k\right) > 0. \quad (2.33)$$

Similarly, if $\arg(p(z_0)) = -\pi\alpha/2$, then we obtain that

$$\arg\left(p(z_0) + \frac{z_0^3 f'(z_0)}{f^2(z_0)} p'(z_0)\right) = \arg(p(z_0)) + \arg\left(1 + i \frac{z_0^2 f'(z_0)}{f^2(z_0)} \alpha k\right) < -\frac{\pi}{2} \alpha, \quad (2.34)$$

because

$$\operatorname{Re} \frac{z_0^2 f'(z_0)}{f^2(z_0)} \alpha k < 0 \text{ and therefore } \arg\left(1 + i \frac{z_0^2 f'(z_0)}{f^2(z_0)} \alpha k\right) < 0. \quad (2.35)$$

Thus we see that (2.32) and (2.34) contradict our condition (2.26). Consequently, we conclude that

$$|\arg(p(z))| < \frac{\pi}{2} \alpha \quad (z \in U). \quad (2.36) \quad \square$$

Taking $p(z) = z f'(z) / f(z)$ in Theorem 2.7, we have the following corollary.

COROLLARY 2.8. *If $f(z) \in \mathbb{A}$ satisfying*

$$\left| \arg\left(\frac{z f'(z)}{f(z)} + \frac{z^3 f'(z)}{f^3(z)} \left((z f'(z))' - \frac{z(f'(z))^2}{f(z)}\right)\right) \right| < \frac{\pi}{2} \alpha \quad (z \in U), \quad (2.37)$$

where $0 < \alpha < 1$ and $f(z) \in B(\alpha)$, then $f(z) \in \tilde{S}_\alpha^*$.

Taking $p(z) = 1 + z f''(z) / f'(z)$ in Theorem 2.7, we have the following corollary.

COROLLARY 2.9. *If $f(z) \in \mathbb{A}$ satisfying*

$$\left| \arg\left(\frac{(z f'(z))'}{f'(z)} + \frac{z^3}{f^3(z)} \left((z f''(z))' - \frac{z(f''(z))^2}{f'(z)}\right)\right) \right| < \frac{\pi}{2} \alpha, \quad (2.38)$$

where $0 < \alpha < 1$ and $f(z) \in B(\alpha)$, then $f(z) \in \tilde{C}_\alpha$.

REFERENCES

- [1] I. S. Jack, *Functions starlike and convex of order α* , J. London Math. Soc. (2) **3** (1971), 469–474. MR 43#7611. Zbl 224.30026.
- [2] M. Nunokawa, *On some angular estimates of analytic functions*, Math. Japon. **41** (1995), no. 2, 447–452. MR 96c:30013. Zbl 822.30014.
- [3] S. Ozaki and M. Nunokawa, *The Schwarzian derivative and univalent functions*, Proc. Amer. Math. Soc. **33** (1972), 392–394. MR 45#8821. Zbl 233.30011.

B. A. FRASIN: SCHOOL OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCES AND TECHNOLOGY, UKM, BANGI 43600 SELANGOR, MALAYSIA

M. DARUS: SCHOOL OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCES AND TECHNOLOGY, UKM, BANGI 43600 SELANGOR, MALAYSIA

E-mail address: maslina@pkrisc.cc.ukm.my