IJMMS 25:3 (2001) 213–216
PII. S0161171201004744
http://ijmms.hindawi.com


# STRONG UNIQUE CONTINUATION OF EIGENFUNCTIONS FOR $p$-LAPLACIAN OPERATOR

**ISLAM EDDINE HADI and N. TSOULI**

(Received 28 January 2000)

ABSTRACT. We show the strong unique continuation property of the eigenfunctions for $p$-Laplacian operator in the case $p < N$.

2000 Mathematics Subject Classification. Primary 35J15.

**1. Introduction.** This paper is primarily concerned with the problem:

$$-\operatorname{div}\left(|\nabla u|^{p-2}\nabla u\right) + V|u|^{p-2}u = 0 \quad \text{in } \Omega, \tag{1.1}$$

where $\Omega$ is a bounded domain in $\mathbb{R}^N$ and the weight functions $V$ is assumed to be not equivalent to zero and to lie in $L^{N/p}(\Omega)$.

Also, we know that the unique continuation property is defined by a different form. In this work, we are interested to study a family of functions which enjoys the strong unique continuation property (SUCP), that is, functions besides possibly the zero functions has a zero of infinite order.

**DEFINITION 1.1.** A function $u \in L^p(\Omega)$ has a zero of infinite order in $p$-mean at $x_0 \in \Omega$, if for each $n \in \mathbb{N}$,

$$\int_{|x-x_0|\le R} |u|^p = 0\left(R^n\right) \quad \text{as } R \longrightarrow 0. \tag{1.2}$$

There is an extensive literature on unique continuation. We refer to the work of Jerison-Kenig on the unique continuation for Shrödinger operators (cf. [3]). The same work is done by Gossez and Figueiredo, but for linear elliptic operator in the case $V \in L^{N/2}$, where $N > 2$, (cf. [1]). Also, Loulit extended this property to $N = 2$ by introducing Orlicz's space, (cf. [2, 5]). In this work, we generalize this property for the $p$-Laplacian in the case $V \in L^{N/p}(\Omega)$ and $p < N$.

**2. Strong unique continuation theorem.** In this section, we proceed to establish the strong unique continuation property of the eigenfunctions for the $p$-Laplacian operator in the case $V \in L^{N/p}(\Omega)$ and $p < N$.

**THEOREM 2.1.** *Let $u \in W^{1,p}_{\mathrm{loc}}(\Omega)$ solution of (1.1). If $u = 0$ on a set $E$ of positive measure, then $u$ has a zero of infinite order in $p$-mean.*

To prove this theorem we need the following lemmas.

**Lemma 2.2.** *Let $g \in W_0^{1,p}(\Omega)$ and $V \in L^{N/p}$. Then for each $\epsilon > 0$ there exists a positive constant $k_\epsilon$ such that*

$$\int_\Omega V|g|^p \le \epsilon \int_\Omega |\nabla g|^p + k_\epsilon \int_\Omega |g|^p. \tag{2.1}$$

**Proof.** Set

$$G = \left\{ x \in \frac{\Omega}{V(x)} \ge c \right\}. \tag{2.2}$$

So

$$\int_\Omega V|g|^p \le \int_G V|g|^p + k \int_\Omega |g|^p. \tag{2.3}$$

By using the Hölder and Poincaré's inequalities, we get

$$\int_\Omega V|g|^p \le c\|\chi_G V\|_L^{N/p} \int_\Omega |\nabla g|^p + k \int_\Omega |g|^p. \tag{2.4}$$

But $\|\cdot\|$ is absolutely continuous. So, given $\epsilon > 0$, there exists $k$ such that $c\|\chi_G V\| \le \epsilon$. Which gives immediately the inequality (2.1). □

**Lemma 2.3.** *Let $B_r$ and $B_{2r}$ be two concentric balls contained in $\Omega$. Then*

$$\int_{B_r} |\nabla u|^p \le \frac{c}{r^p} \int_{B_{2r}} |u|^p, \tag{2.5}$$

*where the constant $c$ does not depend on $r$.*

**Proof.** Take $\varphi \in C_0^\infty(\Omega)$, with $\operatorname{supp}\varphi \subset B_{2r}$, $\varphi(x) = 1$ for $x \in B_r$ and $|\nabla\varphi| \le c/r$. Using $\varphi^p u$ as a test function in (1.1), we get

$$\int_{B_{2r}} -\operatorname{div}(|\nabla u|^{p-2}\nabla u)\varphi^p u + \int_{B_{2r}} V|u|^{p-2} u \varphi^p u = 0. \tag{2.6}$$

So

$$\int_{B_{2r}} |\nabla u|^p \varphi^p = -p \int_{B_{2r}} |\nabla u|^{p-2} \varphi^{p-2} \nabla u \cdot \nabla\varphi (\varphi u) - \int_{B_{2r}} V|\varphi u|^p. \tag{2.7}$$

Using Young's inequalities for $(((p-1)/p) + 1/p = 1)$, we can estimate the first integral in the right-hand side of (2.7) by

$$(p-1)\epsilon^{p/(p-1)} \int_{B_{2r}} |\nabla u|^p \varphi^p + \epsilon^{-p} \int_{B_{2r}} |\nabla\varphi|^p |u|^p. \tag{2.8}$$

Also by the result of Lemma 2.2, we can estimate the second integral in the right-hand side of (2.7) by

$$\epsilon \int_{B_{2r}} |\nabla(\varphi u)|^p + c_\epsilon \int_{B_{2r}} |\varphi u|^p. \tag{2.9}$$

Using these estimates in (2.7), we have

$$\begin{aligned} \int_{B_{2r}} |\nabla u|^p \varphi^p \le{}& ((p-1)\epsilon^{p/(p-1)} + \epsilon) \int_{B_{2r}} |\nabla u|^p |\varphi|^p \\ &+ (\epsilon^{-p} + \epsilon) \int_{B_{2r}} |u|^p |\nabla\varphi|^p + c_\epsilon \int_{B_{2r}} |u|^p |\varphi|^p. \end{aligned} \tag{2.10}$$

Using the fact that $|\nabla\varphi| \le c/r$, $|\varphi| \le c/r$, and $\varphi = 1$ in $B_r$, we have immediately inequality (2.5). □

**Lemma 2.4.** *Let $u \in W^{1,1}(B_r)$, where $B_r$ is the ball of radius $r$ in $R^N$ and let $E = \{x \in B_r : u(x) = 0\}$. Then there exists a constant $\beta$ depending only on $N$ such that*

$$\int_A |u| \le \beta \frac{r^N}{|E|} |A|^{1/N} \int_{B_r} |\nabla u| \tag{2.11}$$

*for all ball $B_r$, $u$ as above and all measurable sets $A \subset B_r$.*

To prove this lemma see [4].

**Proof of Theorem 2.1.** We know that almost every point of $E$ is a point of density of $E$. Let $x_0 \in E$ be such a point. This means that

$$\lim_{r \to 0} \frac{|E \cap B_r|}{|B_r|} = 1, \tag{2.12}$$

where $B_r$ denotes the ball of radius $r$ centered at $x_0$ and $|S|$ denotes the Lebesgue measure of a set $S$. So, given $\epsilon > 0$ there is an $r_0 = r_0(\epsilon)$ such that

$$\frac{|E^c \cap B_r|}{|B_r|} < \epsilon, \qquad \frac{|E \cap B_r|}{|B_r|} > 1 - \epsilon \quad \text{for } r \le r_0, \tag{2.13}$$

where $E^c$ denotes the complement of the set $E$. Taking $r_0$ smaller, if necessary, we can assume $B_{r_0} \subset \Omega$. Since $u = 0$ on $E$, by Lemma 2.4 and (2.13) we have

$$\begin{aligned} \int_{B_r} |u|^p = \int_{B_r \cap E^c} |u|^p &\le \beta \frac{r^N}{|E \cap B_r|} |E^c \cap B_r|^{1/N} \int_{B_r} |\nabla (u)^p| \\ &\le p\beta \frac{r^N}{|B_r|^{(1-1/N)}} \frac{\epsilon^{1/N}}{1-\epsilon} \int_{B_r} |u|^{p-1} |\nabla u|. \end{aligned} \tag{2.14}$$

By Hölder's inequality

$$\int_{B_r} |u|^p \le c \frac{\epsilon^{1/N}}{1-\epsilon} r \left( \int_{B_r} |\nabla u|^p \right)^{1/p} \left( \int_{B_r} |u|^p \right)^{(p-1)/p}, \tag{2.15}$$

and by using the Young's inequality, we get

$$\int_{B_r} |u|^p \le c \frac{\epsilon^{1/N}}{1-\epsilon} r \left( r^{p-1} \int_{B_r} |\nabla u|^p + \frac{p-1}{r} \int_{B_r} |u|^p \right). \tag{2.16}$$

Finally, by Lemma 2.3, we have

$$\int_{B_r} |u|^p \le c \frac{\epsilon^{1/N}}{1-\epsilon} \int_{B_{2r}} |u|^p, \tag{2.17}$$

where $c$ is independent of $\epsilon$ and of $r$ as $r \to 0$.

Now let us introduce the following functions:

$$f(r) = \int_{B_r} |u|^p. \tag{2.18}$$

And let us fix $n \in \mathbb{N}$, choose $\epsilon > 0$ such that $(c\epsilon^{1/N})/(1-\epsilon) \le 2^{-n}$. Observe that consequently $r_0$ depends on $n$. Then (2.17) can be written as

$$f(r) \le 2^{-n} f(2r) \quad \text{for } r \le r_0. \tag{2.19}$$

Iterating (2.19), we get

$$f(\rho) \le 2^{-kn} f(2^k \rho), \quad \text{if } 2^{k-1}\rho \le r_0. \tag{2.20}$$

Now given $0 < r < r_0(n)$ and choose $k \in \mathbb{N}$ such that

$$2^{-k} r_0 \le r \le 2^{-k+1} r_0. \tag{2.21}$$

From (2.20), we obtain

$$f(r) \le 2^{-kn} f(2^k r) \le 2^{-kn} f(2r_0). \tag{2.22}$$

Since $2^{-k} \le r/r_0$, we finally obtain

$$f(r) \le \left(\frac{r}{r_0}\right)^n f(2r_0), \tag{2.23}$$

which shows that $x_0$ is a zero infinite order in $p$-mean. □

## References


[1] D. G. de Figueiredo and J.-P. Gossez, *Strict monotonicity of eigenvalues and unique continuation*, Comm. Partial Differential Equations **17** (1992), no. 1-2, 339–346. MR 93b:35098. Zbl 777.35042.

[2] J.-P. Gossez and A. Loulit, *A note on two notions of unique continuation*, Bull. Soc. Math. Belg. Sér. B **45** (1993), no. 3, 257–268. MR 96k:35034. Zbl 828.35035.

[3] D. Jerison and C. E. Kenig, *Unique continuation and absence of positive eigenvalues for Schrödinger operators.* With an appendix by E. M. Stein, Ann. of Math. (2) **121** (1985), no. 3, 463–494. MR 87a:35058. Zbl 593.35119.

[4] O. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*, Translated from the Russian by Scripta Technica, Inc. Leon Ehrenpreis (trans. ed.), Academic Press, New York, London, 1968. MR 39#5941. Zbl 164.13002.

[5] A. Loulit, *Inégalités avec Poids et Problèmes de Continuation Unique*, Thèse de Doctorat, Université Libre de Bruxelles, 1995 (French).



Islam Eddine Hadi: Département de Mathématiques et Informatique Faculté des Sciences, Université Mohamed I, Oujda, Morocco

*E-mail address*: hadi@sciences.univ-oujda.ac.ma

N. Tsouli: Département de Mathématiques et Informatique Faculté des Sciences, Université Mohamed I, Oujda, Morocco

*E-mail address*: tsouli@sciences.univ-oujda.ac.ma