ON HENSTOCK-DUNFORD AND HENSTOCK-PETTIS INTEGRALS

YE GUOJU and AN TIANQING

(Received 22 October 1998)

ABSTRACT. We give the Riemann-type extensions of Dunford integral and Pettis integral, Henstock-Dunford integral and Henstock-Pettis integral. We discuss the relationships between the Henstock-Dunford integral and Dunford integral, Henstock-Pettis integral and Pettis integral. We prove the Harnack extension theorems and the convergence theorems for Henstock-Dunford and Henstock-Pettis integrals.

2000 Mathematics Subject Classification. Primary 26A39, 28B05; Secondary 28B20, 46G10, 46G12.

1. Introduction. During 1957-1958, R. Henstock and J. Kurzweil, independently, gave a Riemann-type integral called the Henstock-Kurzweil integral (or Henstock integral) (see [7]). It is a kind of nonabsolute integral and contains the Lebesgue integral. It has been proved that this integral is equivalent to the special Denjoy integral [7]. The Dunford, Pettis integrals are generalizations of the Lebesgue integral to Banach-valued functions. In [5], R. A. Gordon gave two Denjoy-type extensions of the Dunford, Pettis integrals, the Denjoy-Dunford and Denjoy-Pettis integrals, and discussed their properties.

In this paper, we give the Riemann-type extensions of Dunford, Pettis integrals, the Henstock-Dunford, Henstock-Pettis integrals, and discuss the relationships between the Henstock-Dunford integral and Dunford integral, Henstock-Pettis integral and Pettis integral. We prove the Harnack extension theorems and the convergence theorems for Henstock-Dunford and Henstock-Pettis integrals.

Throughout this paper, X denotes a real Banach space and X^* its dual. $B(X^*) = \{x^* \in X^* : \|x^*\| \leq 1\}$ is the unit ball in X^* . $I_0 = [a, b]$ is a closed interval in $\mathbb{R}$.

We first give some preliminaries. A partition D of $[a, b]$ is a finite collection of interval-point pairs (I, t) with the intervals nonoverlapping and their union $[a, b]$. Here t is the associated point of I . We write $D = \{(I, t)\}$, it is said to be δ -fine partition of $[a, b]$ if for each interval-point pair (I, t) , we have $t \in I \subset (t - \delta(t), t + \delta(t))$.

DEFINITION 1.1 (see [7]). A function $f : [a, b] \rightarrow \mathbb{R}$ is Henstock integrable if there exists a function $F : [a, b] \rightarrow \mathbb{R}$ such that for every $\epsilon > 0$ there is a function $\delta(t) > 0$ such that for any δ -fine partition $D = \{[u, v]; t\}$ of $[a, b]$, we have

$$\left| \sum [f(t)(v - u) - F(u, v)] \right| < \epsilon, \quad (1.1)$$

where the sum $\sum$ is understood to be over $D = \{([u, v], t)\}$ and $F(u, v) = F(v) - F(u)$. We write $(H) \int_{I_0} f = F(I_0)$.

The function f is said to be Henstock integrable on the set $E \subset [a, b]$ if the function $f\chi_E$ is Henstock integrable on $[a, b]$. We write $(H) \int_{I_0} f\chi_E = (H) \int_E f$.

DEFINITION 1.2 (see [1, 5, 7]). A function $f : [a, b] \rightarrow \mathbb{R}$ is Denjoy (or special Denjoy) integrable if there exists an ACG (or ACG*) function $F : [a, b] \rightarrow \mathbb{R}$ such that $D_{\text{ap}}F(t) = f(t)$ (or $F'(t) = f(t)$) almost everywhere on $[a, b]$. Where $D_{\text{ap}}F(t)$ denotes the approximate derivative of F at t . We write $(D) \int_{I_0} f = F(I_0)$ (or $(D^*) \int_{I_0} f = F(I_0)$).

The function f is said to be Denjoy (or special Denjoy) integrable on the set $E \subset [a, b]$ if the function $f\chi_E$ is Denjoy (or special Denjoy) integrable on $[a, b]$. We write $(D) \int_{I_0} f\chi_E = (D) \int_E f$ (or $(D^*) \int_{I_0} f\chi_E = (D^*) \int_E f$).

If f is special Denjoy integrable, then f is Denjoy integrable.

LEMMA 1.3 (see [7]). A function $f : [a, b] \rightarrow \mathbb{R}$ is Henstock integrable on $[a, b]$ if and only if f is the special Denjoy integrable on $[a, b]$.

DEFINITION 1.4 (see Gordon [5]). (a) A function $f : [a, b] \rightarrow X$ is Denjoy-Dunford integrable on $[a, b]$ if for each x^* in X^* the function x^*f is Denjoy integrable on $[a, b]$ and if for every interval I in $[a, b]$ there exists a vector x_I^{**} in X^{**} such that $x_I^{**}(x^*) = \int_I x^*f$ for all x^* in X^* . We write $x_{I_0}^{**} = (DD) \int_{I_0} f = F(I_0)$ and F is called the primitive of f on I_0 .

(b) A function $f : [a, b] \rightarrow X$ is Denjoy-Pettis integrable on $[a, b]$ if f is Denjoy-Dunford integrable on $[a, b]$ and if $x_I^{**} \in X$ for every interval I in $[a, b]$. We write $x_{I_0}^{**} = (DP) \int_{I_0} f = F(I_0)$ and F is called the primitive of f on I_0 .

The function f is said to be integrable in one of the above senses on the set $E \subset [a, b]$ if the function $f\chi_E$ is integrable in that sense on $[a, b]$.

LEMMA 1.5 (see [3]). A function $f : [a, b] \rightarrow X$ is Denjoy-Dunford integrable on $[a, b]$ if and only if x^*f is Denjoy integrable on $[a, b]$ for all $x^* \in X^*$.

2. Definition and properties. In the following, we give the Riemann-type extensions of Dunford, Pettis integrals, and discuss the relationships between Henstock-Dunford integral and Dunford integral, Henstock-Pettis integral and Pettis integral.

DEFINITION 2.1. (a) A function $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$ if for each x^* in X^* the function x^*f is Henstock integrable on $[a, b]$ and if for every interval I in $[a, b]$ there exists a vector x_I^{**} in X^{**} such that $x_I^{**}(x^*) = \int_I x^*f$ for all x^* in X^* . We write $x_{I_0}^{**} = (HD) \int_{I_0} f = F(I_0)$ and F is called the primitive of f on I_0 .

(b) A function $f : [a, b] \rightarrow X$ is Henstock-Pettis integrable on $[a, b]$ if f is Henstock-Dunford integrable on $[a, b]$ and if $x_I^{**} \in X$ for every interval I in $[a, b]$. We write $x_{I_0}^{**} = (HP) \int_{I_0} f = F(I_0)$ and F is called the primitive of f on I_0 .

The function f is said to be integrable in one of the above senses on the set $E \subset [a, b]$ if the function $f\chi_E$ is integrable in that sense on $[a, b]$.

By the above definitions and Definition 1.4, it is easy to see that if f is Henstock-Dunford (or Henstock-Pettis) integrable on I_0 , then f is Denjoy-Dunford (or Denjoy-Pettis) integrable.

THEOREM 2.2. *A function $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$ if and only if x^*f is Henstock integrable on $[a, b]$ for all $x^* \in X^*$.*

PROOF. If f is Henstock-Dunford integrable on $[a, b]$, for every $x^* \in X^*$, by Definition 2.1, x^*f is Henstock integrable on $[a, b]$. Conversely, if x^*f is Henstock integrable on $[a, b]$. It follows from Lemma 1.3 that x^*f is Denjoy integrable on $[a, b]$ and $(D) \int_a^b x^*f = (H) \int_a^b x^*f$. It follows from Lemma 1.5 that f is Denjoy-Dunford integrable on $[a, b]$, and for every interval I in $[a, b]$ there exists a vector x_I^{**} in X^{**} such that $x_I^{**}(x^*) = (D) \int_I x^*f$ for all x^* in X^* , that is, $x_I^{**}(x^*) = (H) \int_I x^*f$ for all x^* in X^* . So f is Henstock-Dunford integrable on $[a, b]$. $\square$

THEOREM 2.3. *If the function $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$, then each perfect set in $[a, b]$ contains a portion on which f is Dunford integrable.*

PROOF. Since the function $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$, then for each $x^* \in X^*$, x^*f is Henstock integrable on $[a, b]$. It follows from [8] that each perfect set in $[a, b]$ contains a portion on which x^*f is Lebesgue integrable. So f is Dunford integrable on a portion. $\square$

THEOREM 2.4. *If the function $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$, then there is a sequence $\{X_k\}$ of closed subsets such that $X_k \subset X_{k+1}$ for all k , $\cup_{k=1}^{\infty} X_k = [a, b]$, f is Dunford integrable on each X_k and*

$$\lim_{k \rightarrow \infty} (\text{Dunford}) \int_{X_k \cap [a, x]} f(t) dt = (HD) \int_a^x f(t) dt \text{ weakly} \quad (2.1)$$

uniformly on $[a, b]$.

PROOF. It follows from Theorem 2.2 that a function $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$ if and only if x^*f is Henstock integrable on $[a, b]$ for all $x^* \in X^*$. From [8], x^*f is Henstock integrable on $[a, b]$, then there is a sequence $\{X_k\}$ of closed subsets such that $X_k \subset X_{k+1}$ for all k , $\cup_{k=1}^{\infty} X_k = [a, b]$, x^*f is Lebesgue integrable on each X_k and

$$\lim_{k \rightarrow \infty} (L) \int_{X_k \cap [a, x]} x^*f(t) dt = (H) \int_a^x x^*f(t) dt \quad (2.2)$$

uniformly on $[a, b]$ for each $x^* \in X^*$. So we obtain that f is Dunford integrable on each X_k and

$$\lim_{k \rightarrow \infty} (\text{Dunford}) \int_{X_k \cap [a, x]} f(t) dt = (HD) \int_a^x f(t) dt \text{ weakly} \quad (2.3)$$

uniformly on $[a, b]$. $\square$

THEOREM 2.5. *If the function $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$, then there exists a sequence $\{X_k\}$ of closed sets, $\cup_{k=1}^{\infty} X_k = [a, b]$, f is Dunford integrable on each X_k .*

PROOF. Since f is Henstock-Dunford integrable on $[a, b]$, by Definition 2.1, for each $x^* \in X^*$, x^*f is Henstock integrable on $[a, b]$, and for every interval $I \subset [a, b]$,

$\int_I x^* f = x^* \int_I f$, and $F(I) = \int_I f \in X$. Since $x^* f$ is Henstock integrable, then $x^* F$ is ACG^* . So there is a sequence $\{X_k\}$ of closed subsets such that $\cup_{k=1}^\infty X_k = [a, b]$ and $x^* F$ is VB^* on each X_k . From [7, Lemma 6.18], $x^* f$ is Lebesgue integrable on each X_k . So we obtain that f is Dunford integrable on each X_k . $\square$

THEOREM 2.6. *Suppose that X contains no copy of c_0 and $f : [a, b] \rightarrow X$. If the function f is Henstock-Pettis integrable on $[a, b]$, then each perfect set in $[a, b]$ contains a portion on which f is Pettis integrable.*

PROOF. Since the function $f : [a, b] \rightarrow X$ is Henstock-Pettis integrable on $[a, b]$, then f is Denjoy-Pettis integrable on $[a, b]$. It follows from [5, Theorem 38] that each perfect set in $[a, b]$ contains a portion on which f is Pettis integrable.

In the fact, from [3, Theorem 10], we have that if each Henstock-Pettis integrable function defined on $[a, b]$ is Pettis integrable on a portion of every close set, then X does not contain c_0 . $\square$

THEOREM 2.7. *Suppose that X contains no copy of c_0 and $f : [a, b] \rightarrow X$ is a measurable. If the function $f : [a, b] \rightarrow X$ is Henstock-Pettis integrable on $[a, b]$, then there exists a sequence $\{X_k\}$ of closed sets with $X_k \uparrow [a, b]$ such that for each $x^* \in X^*$, f is Pettis integrable on each X_k , and*

$$\lim_{k \rightarrow \infty} (\text{Pettis}) \int_{X_k} f = (HP) \int_a^b f \text{ weakly.} \quad (2.4)$$

PROOF. Since f is Henstock-Pettis integrable on $[a, b]$, then f is Henstock-Dunford integrable on $[a, b]$. By Theorem 2.4, there is a sequence $\{X_k\}$ of closed subsets such that $X_k \subset X_{k+1}$ for all k , $\cup_{k=1}^\infty X_k = [a, b]$, $x^* f$ is Lebesgue integrable on each X_k and

$$\lim_{k \rightarrow \infty} (L) \int_{X_k \cap [a, x]} x^* f(t) dt = (H) \int_a^x x^* f(t) dt \quad (2.5)$$

uniformly on $[a, b]$ for each $x^* \in X^*$. So we obtain that f is Dunford integrable on each X_k . From [2, Theorem 2.5, page 54], f is Pettis integrable on $[a, b]$ and

$$\lim_{k \rightarrow \infty} (\text{Pettis}) \int_{X_k \cap [a, x]} f(t) dt = (HP) \int_a^x f(t) dt \text{ weakly} \quad (2.6)$$

uniformly on each X_k , that is,

$$\lim_{k \rightarrow \infty} (\text{Pettis}) \int_{X_k} f = (HP) \int_a^b f \text{ weakly.} \quad (2.7)$$

$\square$

In Theorem 2.7, if we remove the condition that f is a measurable, then we have the following theorem.

THEOREM 2.8. *Suppose that X contains no copy of c_0 . If the function $f : [a, b] \rightarrow X$ is Henstock-Pettis integrable on $[a, b]$, then there exists a sequence $\{X_k\}$ of closed sets, $\cup_{k=1}^\infty X_k = [a, b]$, f is Pettis integrable on each X_k .*

PROOF. Since f is Henstock-Pettis integrable on $[a, b]$, by Definition 2.1, for each $x^* \in X^*$, $x^* f$ is Henstock integrable on $[a, b]$, and for every interval $I \subset [a, b]$,

$\int_I x^* f = x^* \int_I f$, and $F(I) = \int_I f \in X$. Since $x^* f$ is Henstock integrable, then $x^* F$ is ACG^* . So there is a sequence $\{X_k\}$ of closed subsets such that $\cup_{k=1}^\infty X_k = [a, b]$ and $x^* F$ is VB^* on each X_k . For each $k \in \mathbb{N}$, let $(a, b) - X_k = \cup_{n=1}^\infty (c_n^k, d_n^k)$. Then

$$\sum_{n=1}^\infty \left| x^* \int_{c_n^k}^{d_n^k} f \right| < \infty. \quad (2.8)$$

Since X contains no copy of c_0 , by Bessaga-Pelczynski theorem [2, page 22], $\sum_{n=1}^\infty \int_{c_n^k}^{d_n^k} f$ is unconditionally convergent in norm. Also

$$\sum_{n=1}^\infty \sup_{[a_n^k, b_n^k] \subset [c_n^k, d_n^k]} \left| x^* \int_{a_n^k}^{b_n^k} f \right| < \infty. \quad (2.9)$$

By Harnack extension theorem [7, page 41], we have

$$\int_{X_k} x^* f = \int_a^b x^* f - \sum_{n=1}^\infty \int_{c_n^k}^{d_n^k} x^* f = x^* \left(\int_a^b f - \sum_{n=1}^\infty \int_{c_n^k}^{d_n^k} f \right). \quad (2.10)$$

Hence $\int_{X_k} f = \int_a^b f - \sum_{n=1}^\infty \int_{c_n^k}^{d_n^k} f \in X$ and $\int_{X_k} x^* f = x^* \int_{X_k} f$.

So, for every closed set $H \subset X_k$, we have $\int_H x^* f = x^* \int_H f$ and $\int_H f \in X$. Since $\int_a^b f \chi_{X_k} = \int_{X_k} f \in X$, $\int_a^b f \chi_H = \int_H f \in X$, then for every closed interval $I \subset [a, b]$, $\int_I f \chi_{X_k} = \int_{I \cap X_k} f \in X$. By [5, Theorem 23, page 79], $f \chi_{X_k}$ is Pettis integrable on $[a, b]$, that is, f is Pettis integrable on each X_k . $\square$

3. The extension theorems and convergence theorems. Now we consider the extension theorems and convergence theorems of the Henstock-Dunford and Henstock-Pettis integrals.

THEOREM 3.1. *Let E be a closed subset in $[a, b]$ and $(a, b) - E$ the union of $\{(a_k, b_k)\}$, $k = 1, 2, \dots$. If $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on E and each interval $[a_k, b_k]$ with*

$$\sum_{k=1}^\infty \omega \left(\int_{a_k}^t x^* f, [a_k, b_k] \right) < \infty \quad (3.1)$$

for each $x^ \in X^*$, then f is Henstock-Dunford integrable on $[a, b]$ and*

$$\left\langle x^*, (HD) \int_a^b f \right\rangle = \left\langle x^*, (HD) \int_a^b f \chi_E \right\rangle + \sum_{k=1}^\infty \left\langle x^*, (HD) \int_{a_k}^{b_k} f \right\rangle \quad (3.2)$$

for each $x^ \in X^*$.*

PROOF. From the conditions of Theorem 3.1, we have the function $x^* f$ satisfies the hypothesis of [7, Corollary 7.11]. So we have $x^* f$ is Henstock integrable on $[a, b]$ and

$$(H) \int_a^b x^* f = (H) \int_a^b x^* f \chi_E + \sum_{k=1}^\infty (H) \int_{a_k}^{b_k} x^* f. \quad (3.3)$$

It follows from [Theorem 2.2](#) that f is Henstock-Dunford integrable on $[a, b]$ and the above equality means that

$$\left\langle x^*, (HD) \int_a^b f \right\rangle = \left\langle x^*, (HD) \int_a^b f \chi_E \right\rangle + \sum_{k=1}^{\infty} \left\langle x^*, (HD) \int_{a_k}^{b_k} f \right\rangle \quad (3.4)$$

for each $x^* \in X^*$. $\square$

THEOREM 3.2. *Let E be a closed subset in $[a, b]$ and $\{(a_k, b_k)\}$ be an enumeration of the intervals contiguous to E in (a, b) . Suppose that $f : [a, b] \rightarrow X$ is Henstock-Pettis integrable on E and each interval $[a_k, b_k]$. If $\sum_{k=1}^{\infty} \omega(\int_{a_k}^{b_k} x^* f, [a_k, b_k]) < \infty$ for each $x^* \in X^*$ and the series $\sum_{k=1}^{\infty} (HP) \int_{[a_k, b_k] \cap J} f$ is unconditionally convergent for every subinterval J of $[a, b]$, then f is Henstock-Pettis integrable on $[a, b]$ and*

$$(HP) \int_a^b f = (HP) \int_a^b f \chi_E + \sum_{k=1}^{\infty} (HP) \int_{a_k}^{b_k} f. \quad (3.5)$$

PROOF. From [Theorem 3.1](#), we have the function f is Henstock-Dunford integrable on $[a, b]$ and $(H) \int_a^b x^* f = (H) \int_a^b x^* f \chi_E + \sum_{k=1}^{\infty} (H) \int_{a_k}^{b_k} x^* f$. To show that f is in fact Henstock-Pettis integrable on $[a, b]$. We need to show that $(HD) \int_J f$ belongs to X for each closed interval J in $[a, b]$.

Let $E_0 = E \cap J$. Then E_0 is a closed set. Since $f \chi_E$ is Henstock-Pettis integrable on J , then $f \chi_{E_0}$ is Henstock-Pettis integrable on J , that is, f is Henstock-Pettis integrable on E_0 . And $\{(a_k, b_k) \cap J\}$ is an enumeration of the intervals contiguous to E_0 in J , so f is Henstock-Pettis integrable on them and $\sum_k (HP) \int_{[a_k, b_k] \cap J} f$ is an unconditionally convergent series in X . Now, if we apply [Theorem 3.1](#) to E_0 in J , we get

$$\left\langle x^*, (HD) \int_J f \right\rangle = \left\langle x^*, (HP) \int_J f \chi_{E_0} \right\rangle + \sum_{k=1}^{\infty} \left\langle x^*, (HP) \int_{[a_k, b_k] \cap J} f \right\rangle \quad (3.6)$$

for each $x^* \in X^*$, that is,

$$\left\langle x^*, (HD) \int_J f \right\rangle = \left\langle x^*, (HP) \int_J f \chi_{E_0} + \sum_{k=1}^{\infty} (HP) \int_{[a_k, b_k] \cap J} f \right\rangle \quad (3.7)$$

for each $x^* \in X^*$. We conclude that

$$(HD) \int_J f = (HP) \int_J f \chi_{E_0} + \sum_{k=1}^{\infty} (HP) \int_{[a_k, b_k] \cap J} f. \quad (3.8)$$

Hence, f is Henstock-Pettis integrable on $[a, b]$ and

$$(HP) \int_a^b f = (HP) \int_a^b f \chi_{E_0} + \sum_{k=1}^{\infty} (HP) \int_{[a_k, b_k] \cap J} f. \quad (3.9)$$

$\square$

COROLLARY 3.3. *Suppose that X contains no copy of c_0 . Let E be a closed subset in $[a, b]$ and $\{(a_k, b_k)\}$ be an enumeration of the intervals contiguous to E in (a, b) . Suppose that $f : [a, b] \rightarrow X$ is Henstock-Pettis integrable on E and each interval $[a_k, b_k]$.*

If $\sum_{k=1}^{\infty} \omega(\int_{a_k}^t x^* f, [a_k, b_k]) < \infty$ for each $x^* \in X^*$, then f is Henstock-Pettis integrable on $[a, b]$ and

$$(HP) \int_a^b f = (HP) \int_a^b f \chi_E + \sum_{k=1}^{\infty} (HP) \int_{a_k}^{b_k} f. \quad (3.10)$$

THEOREM 3.4. Suppose that X is weakly sequentially complete and $f : [a, b] \rightarrow X$ is Henstock-Dunford integrable on $[a, b]$. If f is measurable, then f is Henstock-Pettis integrable on $[a, b]$.

PROOF. It is similar to the proof of [5, Theorem 40]. □

LEMMA 3.5 (see [1, 5]). Let Γ be a family of open intervals in (a, b) and suppose that Γ has the following properties:

- (1) if (α, β) and (β, γ) belong to Γ , then (α, γ) belongs to Γ ;
- (2) if (α, β) belong to Γ , then every open interval in (α, β) belongs to Γ ;
- (3) if (α, β) belong to Γ for every interval in $[\alpha, \beta] \subset (c, d)$, then (c, d) belongs to Γ ;
- (4) if all of the intervals contiguous to the perfect set $E \subset [a, b]$ belong to Γ , then there exists an interval I in Γ such that $I \cap E \neq \emptyset$.

Then Γ contains the interval (a, b) .

LEMMA 3.6. Suppose that $f_n : [a, b] \rightarrow \mathbb{R}$, $f : [a, b] \rightarrow \mathbb{R}$, and

- (1) $f_n \rightarrow f$ almost everywhere on $[a, b]$ as $n \rightarrow \infty$, where each f_n is Henstock (or D^*) integrable on $[a, b]$;
 - (2) the primitives F_n of f_n are continuous uniformly in n and ACG^* uniformly in n .
- Then f is Henstock (or D^*) integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f. \quad (3.11)$$

DEFINITION 3.7. Let $F : [a, b] \rightarrow X$ and let E be a subset of $[a, b]$.

(a) F is BV^* on E if $\sup\{\sum_i \omega(F; [c_i, d_i])\}$ is finite, where the supremum is taken over all finite collections $\{[c_i, d_i]\}$ of nonoverlapping intervals that have endpoints in E , ω denotes the oscillation of F over $[c_i, d_i]$, that is,

$$\omega(F; [c_i, d_i]) = \sup\{\|F(x) - F(y)\|; x, y \in [c_i, d_i]\}. \quad (3.12)$$

(b) F is AC^* on E if for each $\epsilon > 0$ there exists $\delta > 0$ such that $\sum_i \omega(F; [c_i, d_i]) < \epsilon$ whenever $\{[c_i, d_i]\}$ is a finite collection of nonoverlapping intervals that have endpoints in E and satisfy $\sum_i (d_i - c_i) < \delta$.

(c) F is BVG^* on E if E can be expressed as a countable union of sets on each of which F is BV^* .

(d) F is ACG^* on E if F is continuous on E and if E can be expressed as a countable union of sets on each of which F is AC^* .

THEOREM 3.8. Suppose that X is weakly sequentially complete and

- (1) $f_n \rightarrow f$ weakly almost everywhere on $[a, b]$ as $n \rightarrow \infty$, where each f_n is Henstock-Pettis integrable on $[a, b]$;
- (2) the primitives F_n of f_n are continuous uniformly in n and ACG^* uniformly in n .

Then f is Henstock-Pettis integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.13)$$

PROOF. Let

$$\Gamma = \left\{ (\alpha, \beta) \subset [a, b] : f \text{ is Henstock-Pettis integrable on } [\alpha, \beta], \int_{\alpha}^{\beta} f_n \rightarrow \int_{\alpha}^{\beta} f \text{ weakly} \right\}. \quad (3.14)$$

We must show that Γ contains (a, b) and by Lemma 3.5 it is sufficient to verify that Γ satisfies Romanovski's four conditions.

Conditions (1) and (2) are easily verified.

Suppose that (α, β) belongs to Γ for every interval $[\alpha, \beta]$ in (c, d) . For each positive integer $n > 2/(d - c)$, define $I_n = (c + 1/n, d - 1/n)$ and let $x_n = x_{I_n}^{**}$.

Then we have

$$x_{(c,d)}^{**}(x^*) = \int_c^d x^* f = \lim_{n \rightarrow \infty} \int_{I_n} x^* f = \lim_{n \rightarrow \infty} x^*(x_n) \quad (3.15)$$

for each x^* in X^* . Since X is weakly sequentially complete, the sequence $\{x_n\}$ converges weakly to an element x_0 of X and we must have $x_{(c,d)}^{**} = x_0$. It follows easily that (c, d) belongs to Γ and this verifies condition (3).

Now let E be a perfect set in $[a, b]$ such that each of the intervals in $[a, b]$ contiguous to E belongs to Γ .

Since $\{F_n\}$ is continuous uniformly in n and ACG^* uniformly in n , then for each $x^* \in X^*$, $\{x^* F_n\}$ is continuous uniformly in n and ACG^* uniformly in n , and $x^* f_n \rightarrow x^* f$ almost everywhere in $[a, b]$. It follows from [1] that $x^* f$ is special Denjoy integrable on $[a, b]$. So there exists an interval $[u, v]$ with $u, v \in E$ and $E \cap (u, v) \neq \emptyset$ such that $\{F_n\}$ is AC^* uniformly in n on $P = E \cap (u, v)$ and the series $\sum_k \omega(F_n; [u_k, v_k])$ unconditionally converges where $(u, v) - E = \cup_k (u_k, v_k)$. Hence $\sum_k \omega(\int_{u_k}^t x^* f_n; [u_k, v_k]) < \infty$ for each $x^* \in X^*$. By Corollary 3.3, we have

$$\int_u^v f_n = \int_P f_n + \sum_k \int_{u_k}^{v_k} f_n. \quad (3.16)$$

$\{F_n\}$ is AC^* uniformly in n on P , $\{x^* F_n : x^* \in B(X^*), n \in \mathbb{N}\}$ is AC^* uniformly in n on P . So $\{x^* f_n : x^* \in B(X^*), n \in \mathbb{N}\}$ is uniformly integrable on P (see [2]), that is, for $E \subset P$,

$$\lim_{|E| \rightarrow 0} \int_E |x^* f_n| = 0 \quad \text{uniformly in } x^* \in B(X^*) \text{ and } n. \quad (3.17)$$

It follows from [4, Theorem 3] that f is Pettis integrable on P and $\int_P f_n \rightarrow \int_P f$ weakly. Since $\{F_n\}$ is AC^* uniformly in n on P , so for every $\epsilon > 0$ there exists N such that $\sum_{k=N}^{\infty} \|\int_{u_k}^{v_k} f_n\| < \epsilon$, $n = 1, 2, \dots$. For every $x^* \in B(X^*)$, we have $\sum_{k=N}^{\infty} |\int_{u_k}^{v_k} x^* f_n| < \epsilon$, $n = 1, 2, \dots$. So $\sum_{k=N}^{\infty} |\int_{u_k}^{v_k} x^* f| < \epsilon$. Since X is weakly sequentially complete and X does not contain c_0 , hence $\sum_k \int_{u_k}^{v_k} f$ unconditionally converges. By (3.16),

$$x^* \int_u^v f_n = x^* \int_P f_n + x^* \sum_k \int_{u_k}^{v_k} f_n. \quad (3.18)$$

Let $n \rightarrow \infty$, we have

$$x_{(u,v)}^{**}(x^*) = x^* \int_P f + x^* \sum_k \int_{u_k}^{v_k} f. \quad (3.19)$$

Hence

$$x_{(u,v)}^{**} = \int_P f + \sum_k \int_{u_k}^{v_k} f \in X, \quad (3.20)$$

that is, f is Henstock-Pettis integrable on $[u, v]$. So $(u, v) \in \Gamma$. This shows that (u, v) belongs to Γ and Γ satisfies condition (4). This completes the proof. $\square$

THEOREM 3.9. Suppose that X is weakly sequentially complete and $f_n \rightarrow f$ weakly almost everywhere on $[a, b]$ as $n \rightarrow \infty$, where each f_n is Henstock-Pettis integrable on $[a, b]$. If there is a scalar function g with $\|f_n(\cdot)\| \leq g(\cdot)$ almost everywhere for all n and if $\int g < \infty$, then f is Henstock-Pettis integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.21)$$

PROOF. It is similar to the proof of Theorem 3.8. $\square$

DEFINITION 3.10. Let $\{f_\alpha\}$ be a family of Henstock-Pettis integrable functions defined on $[a, b]$. The family $\{x^* f_\alpha : x^* \in B(X^*)\}$ is uniformly integrable in the generalized sense on $[a, b]$, if for each perfect set $E \subset [a, b]$ there exists an interval $[c, d] \subset [a, b]$ with $c, d \in E$ and $E \cap (c, d) \neq \emptyset$ such that $\{x^* f_\alpha : x^* \in B(X^*)\}$ is uniformly integrable on $P = E \cap (c, d)$ and for every α the series $\sum_k \int_{c_k}^{d_k} f_\alpha$ is unconditionally convergent where $(c, d) - E = \cup_k (c_k, d_k)$.

THEOREM 3.11. Suppose that X is weakly sequentially complete and

- (1) $f_n \rightarrow f$ weakly almost everywhere on $[a, b]$ as $n \rightarrow \infty$, where each f_n is Henstock-Pettis integrable on $[a, b]$.
- (2) The family $\{x^* f_n : x^* \in B(X^*), n \in \mathbb{N}\}$ is uniformly integrable in the generalized sense on $[a, b]$.
- (3) For each $x^* \in X^*$, $\lim_{n \rightarrow \infty} \int_c^d x^* f_n = \int_c^d x^* f$ uniformly for every $[c, d] \subset [a, b]$. Then f is Henstock-Pettis integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.22)$$

PROOF. It is similar to the proof of Theorem 3.8. The only difference is that the family $\{x^* f_n : x^* \in B(X^*), n \in \mathbb{N}\}$ is uniformly integrable in the generalized sense on $[a, b]$, then there is a portion $P = E \cap I$ of E such that the family $|x^* f_n|_{\chi_E}$ is uniformly integrable on P . So f is Pettis integrable on P . $\square$

THEOREM 3.12. Suppose that X is weakly sequentially complete and

- (1) $f_n \rightarrow f$ weakly almost everywhere on $[a, b]$ as $n \rightarrow \infty$, where each f_n is Henstock-Pettis integrable on $[a, b]$ and f is measurable,
- (2) the primitives F_n of f_n are weakly continuous uniformly in n and weakly ACG^* uniformly in n , that is, for every $x^* \in X^*$, $x^* F_n$ are continuous uniformly in n and ACG^* uniformly in n .

Then f is Henstock-Pettis integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.23)$$

PROOF. For each x^* in X^* , we have

- (1) $x^* f_n \rightarrow x^* f$ almost everywhere on $[a, b]$ as $n \rightarrow \infty$, where each $x^* f_n$ is Henstock integrable on $[a, b]$,
- (2) the primitives $x^* F_n$ of $x^* f_n$ are continuous uniformly in n and ACG^* uniformly in n . It follows from Lemma 3.6 that $x^* f$ is Henstock integrable on $[a, b]$ and

$$\int_a^b x^* f_n \rightarrow \int_a^b x^* f \text{ as } n \rightarrow \infty. \quad (3.24)$$

By Theorem 2.2, f is Henstock-Dunford integrable on $[a, b]$. Since X is weakly sequentially complete and f is measurable, by Theorem 3.4, f is Henstock-Pettis integrable on $[a, b]$. $\square$

THEOREM 3.13. Suppose that the unit ball $B(X^*)$ of X^* is weak* sequentially compact and

- (1) $f_n \rightarrow f$ weakly almost everywhere in $[a, b]$ as $n \rightarrow \infty$, where each f_n is Henstock-Pettis integrable on $[a, b]$,
 - (2) the primitives F_n of f_n are continuous uniformly in n and ACG^* uniformly in n .
- Then f is Henstock-Pettis integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.25)$$

PROOF. Suppose that $I \subset I_0$. Let C be the weak closure of $\{\int_I f_n : n \in \mathbb{N}\}$. For each x^* in X^* , $\{x^* F_n : n \in \mathbb{N}\}$ is continuous uniformly in n and ACG^* uniformly in n in $[a, b]$, and further $\int_a^b x^* f_n = x^* \int_a^b f_n$. A convergence theorem, namely Lemma 3.6, guarantees that $x^* f$ is Henstock integrable on $[a, b]$ and $\lim_{n \rightarrow \infty} \int_a^b x^* f_n = \int_a^b x^* f$ for each x^* in X^* . We observe that C is bounded and that $C - \{\int_I f_n : n \in \mathbb{N}\}$ contains at most one point. We will prove that C is weakly compact.

Suppose that C is not weakly compact. An appeal to a theorem of James [6, Theorem 1] produces a bounded sequence (x_k^*) in X^* , a sequence (x_n) in C , and an $\epsilon > 0$ such that $x_k^*(x_n) = 0$ for $k > n$ and $x_k^*(x_n) > \epsilon$ for $n \geq k$. By passing to subsequences and relabelling, we can find a subsequence $(\int_I g_n)$ of $(\int_I f_n)$ and a subsequence (y_k^*) of x_k^* such that

$$\begin{aligned} y_k^* \int_I g_n &= \int_I y_k^* g_n = 0 \quad \text{for } k > n, \\ y_k^* \int_I g_n &= \int_I y_k^* g_n > \epsilon \quad \text{for } n \geq k, \\ \lim_{n \rightarrow \infty} \int_I x^* g_n &= \int_I x^* f \quad \forall x^* \text{ in } X^*. \end{aligned} \quad (3.26)$$

Since the unit ball $B(X^*)$ of X^* is weak* sequentially compact, the sequence (y_k^*) has a subsequence $(y_{k_j}^*)$ which weak* converges to y_0^* , so $\lim_{j \rightarrow \infty} y_{k_j}^* f = y_0^* f$ on I_0 ,

$\lim_{j \rightarrow \infty} \gamma_{k_j}^* F = \gamma_0^* F$ on I_0 , that is, $\lim_{j \rightarrow \infty} \int_I \gamma_{k_j}^* f = \int_I \gamma_0^* f$. To force a contradiction, note that for each k , $\lim_{n \rightarrow \infty} \int_I \gamma_k^* f_n = \int_I \gamma_k^* f$. Hence $\int_I \gamma_k^* f \geq \epsilon$ for each k , and $\int_I \gamma_0^* f \geq \epsilon$. On the other hand, notice that since each g_n is Henstock-Pettis integrable, $(\gamma_{k_j}^*)$ weak* converges to γ_0^* , hence

$$\lim_{j \rightarrow \infty} \int_I \gamma_{k_j}^* g_n = \lim_{j \rightarrow \infty} \gamma_{k_j}^* \int_I g_n = \gamma_0^* \int_I g_n = \int_I \gamma_0^* g_n. \quad (3.27)$$

Since this holds for each n , and since $\lim_{n \rightarrow \infty} \int_I \gamma_0^* g_n = \int_I \gamma_0^* f$, we see that $\int_I \gamma_0^* f = 0$. This contradicts the inequality $\int_I \gamma_0^* f \geq \epsilon$, and proves that C is weakly compact. Since $\lim_{n \rightarrow \infty} \int_I x^* f_n = \int_I x^* f$, the sequence $(\int_I f_n)$ of the Henstock-Pettis integrals is weakly Cauchy. It follows from the weak compactness of C that $\lim_{n \rightarrow \infty} \int_I f_n$ exists weakly in X . Denote $F(I) = \int_I f = \lim_{n \rightarrow \infty} \int_I f_n$ weakly, then $x^* F(I) = x^* \int_I f = \int_I x^* f$ for each x^* in X^* . So f is Henstock-Pettis integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.28)$$

□

COROLLARY 3.14. Suppose that X is a reflexive Banach space and

- (1) $f_n \rightarrow f$ weakly almost everywhere on $[a, b]$ as $n \rightarrow \infty$, where each f_n is Henstock-Pettis integrable on $[a, b]$,
- (2) the primitives F_n of f_n are weakly continuous uniformly in n and weakly ACG^* uniformly in n on $[a, b]$.

Then f is Henstock-Pettis integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.29)$$

THEOREM 3.15. If the following conditions are satisfied:

- (1) $\lim_{n \rightarrow \infty} f_n = f$ weakly almost everywhere on $[a, b]$, where each f_n is Henstock-Dunford integrable on $[a, b]$,
- (2) the primitives F_n of f_n are weakly continuous uniformly in n and weakly ACG^* uniformly in n .

Then f is Henstock-Dunford integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.30)$$

PROOF. Since

- (1) $\lim_{n \rightarrow \infty} x^* f_n = x^* f$ almost everywhere on $[a, b]$,
- (2) the primitives $x^* F_n$ of $x^* f_n$ are continuous uniformly in n and ACG^* uniformly in n .

Then, as in the proof of [Theorem 3.12](#), $x^* f$ is Henstock integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b x^* f_n = \int_a^b x^* f. \quad (3.31)$$

By [Theorem 2.2](#), f is Henstock-Dunford integrable on $[a, b]$ and

$$\lim_{n \rightarrow \infty} \int_a^b f_n = \int_a^b f \text{ weakly.} \quad (3.32)$$

□

REFERENCES

- [1] V. G. Čelidze and A. G. Džvaršeišvili, *The Theory of the Denjoy Integral and some Applications*, Series in Real Analysis, vol. 3, World Scientific Publishing Co., New Jersey, 1989, translated from Russian, with a preface and an appendix by P. S. Bullen. [MR 90k:26013](#). [Zbl 744.26006](#).
- [2] J. Diestel and J. J. Uhl, Jr., *Vector Measures*, Mathematical Surveys, no. 15, American Mathematical Society, Rhode Island, 1977, with a foreword by B. J. Pettis. [MR 56#12216](#). [Zbl 369.46039](#).
- [3] J. L. Gámez and J. Mendoza, *On Denjoy-Dunford and Denjoy-Pettis integrals*, *Studia Math.* **130** (1998), no. 2, 115–133. [MR 99k:28013](#). [Zbl 980.39999](#).
- [4] R. F. Geitz, *Pettis integration*, *Proc. Amer. Math. Soc.* **82** (1981), no. 1, 81–86. [MR 82c:28018](#). [Zbl 506.28007](#).
- [5] R. A. Gordon, *The Denjoy extension of the Bochner, Pettis, and Dunford integrals*, *Studia Math.* **92** (1989), no. 1, 73–91. [MR 90b:28011](#). [Zbl 681.28006](#).
- [6] R. C. James, *Weak compactness and reflexivity*, *Israel J. Math.* **2** (1964), 101–119. [MR 31#585](#). [Zbl 127.32502](#).
- [7] P. Y. Lee, *Lanzhou Lectures on Henstock Integration*, Series in Real Analysis, vol. 2, World Scientific Publishing Co., New Jersey, 1989. [MR 92j:26010](#). [Zbl 699.26004](#).
- [8] G. Q. Liu, *On necessary conditions for Henstock integrability*, *Real Anal. Exchange* **18** (1992/93), no. 2, 522–531. [MR 94e:26014](#). [Zbl 788.26008](#).

YE GUOJU: DEPARTMENT OF MATHEMATICS, NORTHWEST NORMAL UNIVERSITY, LANZHOU 730070, CHINA

E-mail address: yeguofu@21cn.com

AN TIANQING: DEPARTMENT OF MATHEMATICS, NORTHWEST NORMAL UNIVERSITY, LANZHOU 730070, CHINA