ABOUT INTERPOLATION OF SUBSPACES OF REARRANGEMENT INVARIANT SPACES GENERATED BY RADEMACHER SYSTEM

SERGEY V. ASTASHKIN

(Received 1 August 2000 and in revised form 27 November 2000)

ABSTRACT. The Rademacher series in rearrangement invariant function spaces “close” to the space L_∞ are considered. In terms of interpolation theory of operators, a correspondence between such spaces and spaces of coefficients generated by them is stated. It is proved that this correspondence is one-to-one. Some examples and applications are presented.

2000 Mathematics Subject Classification. Primary 46B70; Secondary 46B42, 42A55.

1. Introduction. Let

$$r_k(t) = \text{sign} \sin 2^{k-1} \pi t \quad (k = 1, 2, \dots) \quad (1.1)$$

be the Rademacher functions on the segment $[0, 1]$. Define the linear operator

$$Ta(t) = \sum_{k=1}^{\infty} a_k r_k(t) \quad \text{for } a = (a_k)_{k=1}^{\infty} \in l_2. \quad (1.2)$$

It is well known (cf. [23, pages 340–342]) that Ta is an almost everywhere finite function on $[0, 1]$. Moreover, from Khintchine’s inequality it follows that

$$\|Ta\|_{L_p} \asymp \|a\|_2 \quad \text{for } 1 \leq p < \infty, \quad (1.3)$$

where $\|a\|_p = (\sum_{k=1}^{\infty} |a_k|^p)^{1/p}$. The symbol $\asymp$ means the existence of two-sided estimates with constants depending only on p . Also, it can easily be checked that

$$\|Ta\|_{L_\infty} = \|a\|_1. \quad (1.4)$$

A more detailed information on the behaviour of Rademacher series can be obtained by treating them in the framework of general rearrangement invariant spaces.

Recall that a Banach space X of measurable functions $x = x(t)$ on $[0, 1]$ is said to be a rearrangement invariant space (r.i.s.) if the inequality $x^*(t) \leq y^*(t)$, for $t \in [0, 1]$ and $y \in X$, implies $x \in X$ and $\|x\| \leq \|y\|$. Here and in what follows $z^*(t)$ is the nonincreasing rearrangement of a function $|z(t)|$ with respect to the Lebesgue measure denoted by meas [10, page 83].

Important examples of r.i.s.’s are Marcinkiewicz and Orlicz spaces. Let $\mathcal{P}$ denote the cone of nonnegative increasing concave functions on the semiaxis $(0, \infty)$.

If $\varphi \in \mathcal{P}$, then the Marcinkiewicz space $M(\varphi)$ consists of all measurable functions $x = x(t)$ such that

$$\|x\|_{M(\varphi)} = \sup \left\{ \frac{1}{\varphi(t)} \int_0^t x^*(s) ds : 0 < t \leq 1 \right\} < \infty. \quad (1.5)$$

If $S(t)$ is a nonnegative convex continuous function on $[0, \infty)$, $S(0) = 0$, then the Orlicz space L_S consists of all measurable functions $x = x(t)$ such that

$$\|x\|_S = \inf \left\{ u > 0 : \int_0^1 S\left(\frac{|x(t)|}{u}\right) dt \leq 1 \right\} < \infty. \quad (1.6)$$

In particular, if $S(t) = t^p$ ($1 \leq p < \infty$), then $L_S = L_p$.

For any r.i.s. X on $[0, 1]$ we have $L_\infty \subset X \subset L_1$ [10, page 124]. Let X^0 denote the closure of L_∞ in an r.i.s. X .

In problems discussed below, a special role is played by the Orlicz space L_N , where $N(t) = \exp(t^2) - 1$ or, more precisely, by the space $G = L_N^0$. In [19], V. A. Rodin and E. M. Semenov proved a theorem about the equivalence of Rademacher system to the standard basis in the space l_2 .

THEOREM 1.1. *Suppose that X is an r.i.s. Then*

$$\|Ta\|_X = \left\| \sum_{k=1}^{\infty} a_k r_k \right\|_X \asymp \|a\|_2 \quad (1.7)$$

if and only if $X \supset G$.

By Theorem 1.1, the space G is the minimal space among r.i.s.'s X such that the Rademacher system is equivalent in X to the standard basis of l_2 .

In this paper, we consider problems related to the behaviour of Rademacher series in r.i.s.'s intermediate between L_∞ and G . Here a major role is played by concepts and methods of interpolation theory of operators.

For a Banach couple (X_0, X_1) , $x \in X_0 + X_1$ and $t > 0$, we introduce the Peetre $\mathcal{H}$ -functional

$$\mathcal{H}(t, x; X_0, X_1) = \inf \left\{ \|x_0\|_{X_0} + t\|x_1\|_{X_1} : x = x_0 + x_1, x_0 \in X_0, x_1 \in X_1 \right\}. \quad (1.8)$$

Let Y_0 be a subspace of X_0 and Y_1 a subspace of X_1 . A couple (Y_0, Y_1) is called a $\mathcal{H}$ -subcouple of a couple (X_0, X_1) if

$$\mathcal{H}(t, y; Y_0, Y_1) \asymp \mathcal{H}(t, y; X_0, X_1), \quad (1.9)$$

with constants independent of $y \in Y_0 + Y_1$ and $t > 0$.

In particular, if $Y_i = P(X_i)$, where P is a linear projector bounded from X_i into itself for $i = 0, 1$, then (Y_0, Y_1) is a $\mathcal{H}$ -subcouple of (X_0, X_1) (see [3] or [21, page 136]). At the same time, there are many examples of subcouples that are not $\mathcal{H}$ -subcouples (see [21, page 589], [22], and Remark 3.2 of this paper).

Let $T(l_1)$ (respectively $T(l_2)$) denote the subspace of L_∞ (of G) consisting of all functions of the form $x = Ta$, where T is given by (1.2) and $a \in l_1$ ($\in l_2$). From (1.4) and Theorem 1.1 it follows that

$$\mathcal{H}(t, Ta; T(l_1), T(l_2)) \asymp \mathcal{H}(t, a; l_1, l_2). \quad (1.10)$$

In spite of the fact that $T(l_1)$ is uncomplemented in L_∞ (see [17] or [11, page 134]) the following assertion holds.

THEOREM 1.2. *The couple $(T(l_1), T(l_2))$ is a $\mathcal{H}$ -subcouple of the couple (L_∞, G) . In other words (see (1.10)),*

$$\mathcal{H}(t, Ta; L_\infty, G) \asymp \mathcal{H}(t, a; l_1, l_2), \quad (1.11)$$

with constants independent of $a = (a_k)_{k=1}^\infty \in l_2$ and $t > 0$.

We will use in the proof of Theorem 1.2 an assertion about the distribution of Rademacher sums. It was proved by S. Montgomery-Smith [13].

THEOREM 1.3. *There exists a constant $A \geq 1$ such that for all $a = (a_k)_{k=1}^\infty \in l_2$ and $t > 0$*

$$\begin{aligned} \text{meas} \left\{ s \in [0, 1] : \sum_{k=1}^{\infty} a_k r_k(s) > \varphi_a(t) \right\} &\leq \exp \left(-\frac{t^2}{2} \right), \\ \text{meas} \left\{ s \in [0, 1] : \sum_{k=1}^{\infty} a_k r_k(s) > A^{-1} \varphi_a(t) \right\} &\geq A^{-1} \exp(-At^2), \end{aligned} \quad (1.12)$$

where $\varphi_a(t) = \mathcal{H}(t, a; l_1, l_2)$.

Now we need some definitions from interpolation theory of operators. We say that a linear operator U is bounded from a Banach couple $\vec{X} = (X_0, X_1)$ into a Banach couple $\vec{Y} = (Y_0, Y_1)$ (in short, $U : \vec{X} \rightarrow \vec{Y}$) if U is defined on $X_0 + X_1$ and acts as bounded operator from X_i into Y_i for $i = 0, 1$.

Let $\vec{X} = (X_0, X_1)$ be a Banach couple. A space X such that $X_0 \cap X_1 \subset X \subset X_0 + X_1$ is called an interpolation space between X_0 and X_1 if each linear operator $U : \vec{X} \rightarrow \vec{X}$ is bounded from X into itself.

To every r.i.s. X assign the sequence space F_X of Rademacher coefficients of functions of the form (1.2) from X :

$$\|(a_k)\|_{F_X} = \left\| \sum_{k=1}^{\infty} a_k r_k \right\|_X. \quad (1.13)$$

Well-known properties of Rademacher functions imply that F_X is an r.i. sequence space [19]. Furthermore, Theorem 1.3 and properties of the $\mathcal{H}$ -functional show that F_X is an interpolation space between l_1 and l_2 (see the proof of Theorem 1.2 later). For interpolation r.i.s. between L_∞ and G the correspondence $X \mapsto F_X$ can be defined by using the real interpolation method.

For every $p \in [1, \infty]$, we denote by $l_p(u_k)$, $u_k \geq 0$ ($k = 0, 1, \dots$) the space of all two-sided sequences of real numbers $a = (a_k)_{k=-\infty}^\infty$ such that the norm $\|a\|_{l_p(u_k)} = \|(a_k u_k)\|_p$ is finite. Let E be a Banach lattice of two-sided sequences, $(\min(1, 2^k))_{k=-\infty}^\infty \in E$. If (X_0, X_1) is a Banach couple, then the space of the real $\mathcal{H}$ -method of interpolation $(X_0, X_1)_{\mathcal{H}}^E$ consists of all $x \in X_0 + X_1$ such that

$$\|x\| = \|(\mathcal{H}(2^k, x; X_0, X_1))_k\|_E < \infty. \quad (1.14)$$

It is readily checked that the space $(X_0, X_1)_{\mathcal{H}}^E$ is an interpolation space between X_0 and X_1 (cf. [15, page 422]). In the special case $E = l_p(2^{-k\theta})$ ($0 < \theta < 1$, $1 \leq p \leq \infty$) we obtain the spaces $(X_0, X_1)_{\theta, p}$ (for the detailed exposition of their properties see [4]).

A couple $\vec{X} = (X_0, X_1)$ is said to be a $\mathcal{H}$ -monotone couple if for every $x \in X_0 + X_1$ and $y \in X_0 + X_1$ there exists a linear operator $U : \vec{X} \rightarrow \vec{X}$ such that $y = Ux$ whenever

$$\mathcal{H}(t, y; X_0, X_1) \leq \mathcal{H}(t, x; X_0, X_1) \quad \forall t > 0. \quad (1.15)$$

As it is well known (cf. [15, page 482]), any interpolation space X with respect to a $\mathcal{H}$ -monotone couple (X_0, X_1) is described by the real $\mathcal{H}$ -method. It means that for some E

$$X = (X_0, X_1)_E^{\mathcal{H}}. \quad (1.16)$$

In particular, by the Sparr theorem [20] the couple (l_1, l_2) is a $\mathcal{H}$ -monotone couple. Therefore, if F is an interpolation space between l_1 and l_2 , then there exists E such that

$$F = (l_1, l_2)_E^{\mathcal{H}}. \quad (1.17)$$

Hence Theorem 1.2 allows to find an r.i.s. that contains Rademacher series with coefficients belonging to an arbitrary interpolation space between l_1 and l_2 . In [19], the similar result was obtained for sequence spaces satisfying more restrictive conditions (see Remark 3.3).

THEOREM 1.4. *Let F be an interpolation sequence space between l_1 and l_2 and $F = (l_1, l_2)_E^{\mathcal{H}}$. Then for the r.i.s. $X = (L_\infty, G)_E^{\mathcal{H}}$ we have*

$$\left\| \sum_{k=1}^{\infty} a_k r_k \right\|_X \asymp \|a\|_F \quad (1.18)$$

with constants independent of $a = (a_k)_{k=1}^{\infty}$.

Combining Theorem 1.4 with the above remarks, we get the following assertion. If F is a sequence space, then

$$\|(a_k)\|_F \asymp \left\| \sum_{k=1}^{\infty} a_k r_k \right\|_X \quad \text{for some r.i.s. } X \quad (1.19)$$

if and only if F is an interpolation space between l_1 and l_2 .

The last result shows that the restriction of the correspondence (1.13) to interpolation r.i.s. between L_∞ and G is bijective.

THEOREM 1.5. *Let r.i.s.'s X_0 and X_1 be two interpolation spaces between L_∞ and G . If*

$$\left\| \sum_{k=1}^{\infty} a_k r_k \right\|_{X_0} \asymp \left\| \sum_{k=1}^{\infty} a_k r_k \right\|_{X_1}, \quad (1.20)$$

then $X_0 = X_1$ and the norms of X_0 and X_1 are equivalent.

In [16, 19], the similar results were obtained by additional conditions with respect to spaces X_0 and X_1 .

2. Proofs

PROOF OF THEOREM 1.2. It is known [10, page 164] that the $\mathcal{H}$ -functional of a couple of Marcinkiewicz spaces is given by the formula

$$\mathcal{H}(t, x; M(\varphi_0), M(\varphi_1)) = \sup_{0 < u \leq 1} \frac{\int_0^u x^*(s) ds}{\max(\varphi_0(u), \varphi_1(u)/t)}. \quad (2.1)$$

If $N(t) = \exp(t^2) - 1$, then the Orlicz space L_N coincides with the Marcinkiewicz space $M(\varphi_1)$, where $\varphi_1(u) = u \log_2^{1/2}(2/u)$ [12]. In addition, $L_\infty = M(\varphi_0)$, where $\varphi_0(u) = u$. Therefore,

$$\mathcal{H}(t, x; L_\infty, G) = \sup_{0 < u \leq 1} \left\{ \frac{1}{u} \int_0^u x^*(s) ds \min \left(1, t \log_2^{-1/2} \left(\frac{2}{u} \right) \right) \right\} \quad \text{for } x \in G. \quad (2.2)$$

Since $x^*(u) \leq 1/u \int_0^u x^*(s) ds$, then from (2.2) it follows that

$$\mathcal{H}(t, x; L_\infty, G) \geq \sup_{k=0,1,\dots} \{x^*(2^{-k}) \min(1, t(k+1)^{-1/2})\}. \quad (2.3)$$

Hence,

$$\mathcal{H}(t, x; L_\infty, G) \geq x^*(2^{-k_t}) \quad \text{for } t \geq 1, \quad (2.4)$$

where $k_t = [t^2] - 1$ ($[z]$ is the integral part of a number z).

Now let $a = (a_k)_{k=1}^\infty \in l_2$ and $x(t) = Ta(t) = \sum_{k=1}^\infty a_k r_k(t)$. By the Holmstedt formula [7],

$$\varphi_a(t) \leq \sum_{k=1}^{[t^2]} a_k^* + t \left\{ \sum_{k=[t^2]+1}^\infty (a_k^*)^2 \right\}^{1/2} \leq B \varphi_a(t), \quad (2.5)$$

where $\varphi_a(t) = \mathcal{H}(t, a; l_1, l_2)$, $(a_k^*)_{k=1}^\infty$ is a nonincreasing rearrangement of the sequence $(|a_k|)_{k=1}^\infty$, and $B > 0$ is a constant independent of $a = (a_k)_{k=1}^\infty$ and $t > 0$.

Assume, at first, that $a \notin l_1$. Then inequality (2.5) shows that

$$\lim_{t \rightarrow 0+} \varphi_a(t) = 0, \quad \lim_{t \rightarrow \infty} \varphi_a(t) = \infty. \quad (2.6)$$

The function φ_a belongs to the class $\mathcal{P}$ [4, page 55]. Therefore it maps the semiaxis $(0, \infty)$ onto $(0, \infty)$ one-to-one, and there exists the inverse function φ_a^{-1} . By Theorem 1.3, we have

$$n_{|x|}(\tau) = \text{meas} \{s \in [0, 1] : |x(s)| > \tau\} \geq \psi(\tau) \quad \text{for } \tau > 0, \quad (2.7)$$

where $\psi(\tau) = A^{-1} \exp\{-A[\varphi_a^{-1}(\tau A)]^2\}$. Passing to rearrangements we obtain

$$x^*(s) \geq \psi^{-1}(s) \quad \text{for } 0 < s < A^{-1}. \quad (2.8)$$

Obviously, by condition $t \geq C_1 = C_1(A) = \sqrt{2 \log_2(2A)}$, it holds

$$2^{-k_t/2} < A^{-1} \quad \text{for } k_t = [t^2] - 1. \quad (2.9)$$

Hence (2.4) and (2.8) imply

$$\mathcal{H}(t, x; L_\infty, G) \geq \psi^{-1}(2^{-k_t}). \quad (2.10)$$

Combining the definition of the function ψ with (2.9), we obtain

$$\begin{aligned}\psi^{-1}(2^{-k_t}) &= A^{-1}\varphi_a(A^{-1/2}\ln^{1/2}(A^{-1}2^{k_t})) \geq A^{-1}\varphi_a\left(\sqrt{\frac{k_t\ln 2}{2A}}\right) \\ &\geq A^{-3/2}\sqrt{\frac{\ln 2}{2}}\varphi_a(\sqrt{k_t}) \geq A^{-3/2}\sqrt{\frac{\ln 2}{2}}t^{-1}\sqrt{k_t}\varphi_a(t).\end{aligned}\quad (2.11)$$

From the inequality $t \geq C_1 \geq \sqrt{2}$ it follows that

$$\frac{\sqrt{k_t}}{t} \geq \frac{\sqrt{[t^2]-1}}{\sqrt{[t^2]+1}} \geq 3^{-1/2}.$$
(2.12)

Therefore, by (2.10), we have

$$\mathcal{H}(t, x; L_\infty, G) \geq C_2\varphi_a(t) \quad \text{for } t \geq C_1, \quad (2.13)$$

where $C_2 = C_2(A) = \sqrt{\ln 2/6}A^{-3/2}$.

If now $t \geq 1$, then the concavity of the $\mathcal{H}$ -functional and the previous inequality yield

$$\mathcal{H}(t, x; L_\infty, G) \geq C_1^{-1}\mathcal{H}(tC_1, x; L_\infty, G) \geq \frac{C_2}{C_1}\varphi_a(C_1t) \geq \frac{C_2}{C_1}\varphi_a(t). \quad (2.14)$$

Using the inequalities $\|a\|_2 \leq \|a\|_1$ ($a \in l_1$) and $\|x\|_G \leq \|x\|_\infty$ ($x \in L_\infty$), the definition of the $\mathcal{H}$ -functional, and Theorem 1.1, we obtain

$$\mathcal{H}(t, x; L_\infty, G) = t\|x\|_G \geq C_3t\|a\|_2 = C_3\varphi_a(t) \quad \text{for } 0 < t \leq 1. \quad (2.15)$$

Thus,

$$\mathcal{H}(t, a; l_1, l_2) \leq C\mathcal{H}(t, Ta; L_\infty, G), \quad (2.16)$$

if $C = \max(C_3^{-1}, C_1/C_2)$.

Suppose now $a \in l_1$. By (2.5), without loss of generality, we can assume that the function φ_a maps the semiaxis $(0, \infty)$ injectively onto the interval $(0, \|a\|_1)$. Hence we can define the mappings $\varphi_a^{-1}: (0, \|a\|_1) \rightarrow (0, \infty)$, $\psi: (0, A^{-1}\|a\|_1) \rightarrow (0, A^{-1})$, and $\psi^{-1}: (0, A^{-1}) \rightarrow (0, A^{-1}\|a\|_1)$. Arguing as above, we get inequality (2.16).

The opposite inequality follows from Theorem 1.1 and relation (1.4). Indeed,

$$\begin{aligned}\mathcal{H}(t, Ta; L_\infty, G) &\leq \inf \{ \|Ta^0\|_\infty + t\|Ta^1\|_G : a = a^0 + a^1, a^0 \in l_1, a^1 \in l_2 \} \\ &\leq D\mathcal{H}(t, a; l_1, l_2).\end{aligned}\quad (2.17)$$

□

PROOF OF THEOREM 1.4. It is sufficient to use Theorem 1.2 and the definition of the real $\mathcal{H}$ -method of interpolation. □

For the proof of Theorem 1.5 we need some definitions and auxiliary assertions. These results are also of some independent interest.

Let $f(t)$ be a function defined on the interval $(0, l)$, where $l = 1$ or $l = \infty$. Then the dilation function of f is defined as follows:

$$\mathcal{M}_f(t) = \sup \left\{ \frac{f(st)}{f(s)} : s, st \in (0, l) \right\}, \quad \text{if } t \in (0, l). \quad (2.18)$$

Since this function is semimultiplicative, then there exist numbers

$$\gamma_f = \lim_{t \rightarrow 0^+} \frac{\ln \mathcal{M}_f(t)}{\ln t}, \quad \delta_f = \lim_{t \rightarrow \infty} \frac{\ln \mathcal{M}_f(t)}{\ln t}. \quad (2.19)$$

A Banach couple $\vec{X} = (X_0, X_1)$ is called a partial retract of a couple $\vec{Y} = (Y_0, Y_1)$ if each element $x \in X_0 + X_1$ is orbitally equivalent to some element $y \in Y_0 + Y_1$. The last means that there exist linear operators $U : \vec{X} \rightarrow \vec{Y}$ and $V : \vec{Y} \rightarrow \vec{X}$ such that $Ux = y$ and $Vy = x$.

PROPOSITION 2.1. *Suppose that $M(\varphi)$ is a Marcinkiewicz space on $[0, 1]$. If $\gamma_\varphi > 0$, then $\vec{X} = (L_\infty, M(\varphi))$ is a $\mathcal{H}$ -monotone couple.*

PROOF. It is sufficient to show that the couple $\vec{X}$ is a partial retract of the couple $\vec{Y} = (L_\infty, L_\infty(\tilde{\varphi}))$, where

$$\|x\|_{L_\infty(\tilde{\varphi})} = \sup_{0 < t \leq 1} \tilde{\varphi}(t) |x(t)|, \quad \tilde{\varphi}(t) = \frac{t}{\varphi}(t). \quad (2.20)$$

Indeed, a partial retract of a $\mathcal{H}$ -monotone couple is a $\mathcal{H}$ -monotone couple [15, page 420], and by the Sparr theorem [20] $\vec{Y}$ is a $\mathcal{H}$ -monotone couple.

First note that the inclusion $L_\infty \subset M(\varphi)$ implies $L_\infty + M(\varphi) = M(\varphi)$. So, let $x \in M(\varphi)$. Without loss of generality [10, page 87], assume that $x(t) = x^*(t)$. Define the operator

$$U_1 y(t) = \sum_{k=1}^{\infty} 2^k \int_0^{2^{-k}} y(s) ds \chi_{(2^{-k}, 2^{-k+1}]}(t) \quad \text{for } y \in M(\varphi). \quad (2.21)$$

Clearly, U_1 maps L_∞ into itself. In addition, the concavity of the function φ and properties of the nonincreasing rearrangement imply

$$\|U_1 y\|_{L_\infty(\tilde{\varphi})} \leq 2 \sup_{k=1, 2, \dots} (\varphi(2^{-k+1}))^{-1} \int_0^{2^{-k}} y^*(s) ds \leq 2 \|y\|_{M(\varphi)}. \quad (2.22)$$

Hence $U_1 : \vec{X} \rightarrow \vec{Y}$. Since $x(t)$ is nonincreasing, then $U_1 x(t) \geq x(t)$. Therefore the linear operator

$$Uy(t) = \frac{x(t)}{U_1 x(t)} U_1 y(t) \quad (2.23)$$

is bounded from the couple $\vec{X}$ into the couple $\vec{Y}$. In addition, $Ux(t) = x(t)$.

Take for V the identity mapping, that is, $Vy(t) = y(t)$. Since $\gamma_f > 0$, then, by [10, page 156], we have

$$\|Vy\|_{M(\varphi)} \leq C \sup_{0 < t \leq 1} \tilde{\varphi}(t) y^*(t) \leq C \sup_{0 < t \leq 1} \tilde{\varphi}(t) |y(t)| = C \|y\|_{L_\infty(\tilde{\varphi})}. \quad (2.24)$$

Therefore $V : \vec{Y} \rightarrow \vec{X}$ and $Vx = x$.

Thus an arbitrary element $x \in M(\varphi)$ is orbitally equivalent to itself as to element of the space $L_\infty + L_\infty(\tilde{\varphi})$. This completes the proof. $\square$

COROLLARY 2.2. *If $\gamma_\varphi > 0$, then $(L_\infty, M(\varphi)^0)$ is a $\mathcal{H}$ -monotone couple.*

PROOF. Assume that x and y belong to the space $M(\varphi)^0$ and

$$\mathcal{H}(t, y; L_\infty, M(\varphi)^0) \leq \mathcal{H}(t, x; L_\infty, M(\varphi)^0) \quad \text{for } t > 0. \quad (2.25)$$

If $z \in M(\varphi)^0$, then

$$\mathcal{H}(t, z; L_\infty, M(\varphi)^0) = \mathcal{H}(t, z; L_\infty, M(\varphi)). \quad (2.26)$$

Therefore,

$$\mathcal{H}(t, y; L_\infty, M(\varphi)) \leq \mathcal{H}(t, x; L_\infty, M(\varphi)) \quad \text{for } t > 0. \quad (2.27)$$

Hence, by [Proposition 2.1](#), there exists an operator $T : (L_\infty, M(\varphi)) \rightarrow (L_\infty, M(\varphi))$ such that $y = Tx$. It is readily seen that $M(\varphi)^0$ is an interpolation space of the couple $(L_\infty, M(\varphi))$. Therefore $T : (L_\infty, M(\varphi)^0) \rightarrow (L_\infty, M(\varphi)^0)$. $\square$

We define now two subcones of the cone $\mathcal{P}$. Denote by $\mathcal{P}_0$ the set of all functions $f \in \mathcal{P}$ such that $\lim_{t \rightarrow 0+} f(t) = \lim_{t \rightarrow \infty} f(t)/t = 0$. If $f \in \mathcal{P}$, then $0 \leq \gamma_f \leq \delta_f \leq 1$ [[10](#), page 76]. Let $\mathcal{P}^{+-}$ be the set of all $f \in \mathcal{P}$ such that $0 < \gamma_f \leq \delta_f < 1$. It is obvious that $\mathcal{P}^{+-} \subset \mathcal{P}_0$.

A couple (X_0, X_1) is called a $\mathcal{H}_0$ -complete couple if for any function $f \in \mathcal{P}_0$ there exists an element $x \in X_0 + X_1$ such that

$$\mathcal{H}(t, x; X_0, X_1) \asymp f(t). \quad (2.28)$$

In other words, the set $\mathcal{H}(X_0 + X_1)$ of all $\mathcal{H}$ -functionals of a $\mathcal{H}_0$ -complete couple (X_0, X_1) contains, up to equivalence, the whole of the subcone $\mathcal{P}_0$.

PROPOSITION 2.3. *The Banach couple $(L_1(0, \infty), L_2(0, \infty))$ is a $\mathcal{H}_0$ -complete couple.*

PROOF. By the Holmstedt formula for functional spaces [[7](#)],

$$\mathcal{H}(t, x; L_1, L_2) \asymp \max \left\{ \int_0^{t^2} x^*(s) ds, t \left[\int_{t^2}^\infty (x^*(s))^2 ds \right]^{1/2} \right\}. \quad (2.29)$$

If $f \in \mathcal{P}_0$, then $g(t) = f(t^{1/2})$ belongs to $\mathcal{P}_0$. We denote $x(t) = g'(t)$. Then $x(t) = x^*(t)$ and

$$\int_0^t x(s) ds = g(t). \quad (2.30)$$

Assume that $f \in \mathcal{P}^{+-}$. If $\delta_f < 1$, then there exists $\varepsilon > 0$ such that for some $C > 0$

$$G(s) = f(s^{1/2}) \leq C \left(\sqrt{\frac{s}{t}} \right)^{1-\varepsilon} f(t^{1/2}), \quad \text{if } s \geq t. \quad (2.31)$$

Since $g \in \mathcal{P}_0$, then $g'(t) \leq g(t)/t$. Therefore for $t > 0$

$$\int_t^\infty (x(s))^2 ds \leq \int_t^\infty \frac{g^2(s)}{s^2} ds \leq C^2 t^{\varepsilon-1} (f(t^{1/2}))^2 \int_t^\infty s^{-1-\varepsilon} ds = C^2 \varepsilon t^{-1} (g(t))^2. \quad (2.32)$$

Combining this with (2.29) and (2.30), we obtain

$$\mathcal{H}(t, x; L_1, L_2) \asymp g(t^2) = f(t). \quad (2.33)$$

Thus $\mathcal{H}(L_1 + L_2) \supset \mathcal{P}^{+-}$. Hence, in particular, the intersection $\mathcal{H}(X_0 + X_1) \cap \mathcal{P}^{+-}$ is not empty. Therefore, by [6, Theorem 4.5.7], (L_1, L_2) is a $\mathcal{H}_0$ -complete Banach couple. This completes the proof. $\square$

Let $\mathcal{H}(l_1 + l_2)$ be the set of all $\mathcal{H}$ -functionals corresponding to the couple (l_1, l_2) . By $\mathcal{F}$ we denote the set of all functions $f \in \mathcal{P}$ such that

$$f(t) = f(1)t \quad \text{for } 0 < t \leq 1, \quad \lim_{t \rightarrow \infty} \frac{f(t)}{t} = 0. \quad (2.34)$$

COROLLARY 2.4. *Up to equivalence,*

$$\mathcal{H}(l_1 + l_2) \supset \mathcal{F}. \quad (2.35)$$

PROOF. It is well known (cf. [4, page 142]) that for $x \in L_1(0, \infty) + L_\infty(0, \infty)$ and $u > 0$

$$\mathcal{H}(u, x; L_1, L_\infty) = \int_0^u x^*(s) ds. \quad (2.36)$$

In addition,

$$L_1 = (L_1, L_\infty)_{l_\infty}^{\mathcal{H}}, \quad L_2 = (L_1, L_\infty)_{l_2(2^{-k/2})}^{\mathcal{H}}. \quad (2.37)$$

The spaces l_∞ and $l_2(2^{-k/2})$ are interpolation spaces with respect to the couple $(l_\infty, l_\infty(2^{-k}))$ [4]. Therefore, by the reiteration theorem (see [5] or [14]),

$$\mathcal{H}(t, x; L_1, L_2) \asymp \mathcal{H}(t, \mathcal{H}(\cdot, x; L_1, L_\infty); l_\infty, l_2(2^{-k/2})) \quad \text{for } x \in L_1 + L_2. \quad (2.38)$$

Introduce the average operator:

$$Qx(t) = \sum_{k=1}^{\infty} \int_{k-1}^k x(s) ds \chi_{(k-1, k]}(t), \quad \text{if } t > 0. \quad (2.39)$$

From (2.36) it follows that

$$\mathcal{H}(t, Qx^*; L_1, L_\infty) = \mathcal{H}(t, x; L_1, L_\infty) \quad (2.40)$$

for all positive integers t . Both functions in (2.40) are concave. Therefore,

$$\mathcal{H}(t, Qx^*; L_1, L_\infty) \asymp \mathcal{H}(t, x; L_1 \cdot L_\infty) \quad \forall t \geq 1. \quad (2.41)$$

Hence (2.38) yields

$$\mathcal{H}(t, Qx^*; L_1, L_2) \asymp \mathcal{H}(t, x; L_1, L_2), \quad \text{if } t \geq 1. \quad (2.42)$$

Now let $f \in \mathcal{F}$. Since $\mathcal{F} \subset \mathcal{P}_0$, then, by Proposition 2.3, there exists a function $x \in L_1(0, \infty) + L_2(0, \infty)$ such that

$$\mathcal{H}(t, x; L_1, L_2) \asymp f(t). \quad (2.43)$$

Clearly, the operator Q is a projector in the spaces L_1 and L_2 with norm 1. Moreover, $Q(L_1) = l_1$ and $Q(L_2) = l_2$. Hence, by the theorem about complemented subcouples

mentioned in [Section 1](#) (see [\[3\]](#) or [\[21, page 136\]](#)),

$$\mathcal{H}(t, Qx^*; L_1, L_2) \asymp \mathcal{H}(t, a; l_1, l_2) \quad \text{for } t > 0, \quad (2.44)$$

where $a = (\int_{k-1}^k x^*(s) ds)_{k=1}^\infty$.

Thus [\(2.42\)](#) and [\(2.43\)](#) imply

$$\mathcal{H}(t, a; l_1, l_2) \asymp f(t) \quad \text{for } t \geq 1. \quad (2.45)$$

The last relation also holds if $0 < t \leq 1$. Indeed, in this case

$$\mathcal{H}(t, a; l_1, l_2) = t \|a\|_2 = t \mathcal{H}(1, a; l_1, l_2) \asymp t f(1) = f(t). \quad (2.46)$$

This completes the proof. $\square$

PROOF OF THEOREM 1.5. As it was already mentioned in the proof of [Theorem 1.2](#), the Orlicz space $L_N, N(t) = \exp(t^2) - 1$, coincides with the Marcinkiewicz space $M(\varphi_1)$, for $\varphi_1(u) = u \log_2^{1/2}(2/u)$. Since $\gamma_{\varphi_1} = 1$, then [Corollary 2.2](#) implies that the couple (L_∞, G) is a $\mathcal{H}$ -monotone couple. Hence,

$$X_0 = (l_\infty, G)_{E_0}^{\mathcal{H}}, \quad X_1 = (l_\infty, G)_{E_1}^{\mathcal{H}}, \quad (2.47)$$

for some parameters of the real $\mathcal{H}$ -method of interpolation E_0 and E_1 . By [Theorem 1.4](#),

$$\left\| \sum_{k=1}^{\infty} a_k r_k \right\|_{X_i} \asymp \| (a_k) \|_{F_i}, \quad (2.48)$$

where $F_i = (l_1, l_2)_{E_i}^{\mathcal{H}} (i = 0, 1)$. So

$$(l_1, l_2)_{E_0}^{\mathcal{H}} = (l_1, l_2)_{E_1}^{\mathcal{H}}. \quad (2.49)$$

Equation [\(2.49\)](#) means that the norms of spaces E_0 and E_1 are equivalent on the set $\mathcal{H}(l_1 + l_2)$. It is readily to check that this set coincides, up to the equivalence, with the set $\mathcal{H}(L_\infty + G)$ of all $\mathcal{H}$ -functionals corresponding to the couple (L_∞, G) . More precisely,

$$\mathcal{H}(l_1 + l_2) = \mathcal{H}(L_\infty + G) = \mathcal{F}. \quad (2.50)$$

In fact, by [Theorem 1.2](#) and [Corollary 2.2](#), $\mathcal{F} \subset \mathcal{H}(l_1 + l_2) \subset \mathcal{H}(L_\infty + G)$. On the other hand, since $L_\infty \subset G$ with the constant 1 and L_∞ is dense in G , then $\mathcal{H}(L_\infty + G) \subset \mathcal{F}$ [\[15, page 386\]](#).

Now let $x \in X_0$. By [\(2.47\)](#), we have $(\mathcal{H}(2^k, x; L_\infty, G))_k \in X_0$. Using [\(2.50\)](#), we can find $a \in l_2$ such that

$$\mathcal{H}(2^k, a; l_1, l_2) \asymp \mathcal{H}(2^k, x; L_\infty, G) \quad (2.51)$$

for all positive integers k . Since a parameter of $\mathcal{H}$ -method is a Banach lattice, then this implies $(\mathcal{H}(2^k, a; l_1, l_2))_k \in E_0$. Therefore, by [\(2.49\)](#), $(\mathcal{H}(2^k, a; l_1, l_2))_k \in E_1$, that is, $(\mathcal{H}(2^k, x; L_\infty, G))_k \in E_1$ or $x \in X_1$. Thus $X_0 \subset X_1$. Arguing as above, we obtain the converse inclusion, and $X_0 = X_1$ as sets. Since X_0 and X_1 are Banach lattices, then their norms are equivalent. This completes the proof. $\square$

3. Final remarks and examples

REMARK 3.1. Combining Theorems 1.2, 1.4, and 1.5 with results obtained in [8], we can prove similar assertions for lacunary trigonometric series. Moreover, taking into account the main result of [1], we can extend Theorems 1.2, 1.4, and 1.5 to Sidon systems of characters of a compact abelian group.

REMARK 3.2. In Theorem 1.2, we cannot replace the space G by L_q with some $q < \infty$. Indeed, suppose that the couple $(T(l_1), T(l_2))$ is a $\mathcal{H}$ -subcouple of the couple (L_∞, L_q) , that is,

$$\mathcal{H}(t, a; l_1, l_2) \asymp \mathcal{H}(t, Ta; L_\infty, L_q). \quad (3.1)$$

Let $E = l_p(2^{-\theta k})$, where $0 < \theta < 1$ and $p = q/\theta$. Applying the $\mathcal{H}$ -method of interpolation $(\cdot, \cdot)_E^{\mathcal{H}}$ to the couples (l_1, l_2) and (L_∞, L_q) , we obtain

$$\|Ta\|_p \asymp \|a\|_{r,p} = \left\{ \sum_{k=1}^{\infty} (a_k^*)^p k^{p/r-1} \right\}^{1/p}. \quad (3.2)$$

Since $r = 2/(2-\theta) < 2$ [4, page 142], then this contradicts with (1.3).

REMARK 3.3. Clearly, a partial retract of a couple $\vec{Y} = (Y_0, Y_1)$ is a $\mathcal{H}$ -subcouple of $\vec{Y}$. The opposite assertion is not true, in general (nevertheless, some interesting examples of $\mathcal{H}$ -subcouples and partial retracts simultaneously are given in [9]). Indeed, by Theorem 1.2, the subcouple (l_1, l_2) is a $\mathcal{H}$ -subcouple of the couple (L_∞, G) . Assume that (l_1, l_2) is a partial retract of this couple. Then (see the proof of Proposition 2.1) (l_1, l_2) is a partial retract of the couple $(L_\infty, L_\infty(\log_2^{-1/2}(2/t)))$, as well. Therefore, by Lemma 1 from [2] and [4, page 142] it follows that

$$[l_1, l_2]_\theta = (l_1, l_2)_{\theta, \infty} = l_{p, \infty}, \quad (3.3)$$

where $[l_1, l_2]_\theta$ is the space of the complex method of interpolation [4], $0 < \theta < 1$, and $p = 2/(2-\theta)$. On the other hand, it is well known [4, page 139] that

$$[l_1, l_2]_\theta = l_p \quad \text{for } p = \frac{2}{2-\theta}. \quad (3.4)$$

This contradiction shows that the couple (l_1, l_2) is not a partial retract of the couple (L_∞, G) .

Using Theorem 1.4, we can find coordinate sequence spaces of coefficients of Rademacher series belonging to certain r.i.s.'s.

EXAMPLE 3.4. Let X be the Marcinkiewicz space $M(\varphi)$, where $\varphi(t) = t \log_2 \log_2(16/t)$, $0 < t \leq 1$. Show that

$$\left\| \sum_{k=1}^{\infty} a_k r_k \right\|_{M(\varphi)} \asymp \|a\|_{l_1(\log)}, \quad (3.5)$$

where $l_1(\log)$ is the space of all sequences $a = (a_k)_{k=1}^{\infty}$ such that the norm

$$\|a\|_{l_1(\log)} = \sup_{k=1,2,\dots} \log_2^{-1}(2k) \sum_{i=1}^k a_i^* \quad (3.6)$$

is finite. Taking into account [Theorem 1.4](#), it is sufficient to check that

$$(l_1, l_2)_F^{\mathcal{H}} = l_1(\log), \quad (3.7)$$

$$(l_\infty, G)_F^{\mathcal{H}} = M(\varphi), \quad (3.8)$$

for some parameter F of the $\mathcal{H}$ -method of interpolation. More precisely, we will prove that (3.7) and (3.8) are true for $F = l_\infty(u_k)$, where $u_k = 1/(k+1)$ ($k \geq 0$) and $u_k = 1$ ($k < 0$).

By the Holmstedt formula (2.5),

$$\varphi_a(2^k) \leq \sum_{i=1}^{2^{2k}} a_i^* + 2^k \left[\sum_{i=2^{2k+1}}^{\infty} (a_i^*)^2 \right]^{1/2} \leq B\varphi_a(2^k) \quad \text{for } k = 0, 1, 2, \dots, \quad (3.9)$$

where, as before, $\varphi_a(t) = \mathcal{H}(t, a; l_1, l_2)$. Without loss of generality, assume that $a_i = a_i^*$. If $\|a\|_{l_1(\log)} = R < \infty$, then by (3.6),

$$\sum_{i=1}^{2^{2k}} a_i^* \leq 2R(k+1). \quad (3.10)$$

In particular, this implies $a_{2^{2k}} \leq 2^{-2k+1}R(k+1)$, for nonnegative integer k . Using (3.10), we obtain

$$\begin{aligned} \sum_{i=2^{2k+1}}^{\infty} a_i^2 &= \sum_{j=k}^{\infty} \sum_{i=2^{2j+1}}^{2^{2(j+1)}} a_i^2 \leq 3 \sum_{j=k}^{\infty} 2^{2j} a_{2^{2j}}^2 \leq 12R^2 \sum_{j=k}^{\infty} 2^{-2j} (j+1)^2 \\ &\leq 192R^2 \int_{k+1}^{\infty} x^2 2^{-2x} dx \leq 144R^2 (k+1)^2 2^{-2k}. \end{aligned} \quad (3.11)$$

Hence the second term in (3.9) does not exceed $12R(k+1)$. Therefore, if $E = (l_1, l_2)_F^{\mathcal{H}}$, then (3.10) implies

$$\|a\|_E = \sup_{k=0,1,\dots} \frac{\varphi_a(2^k)}{k+1} \leq 14\|a\|_{l_1(\log)}. \quad (3.12)$$

Conversely, if $2^{2j} + 1 \leq k \leq 2^{2(j+1)}$ for some $j = 0, 1, 2, \dots$, then from (3.9) it follows that

$$\sum_{i=1}^k a_i \leq B\varphi_a(2^{j+1}) \leq \sum_{i=1}^{2^{2(j+1)}} a_i \leq B\|a\|_E(j+2) \leq 2B\log_2(2k)\|a\|_E. \quad (3.13)$$

Therefore, $\|a\|_{l_1(\log)} \leq 2B\|a\|_E$ and (3.7) is proved.

We pass now to function spaces. At first, we introduce one more interpolation method which is, actually, a special case of the real method of interpolation. For a function $\varphi \in \mathcal{P}$ and an arbitrary Banach couple (X_0, X_1) define generalized Marcinkiewicz space as follows:

$$M_\varphi(X_0, X_1) = \left\{ x \in X_0 + X_1 : \sup_{t>0} \frac{\mathcal{H}(t, x; X_0, X_1)}{\varphi(t)} < \infty \right\}. \quad (3.14)$$

Let $\varphi_0(t) = \min(1, t)$, $\varphi_1(t) = \min(1, t \log_2^{1/2}[\max(2, 2/t)])$, and $N(t) = \exp(t^2) - 1$, as before. By equation (2.36), we have

$$L_\infty = M_{\varphi_0}(L_1, L_\infty), \quad L_N = M_{\varphi_1}(L_1, L_\infty), \quad (3.15)$$

(here L_∞ and L_N are functional spaces on the segment $[0, 1]$). In addition, using similar notation, it is easy to check that

$$(X_0, X_1)_F^{\mathcal{K}} = M_\rho(X_0, X_1), \quad (3.16)$$

for an arbitrary Banach couple (X_0, X_1) and $\rho(t) = \log_2(4+t)$. Hence, by the reiteration theorem for generalized Marcinkiewicz spaces [15, page 428], we obtain

$$(L_\infty, L_N)_F^{\mathcal{K}} = M_\rho(M_{\varphi_0}(L_1, L_\infty), M_{\varphi_1}(L_1, L_\infty)) = M_{\varphi_\rho}(L_1, L_\infty) = M(\varphi_\rho), \quad (3.17)$$

where $\varphi_\rho(t) = \varphi_0(t)\rho(\varphi_1(t)/\varphi_0(t))$. A simple calculation gives $\varphi_\rho(t) \asymp \varphi(t)$, if $t > 0$. Thus,

$$(L_\infty, L_N)_F^{\mathcal{K}} = M(\varphi). \quad (3.18)$$

It is readily seen that $\mathcal{H}(t, x; L_\infty, G) = \mathcal{H}(t, x; L_\infty, L_N)$, for all $x \in G$. Therefore, for such x the norm $\|x\|_{M(\varphi)}$ is equal to the norm $\|x\|_Y$, where $Y = (L_\infty, G)_F^{\mathcal{K}}$. On the other hand, for $x \in M(\varphi)$

$$\frac{1}{t \log_2^{1/2}(2/t)} \int_0^t x^*(s) ds \leq \|x\|_{M(\varphi)} \frac{\log_2 \log_2(16/t)}{\log_2^{1/2}(2/t)} \rightarrow 0 \quad \text{as } t \rightarrow 0+. \quad (3.19)$$

This implies that $M(\varphi) \subset G$ [10, page 156]. Thus $Y = M(\varphi)$, and (3.8) is proved. Equivalence (3.5) follows now, as already stated, from (3.7) and (3.8).

REMARK 3.5. Theorems 1.4 and 1.5 strengthen results of [18, 19], where similar assertions are obtained for sequence spaces F satisfying more restrictive conditions. For instance, we can readily show that the norm of the dilation operator

$$\sigma_n a = \left(\underbrace{a_1, \cdot, a_1}_n, \underbrace{a_2, \cdot, a_2}_n, \dots \right) \quad (3.20)$$

in the space $l_1(\ln)$ (see Example 3.6) is equal to n . Therefore, condition (11) from [19] fails for this space and the theorems obtained in [18, 19] cannot be applied to it. Similarly, the Marcinkiewicz space $M(\varphi)$ from Example 3.4 does not satisfy the conditions of Theorem 8 of [19].

Using Theorems 1.4 and 1.5, we can derive certain interpolation relations.

EXAMPLE 3.6. Let $\varphi \in \mathcal{P}$ and $1 \leq p < \infty$. Recall that the Lorentz space $\Lambda_p(\varphi)$ consists of all measurable functions $x = x(s)$ such that

$$\|x\|_{\varphi, p} = \left\{ \int_0^1 (x^*(s))^p d\varphi(s) \right\}^{1/p} < \infty. \quad (3.21)$$

In [19], V. A. Rodin and E. M. Semenov proved that

$$\left\| \sum_{k=1}^{\infty} a_k r_k \right\|_{\varphi, p} \asymp \| (a_k) \|_p, \quad (3.22)$$

where $\varphi(s) = \log_2^{1-p}(2/s)$ and $1 < p < 2$. Moreover, the space $\Lambda_p(\varphi)$ is the unique r.i.s. having this property. Note that $l_p = (l_1, l_2)_{\theta, p}$, where $\theta = 2(p-1)/p$ [4, page 142]. Therefore, by Theorem 1.4, we obtain

$$(L_{\infty}, G)_{\theta, p} = \Lambda_p(\varphi) \quad (3.23)$$

for the same p and θ .

ACKNOWLEDGEMENT. The author is grateful to Prof. S. Montgomery-Smith for useful advices and to referees for their suggestions and remarks.

REFERENCES

- [1] N. Asmar and S. Montgomery-Smith, *On the distribution of Sidon series*, Ark. Mat. **31** (1993), no. 1, 13–26. [MR 94i:43006](#). [Zbl 836.43011](#).
- [2] S. V. Astashkin, *$\mathcal{H}$ -monotone weighted couples generated by space noninvariant concerning to shift*, Internat. Sem. Differential Equations (Samara), vol. 2, 1998, pp. 19–25.
- [3] M. S. Baouendi and C. Goulaouic, *Commutation de l'intersection et des foncteurs d'interpolation*, C. R. Acad. Sci. Paris Sér. A-B **265** (1967), 313–315 (French). [MR 37#727](#). [Zbl 166.10702](#).
- [4] J. Bergh and J. Löfström, *Interpolyatsionnye Prostranstva [Interpolation Spaces]*, Mir, Moscow, 1980 (Russian), Vvedenie (An introduction). Translated from English by V. S. Krjučkov and P. I. Lizorkin. [MR 82c:46083](#).
- [5] J. A. Brudnyi and N. J. Krugljak, *Functors of real interpolation*, Dokl. Akad. Nauk SSSR **256** (1981), no. 1, 14–17 (Russian). [MR 82g:46115](#).
- [6] ———, *Interpolation Functors and Interpolation Spaces. Vol. I*, North-Holland Mathematical Library, vol. 47, North-Holland Publishing Co., Amsterdam, 1991, translated from Russian by Natalie Wadhwa with a preface by Jaak Peetre. [MR 93b:46141](#). [Zbl 743.46082](#).
- [7] T. Holmstedt, *Interpolation of quasi-normed spaces*, Math. Scand. **26** (1970), 177–199. [MR 54#3440](#). [Zbl 193.08801](#).
- [8] J. Jakubowski and S. Kwapien, *On multiplicative systems of functions*, Bull. Acad. Polon. Sci. Sér. Sci. Math. **27** (1979), no. 9, 689–694. [MR 82c:60014](#). [Zbl 493.42036](#).
- [9] S. V. Kislyakov and K. Shu, *Real interpolation and singular integrals*, Algebra i Analiz **8** (1996), no. 4, 75–109, translation in St. Petersburg Math. J. **8** (1997), no. 4, 593–615. [MR 98c:46161](#).
- [10] S. G. Kreĭn, J. I. Petunin, and E. M. Semënov, *Interpolyatsiya Lineĭnykh Operatorov [Interpolation of Linear Operators]*, Nauka, Moscow, 1978 (Russian). [MR 81f:46086](#). [Zbl 499.46044](#).
- [11] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces. II. Function spaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas], vol. 97, Springer-Verlag, Berlin, 1979. [MR 81c:46001](#). [Zbl 403.46022](#).
- [12] G. G. Lorentz, *Relations between function spaces*, Proc. Amer. Math. Soc. **12** (1961), 127–132. [MR 23#A511](#). [Zbl 124.31704](#).
- [13] S. J. Montgomery-Smith, *The distribution of Rademacher sums*, Proc. Amer. Math. Soc. **109** (1990), no. 2, 517–522. [MR 91a:60034](#). [Zbl 696.60013](#).
- [14] P. Nilsson, *Reiteration theorems for real interpolation and approximation spaces*, Ann. Mat. Pura Appl. (4) **132** (1982), 291–330. [MR 86c:46089](#). [Zbl 514.46049](#).

- [15] V. I. Ovchinnikov, *The method of orbits in interpolation theory*, Math. Rep. **1** (1984), no. 2, i-x and 349–515. [MR 88d:46136](#). [Zbl 875.46007](#).
- [16] V. A. Rodin, *Fourier series in symmetric space and interpolation of operators*, Ph.D. thesis, Cand. Phys.-Math. Sci., Voronezh, 1973.
- [17] V. A. Rodin and E. M. Semënov, *The complementability of a subspace that is generated by the Rademacher system in a symmetric space*, Funktsional. Anal. i Prilozhen. **13** (1979), no. 2, 91–92 (Russian). [MR 80j:46048](#).
- [18] V. A. Rodin and E. M. Semenov (eds.), *Distribution functions of Rademacher series*, Vseross School on Stoch. Methods, vol. 3, Tuapse, 1996.
- [19] V. A. Rodin and E. M. Semyonov, *Rademacher series in symmetric spaces*, Anal. Math. **1** (1975), no. 3, 207–222. [MR 52#8905](#). [Zbl 315.46031](#).
- [20] G. Sparr, *Interpolation of weighted L_p -spaces*, Studia Math. **62** (1978), no. 3, 229–271. [MR 80d:46055](#). [Zbl 393.46029](#).
- [21] H. Triebel, *Teoriya Interpolyatsii, Funktsional'nye Prostranstva, Differentsial'nye Operatory* [Interpolation Theory. Function Spaces. Differential Operators], Mir, Moscow, 1980 (Russian), translated from English. [Zbl 531.46025](#).
- [22] R. Wallstén, *Remarks on interpolation of subspaces*, Function Spaces and Applications (Lund, 1986), Lecture Notes in Math., vol. 1302, Springer-Verlag, Berlin, 1988, pp. 410–419. [MR 89h:46042](#). [Zbl 662.46079](#).
- [23] A. Zigmund, *Trigonometricheskie Ryady. Tomy I* [Trigonometric Series], Izdat. “Mir”, Moscow, 1965 (Russian), translated from English by O. S. Ivasev Musatov. [MR 31#2554](#). [Zbl 121.05804](#).

SERGEY V. ASTASHKIN: DEPARTMENT OF MATHEMATICS, SAMARA STREET UNIVERSITY, ACADEMIC PAVLOV STREET, 1, SAMARA, 443011, RUSSIA
E-mail address: astashkn@ssu.samara.ru