GENERALIZED PERIODIC AND GENERALIZED BOOLEAN RINGS

HOWARD E. BELL and ADIL YAQUB

(Received 21 August 2000)

ABSTRACT. We prove that a generalized periodic, as well as a generalized Boolean, ring is either commutative or periodic. We also prove that a generalized Boolean ring with central idempotents must be nil or commutative. We further consider conditions which imply the commutativity of a generalized periodic, or a generalized Boolean, ring.

2000 Mathematics Subject Classification. 16D70, 16U80.

Throughout, R denotes a ring, N the set of nilpotents, C the center, and E the set of idempotents of R . A ring R is called *periodic* if for every x in R , there exist distinct positive integers m, n such that $x^m = x^n$. We now formally state the definitions of a generalized periodic ring and a generalized Boolean ring.

DEFINITION 1. A ring R is called *generalized periodic* if for every x in R such that $x \notin (N \cup C)$, we have $x^n - x^m \in (N \cap C)$, for some positive integers m, n of opposite parity.

DEFINITION 2. A ring R is called *generalized Boolean* if for every x in R such that $x \notin (N \cup C)$, there exists an even positive integer n such that $x - x^n \in (N \cap C)$.

THEOREM 3. *If R is a generalized periodic ring, then R is either commutative or periodic.*

PROOF. Let N and C denote the set of nilpotents and the center of R , respectively. We distinguish three cases.

CASE 1 ($N \subseteq C$). Then $x \notin C$ implies $x \notin (N \cup C)$, and hence there exist distinct positive integers m, n such that $x^m - x^n \in N$, with $n > m$. Suppose $(x^m - x^n)^k = 0$. Then, as is readily verified,

$$(x - x^{n-m+1})^k x^{k(m-1)} = 0, \quad (1)$$

which, in turn, implies that

$$\begin{aligned} (x - x^{n-m+1})^{km} &= (x - x^{n-m+1})^k x^{k(m-1)} g(x) \\ &= 0, \end{aligned} \quad (2)$$

where

$$g(\lambda) \in \mathbb{Z}[\lambda]. \quad (3)$$

We have thus shown that

$$x - x^{n-m+1} \in N, \quad \forall x \notin C, \quad (n - m + 1 > 1). \quad (4)$$

Recall that, in our present case, we assumed that $N \subseteq C$, and hence by (4),

$$x - x^{n-m+1} \in C, \quad \forall x \notin C, \quad (n - m + 1 > 1). \quad (5)$$

Since (5) is trivially satisfied if $x \in C$, we see that

$$x - x^{n(x)} \in C, \quad \text{for some } n(x) > 1, \text{ where } x \in R \text{ (arbitrary)}. \quad (6)$$

Therefore, R is commutative, by a well-known theorem of Herstein [3].

CASE 2 ($C \subseteq N$). Then $x \notin N$ implies $x \notin (N \cup C)$, and hence there exist distinct positive integers m, n such that $x^n - x^m \in N$, with $n > m$. Repeating the argument used to prove (4), we see that

$$x - x^{n-m+1} \in N, \quad \forall x \notin N, \quad (n - m + 1 > 1). \quad (7)$$

Since (7) is trivially satisfied for all $x \in N$, we conclude that

$$x - x^{k(x)} \in N, \quad \text{for some } k(x) > 1, \text{ where } x \in R \text{ (arbitrary)}. \quad (8)$$

By a well-known theorem of Chacron [2], equation (8) implies that R is periodic.

CASE 3 ($C \not\subseteq N$ and $N \not\subseteq C$). In this case, let

$$z \in C \setminus N, \quad u \in N \setminus C. \quad (9)$$

Equation (9) readily implies that $z + u \notin C$ and $z + u \notin N$, and hence (see Definition 1)

$$(z + u)^n - (z + u)^m \in N, \quad \text{for some integers } n > m \geq 1. \quad (10)$$

Since z commutes with the nilpotent element u , (10) implies that

$$z^n - z^m + u' \in N, \quad \text{where } u' \in N, \quad u' \text{ commutes with } z. \quad (11)$$

Hence $z^n - z^m \in N$, with $n > m \geq 1$. Now, a repetition of the argument used in the proof of (4) shows that

$$z - z^{n-m+1} \in N, \quad \forall z \in C \setminus N, \quad (n - m + 1 > 1). \quad (12)$$

Trivially,

$$x - x^k \in N, \quad \forall x \in N, \quad \forall k \in \mathbb{Z}^+. \quad (13)$$

Finally, if $x \notin (N \cup C)$, then

$$x^n - x^m \in N, \quad \text{for some integers } n > m \geq 1. \quad (14)$$

Again, repeating the argument used in the proof of (4), we see that

$$x - x^{n-m+1} \in N, \quad \forall x \notin (N \cup C), \quad (n - m + 1 > 1). \quad (15)$$

Combining (12), (13), and (15), we conclude that

$$x - x^{k(x)} \in N, \quad \text{for some } k(x) > 1, \text{ where } x \in R \text{ (arbitrary)}. \quad (16)$$

Thus, by Chacron's theorem [2], R is periodic. This completes the proof. $\square$

COROLLARY 4. *If R is a generalized Boolean ring, then R is either commutative or periodic.*

This follows at once, since a generalized Boolean ring is necessarily a generalized periodic ring (see Definitions 1 and 2).

Before proving the next theorem, we prove the following lemma.

LEMMA 5. *Let R be a generalized periodic ring. If e is any nonzero central idempotent in R and $a \in N$, then $ea \in C$.*

PROOF. The proof is by contradiction. Suppose the lemma is false, and let

$$\eta_0 \in N, \quad e\eta_0 \notin C. \quad (17)$$

Since $e \in C$ and $\eta_0 \in N$, therefore $e\eta_0$ is nilpotent. Let

$$(e\eta_0)^\alpha \in C, \quad \forall \alpha \geq \alpha_0, \quad \alpha_0 \text{ minimal}. \quad (18)$$

Since $e\eta_0 \notin C$ (see (17)), therefore $\alpha_0 > 1$. Let $\eta = (e\eta_0)^{\alpha_0-1}$. Then,

$$\begin{aligned} \eta &= (e\eta_0)^{\alpha_0-1} \in N, \quad \eta \notin C \text{ (by the minimality of } \alpha_0), \\ \eta^k &\in C, \quad \forall k \geq 2, \quad e \in C, \quad e^2 = e \neq 0, \quad e \notin N. \end{aligned} \quad (19)$$

Equation (19) implies that $e + \eta \notin C$ and $e + \eta \notin N$, and hence (see Definition 1)

$$(e + \eta)^{m'} - (e + \eta)^{n'} \in C, \quad (20)$$

where m', n' are of *opposite* parity. Combining (20) and (19), we see that (keep in mind that $e\eta = \eta$; see (19))

$$(m' - n')e\eta \in C, \quad (21)$$

where $m' - n'$ is an odd integer. Equation (19) also implies that $(-e + \eta)$ is not in $(N \cup C)$, so

$$(-e + \eta)^{m''} - (-e + \eta)^{n''} \in N, \quad (22)$$

where m'', n'' are of *opposite* parity. Combining (19) and (22), we see that

$$(-e)^{m''} - (-e)^{n''} \in N, \quad (23)$$

and hence $2e \in N$, since m'' and n'' are of *opposite* parity. Therefore, $(2e)^y = 0$, $y \in \mathbb{Z}^+$, and thus $2^y e = 0$, which implies that

$$2^y e\eta \in C; \quad y \in \mathbb{Z}^+. \quad (24)$$

Now, combining (21) and (24), keeping in mind that $(2^y, m' - n') = 1$, we see that $e\eta \in C$, and hence, by (19), $\eta = e\eta \in C$, which contradicts (19). This contradiction proves the lemma. $\square$

As usual, $[x, y] = xy - yx$ denotes the commutator of x and y .

We are now in a position to prove the following theorem.

THEOREM 6. *Suppose R is a generalized periodic ring, and suppose that there exists an element c in C , with $c \neq 0$, such that*

$$c[x, y] = 0 \text{ implies } [x, y] = 0, \forall x, y \in R. \quad (25)$$

Then R is commutative.

PROOF. We distinguish two cases.

CASE 1 ($c \in N$). In this case, $c^k = 0$ for some positive integer k , and hence

$$c^k[x, y] = 0, \quad \forall x, y \in R. \quad (26)$$

Combining (25) and (26), we see that

$$\begin{aligned} c^k[x, y] = 0 &\Rightarrow c[c^{k-1}x, y] = 0 \Rightarrow [c^{k-1}x, y] = 0 \Rightarrow c^{k-1}[x, y] = 0 \\ &\Rightarrow \dots \Rightarrow c[x, y] = 0 \Rightarrow [x, y] = 0. \end{aligned} \quad (27)$$

Thus, $c^k[x, y] = 0$ implies $[x, y] = 0$, and hence R is commutative.

CASE 2 ($c \notin N$). In view of [Theorem 3](#), we may assume that R is periodic. This implies, in particular, that c^m is idempotent for some positive integer m . Furthermore, $c^m \neq 0$ (since $c \notin N$ in our present case). The net result is (since $c \in C$ also)

$$c^m = e \text{ is a nonzero central idempotent in } R. \quad (28)$$

Let $a \in N$. By [Lemma 5](#) and equation (28), we have $ea \in C$, and hence $[ea, x] = 0$ for all $x \in R$, which implies

$$[c^m a, x] = c^m [a, x] = 0, \quad \forall x \in R. \quad (29)$$

The argument used in [Case 1](#) of [Theorem 6](#) shows that

$$c^m[a, x] = 0 \text{ implies } [a, x] = 0, \quad (30)$$

and hence (see (29))

$$[a, x] = 0 \quad \forall x \in R, \forall a \in N. \quad (31)$$

Thus, R is a periodic ring with the property that $N \subseteq C$. By a well-known theorem of Herstein [\[4\]](#), it follows that R is commutative, and the theorem is proved. $\square$

COROLLARY 7. *Suppose that R is a generalized periodic ring with identity 1. Then, R is commutative.*

[Corollary 7](#) follows at once by taking $c = 1$ in [Theorem 6](#).

Since a generalized Boolean ring is also a generalized periodic ring, therefore we have the following corollary.

COROLLARY 8. *A generalized Boolean ring with identity 1 is necessarily commutative.*

Another corollary is the following result, proved by the authors in [\[1\]](#).

COROLLARY 9. *Suppose that R is a generalized periodic ring containing a central element which is not a zero divisor. Then R is commutative.*

This follows at once, since the hypotheses of this corollary imply the hypotheses of [Theorem 6](#).

THEOREM 10. *Suppose R is a generalized periodic ring. Suppose, further, that there exists a nonzero central element c such that*

$$ca = 0 \text{ implies } a = 0, \forall a \in N. \quad (32)$$

Then R is commutative.

PROOF. In [\[1\]](#), the authors proved the following result:

$$\begin{aligned} &\text{If } R \text{ is a generalized periodic ring, then the nilpotents} \\ &N \text{ form an ideal and } R/N \text{ is commutative.} \end{aligned} \quad (33)$$

Let $x, y \in R$. By [\(33\)](#), for all $\bar{x}, \bar{y}$ in R/N , $\bar{x}\bar{y} = \bar{y}\bar{x}$, and hence $[x, y] \in N$. Taking $a = [x, y] \in N$ in [\(32\)](#), we see that [\(32\)](#) yields

$$c[x, y] = 0 \text{ implies } [x, y] = 0, \forall x, y \in R. \quad (34)$$

The theorem now follows at once from [Theorem 6](#). $\square$

THEOREM 11. *A generalized Boolean ring R with central idempotents is necessarily nil ($R = N$) or commutative ($R = C$).*

PROOF. Since R is also a generalized periodic ring, therefore by [Theorem 3](#), R is commutative or periodic. If R is commutative, there is nothing to prove. So we may assume that R is periodic. We now distinguish two cases.

CASE 1 ($C \subseteq N$). Recall that, by hypothesis, the set E of idempotents is central, and hence $E \subseteq C \subseteq N$ (in the present case). Thus, $E \subseteq N$, and hence $E = \{0\}$. Therefore,

$$\text{zero is the only idempotent of } R. \quad (35)$$

Let $x \in R$. Since R is periodic, therefore x^k is idempotent for some positive integer k , and hence by [\(35\)](#), $x^k = 0$, which proves that R is nil.

CASE 2 ($C \not\subseteq N$). Then, for some $c \in R$, we have

$$c \in C, \quad c \notin N. \quad (36)$$

Again, since R is periodic, c^m is idempotent for some positive integer m . Moreover, $c^m \neq 0$ (since $c \notin N$). The net result is (see [\(36\)](#))

$$e = c^m \text{ is a nonzero central idempotent of } R. \quad (37)$$

Now, suppose $a \in N$. Since $0 \neq e \in C$ and $a \in N$, therefore $e + a \notin N$. Suppose, for the moment, that $a \notin C$. Then $e + a \notin C$ (since $e \in C$), and hence $e + a \notin (N \cup C)$. Therefore, by [Definition 2](#),

$$(e + a) - (e + a)^n \in (N \cap C), \quad \text{for some even integer } n \geq 2. \quad (38)$$

Since R is also a generalized periodic ring, therefore by [Lemma 5](#) (see (37))

$$ea^i \in C, \quad \forall i \in \{1, \dots, n-1\}, \quad (0 \neq e = e^2, e \in C, a \in N). \quad (39)$$

Combining (38) and (39), we see that

$$a - a^n \in C, \quad \forall a \in N \setminus C. \quad (40)$$

Since (40) is trivially satisfied for $a \in (N \cap C)$, therefore

$$a - a^n \in C, \quad \forall a \in N, \quad n \geq 2. \quad (41)$$

We claim that

$$N \subseteq C. \quad (42)$$

The proof is by contradiction. Suppose (42) is false. Then, for some $a \in R$, we have

$$a \in N, \quad a \notin C. \quad (43)$$

Since $a \in N$, there exists a positive integer σ_0 such that

$$a^\sigma \in C, \quad \forall \sigma \geq \sigma_0, \quad \sigma_0 \text{ minimal}. \quad (44)$$

Moreover, since $a \notin C$ (see (43)), therefore $\sigma_0 > 1$. Now, applying (41) to the nilpotent element a^{σ_0-1} , we see that

$$a^{\sigma_0-1} - (a^{\sigma_0-1})^n \in C, \quad \text{for some } n = n(a^{\sigma_0-1}) \geq 2. \quad (45)$$

Furthermore, since $(\sigma_0 - 1)n \geq (\sigma_0 - 1)2 \geq \sigma_0$ (since $\sigma_0 \geq 2$), (44) implies that

$$(a^{\sigma_0-1})^n = a^{(\sigma_0-1)n} \in C. \quad (46)$$

Combining (45) and (46), we conclude that $a^{\sigma_0-1} \in C$, which contradicts the minimality of σ_0 in (44). This contradiction proves (42). Since R is a periodic ring satisfying (42), therefore, by a well-known theorem of Herstein [4], R is commutative. This completes the proof. $\square$

COROLLARY 12. *A generalized Boolean ring with central idempotents and commuting nilpotents is commutative.*

This corollary recovers a result proved by the authors in [1].

COROLLARY 13. *If R is a generalized Boolean ring, and if R is 2-torsion-free, then R is nil or commutative.*

PROOF. We claim that all idempotents of R are central. Suppose not, and suppose e is a noncentral idempotent in R . Then $-e \notin (N \cup C)$, and hence (see [Definition 2](#))

$$(-e) - (-e)^n \in C, \quad n \text{ even}. \quad (47)$$

Thus, $2e \in C$, and hence $[2e, x] = 0$ for all x in R . Since R is 2-torsion-free, $2[e, x] = 0$ implies $[e, x] = 0$, and thus $e \in C$, a contradiction. This contradiction proves that all idempotents of R are central, and hence R is nil or commutative, by [Theorem 11](#). $\square$

THEOREM 14. *Let R be a generalized Boolean ring in which every finite subring is either commutative or nil. Then R is either commutative or nil.*

PROOF. By contradiction. Thus, suppose R is a generalized Boolean ring such that every finite subring of R is either commutative or nil. Suppose, further, that R is not commutative and not nil either. By [Theorem 11](#), there must exist a *noncentral* idempotent element e in R , and hence $e \notin (C \cup N)$. Thus (see [Definition 2](#)), since $-e \notin (C \cup N)$,

$$(-e) - (-e)^n \in (N \cap C), \quad n \text{ even.} \quad (48)$$

This implies that $2e \in (N \cap C)$, and hence $(2e)^k = 2^k e = 0$, for some $k \in \mathbb{Z}^+$. Since $e \notin C$, we must have the following:

$$\text{Either } ex - exe \neq 0 \text{ for some } x \in R, \text{ or } x'e - ex'e \neq 0 \text{ for some } x' \in R. \quad (49)$$

Suppose $u = ex - exe \neq 0$. Then,

$$eu = u \neq 0 = ue = u^2, \quad (u = ex - exe \neq 0). \quad (50)$$

Moreover,

$$2u = [2e, ex] = 0 \quad (\text{since } 2e \in C). \quad (51)$$

Furthermore, the subring generated by e and u is

$$\langle e, u \rangle = \{re + su \mid r, s \in \mathbb{Z}\}. \quad (52)$$

Since $2^k e = 0$ and $2u = 0$, the subring $\langle e, u \rangle$ is *finite*. Indeed,

$$\langle e, u \rangle = \{re + su \mid 1 \leq r \leq 2^k, 1 \leq s \leq 2\}. \quad (53)$$

On the other hand, if $x'e - ex'e \neq 0$ for some $x' \in R$ (the only other possibility), then the subring, $\langle e, v \rangle$, generated by e and $v = x'e - ex'e$ is (as is readily verified)

$$\langle e, v \rangle = \{re + sv \mid 1 \leq r \leq 2^k, 1 \leq s \leq 2\}. \quad (54)$$

Again, $\langle e, v \rangle$ is a *finite* subring of R . Hence, in either case, we found a *finite* subring of R , which is neither commutative (since $e \notin C$), nor nil (since $e \notin N$), contradicting our hypothesis. This contradiction proves the theorem. $\square$

REMARK 15. A careful examination of the proof of [Theorem 14](#) shows that we only need to assume that “every subring S , with $|S| = 2^m$ for some positive integer m , is commutative or nil” in order for the ground generalized Boolean ring R to be commutative or nil. Indeed, $|\langle e, u \rangle| = 2^k \cdot 2 = 2^{k+1}$, since the representation of any x in this subring in the form $x = re + su$; $r, s \in \mathbb{Z}$, is *unique*. For, suppose $x = re + su$ and $x = r'e + s'u$. Then, $(r - r')e = (s' - s)u$. Recall that $2u = 0$, and $ue = 0$. Thus, if $s' - s$ is even, then $(r - r')e = 0$, and hence $re = r'e$, $su = s'u$. On the other hand, if $s' - s$ is odd, then $(r - r')e = u$, and hence $(r - r')ee = ue = 0$. Again, we obtain $re = r'e$, $su = s'u$.

We conclude with the following examples.

EXAMPLE 16. Let

$$R = \left\{ \begin{pmatrix} a & b & c \\ 0 & a^2 & 0 \\ 0 & 0 & a \end{pmatrix} : a, b, c \in \text{GF}(4) \right\}. \quad (55)$$

It is readily verified that the idempotents of R are central and

$$x - x^7 = 0, \quad \forall x \in R \setminus (N \cup C), \quad (56)$$

but R is neither nil nor commutative. Hence, [Theorem 11](#) is not true if we drop the hypothesis that “ n is even” in the definition of a generalized Boolean ring.

EXAMPLE 17. Let

$$R = \left\{ \begin{pmatrix} 0 & a & b \\ 0 & 0 & c \\ 0 & 0 & 0 \end{pmatrix} : a, b, c \in \text{GF}(3) \right\}. \quad (57)$$

This example shows that we cannot drop the hypothesis that “ N is commutative” in [Corollary 12](#). (Note that R is not commutative.)

EXAMPLE 18. Let

$$R = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} : 0, 1 \in \text{GF}(2) \right\}. \quad (58)$$

This example shows that we cannot drop the hypothesis that “the idempotents are central” in [Corollary 12](#). (Note that R is not commutative.) This example also shows that we cannot drop the hypothesis that “ R is 2-torsion-free” in [Corollary 13](#). Note that, in this ring R , $x - x^2 = 0$ for all $x \in R \setminus (N \cup C)$. Even more is true. This ring R also shows that we cannot drop the hypothesis that “ $1 \in R$ ” in [Corollary 7](#), nor the hypothesis that “ $1 \in R$ ” in [Corollary 8](#).

Returning to the ring R in [Example 16](#), we see that this ring further shows that we cannot drop the hypothesis that “ m and n are of *opposite* parity” in the definition of a generalized periodic ring in connection with [Corollary 7](#), or the hypothesis that “ n is even” in the definition of a generalized Boolean ring as far as [Corollary 8](#) is concerned. (Recall that $x - x^7 = 0$ for all $x \in R \setminus (N \cup C)$.)

EXAMPLE 19. Let S be any *noncommutative* ring such that $S^3 = (0)$. (For example, we may take S to be the ring of all 3×3 strictly upper triangular matrices over a field F .) Let $R = \text{GF}(4) \oplus S$. It is readily verified that $x^3 = x^6$ for all x in R , and hence R is indeed a generalized periodic ring. Moreover, the only idempotents of R are $(0, 0)$ and $(1, 0)$, and thus the idempotents of R are certainly central. Had R been a generalized Boolean ring, then, by [Theorem 11](#), R would have to be either nil or commutative, which is certainly false here (recall that S is *not* commutative). This example shows that the set of generalized periodic rings is a wider class than that of generalized Boolean rings, and thus [Theorem 11](#) does not hold for generalized periodic rings.

ACKNOWLEDGEMENT. This work was supported by the Natural Sciences and Engineering Research Council of Canada, Grant No. 3961.

REFERENCES

- [1] H. E. Bell and A. Yaquib, *Generalized periodic rings*, Int. J. Math. Math. Sci. **19** (1996), no. 1, 87–92. [MR 96h:16036](#). [Zbl 842.16019](#).
- [2] M. Chacron, *On a theorem of Herstein*, Canad. J. Math. **21** (1969), 1348–1353. [MR 41 #6905](#). [Zbl 213.04302](#).
- [3] I. N. Herstein, *A generalization of a theorem of Jacobson. III*, Amer. J. Math. **75** (1953), 105–111. [MR 14,613e](#). [Zbl 050.02901](#).
- [4] ———, *A note on rings with central nilpotent elements*, Proc. Amer. Math. Soc. **5** (1954), 620. [MR 16,5c](#). [Zbl 055.26003](#).

HOWARD E. BELL: DEPARTMENT OF MATHEMATICS, BROCK UNIVERSITY, STREET CATHARINES, ONTARIO, CANADA L2S 3A1

E-mail address: hbell@spartan.ac.brocku.ca

ADIL YAQUB: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106, USA

E-mail address: yaqub@math.ucsb.edu