FUZZY BCI-SUBALGEBRAS WITH INTERVAL-VALUED MEMBERSHIP FUNCTIONS

SUNG MIN HONG, YOUNG BAE JUN, SEON JEONG KIM, and GWANG IL KIM

(Received 3 May 2000)

ABSTRACT. The purpose of this paper is to define the notion of an interval-valued fuzzy BCI-subalgebra (briefly, an i-v fuzzy BCI-subalgebra) of a BCI-algebra. Necessary and sufficient conditions for an i-v fuzzy set to be an i-v fuzzy BCI-subalgebra are stated. A way to make a new i-v fuzzy BCI-subalgebra from old one is given. The images and inverse images of i-v fuzzy BCI-subalgebras are defined, and how the images or inverse images of i-v fuzzy BCI-subalgebras become i-v fuzzy BCI-subalgebras is studied.

2000 Mathematics Subject Classification. Primary 06F35, 03B52.

1. Introduction. The notion of BCK-algebras was proposed by Iami and Iséki in 1966. In the same year, Iséki [2] introduced the notion of a BCI-algebra which is a generalization of a BCK-algebra. Since then numerous mathematical papers have been written investigating the algebraic properties of the BCK/BCI-algebras and their relationship with other universal structures including lattices and Boolean algebras. Fuzzy sets were initiated by Zadeh [3]. In [4], Zadeh made an extension of the concept of a fuzzy set by an interval-valued fuzzy set (i.e., a fuzzy set with an interval-valued membership function). This interval-valued fuzzy set is referred to as an i-v fuzzy set. In [4], Zadeh also constructed a method of approximate inference using his i-v fuzzy sets. In [1], Biswas defined interval-valued fuzzy subgroups (i.e., i-v fuzzy subgroups) of Rosenfeld's nature, and investigated some elementary properties. In this paper, using the notion of interval-valued fuzzy set by Zadeh, we introduce the concept of an interval-valued fuzzy BCI-subalgebra (briefly, i-v fuzzy BCI-subalgebra) of a BCI-algebra, and study some of their properties. Using an i-v level set of an i-v fuzzy set, we state a characterization of an i-v fuzzy BCI-subalgebra. We prove that every BCI-subalgebra of a BCI-algebra X can be realized as an i-v level BCI-subalgebra of an i-v fuzzy BCI-subalgebra of X . In connection with the notion of homomorphism, we study how the images and inverse images of i-v fuzzy BCI-subalgebras become i-v fuzzy BCI-subalgebras.

2. Preliminaries. In this section, we include some elementary aspects that are necessary for this paper.

Recall that a *BCI-algebra* is an algebra $(X, *, 0)$ of type $(2, 0)$ satisfying the following axioms:

$$(I) \quad ((x * y) * (x * z)) * (z * y) = 0,$$

$$(II) \quad (x * (x * y)) * y = 0,$$

(III) $x * x = 0$, and

(IV) $x * y = 0$ and $y * x = 0$ imply $x = y$,

for every $x, y, z \in X$.

Note that the equality $0 * (x * y) = (0 * x) * (0 * y)$ holds in a BCI-algebra. A non-empty subset S of a BCI-algebra X is called a *BCI-subalgebra* of X if $x * y \in S$ whenever $x, y \in S$. A mapping $f : X \rightarrow Y$ of BCI-algebras is called a *homomorphism* if $f(x * y) = f(x) * f(y)$ for all $x, y \in X$.

We now review some fuzzy logic concepts. Let X be a set. A *fuzzy set* in X is a function $\mu : X \rightarrow [0, 1]$. Let f be a mapping from a set X into a set Y . Let ν be a fuzzy set in Y . Then the *inverse image* of ν , denoted by $f^{-1}[\nu]$, is the fuzzy set in X defined by $f^{-1}[\nu](x) = \nu(f(x))$ for all $x \in X$. Conversely, let μ be a fuzzy set in X . The *image* of μ , written as $f[\mu]$, is a fuzzy set in Y defined by

$$f[\mu](y) = \begin{cases} \sup_{z \in f^{-1}(y)} \mu(z) & \text{if } f^{-1}(y) \neq \emptyset, \\ 0 & \text{otherwise,} \end{cases} \quad (2.1)$$

for all $y \in Y$, where $f^{-1}(y) = \{x \mid f(x) = y\}$.

An *interval-valued fuzzy set* (briefly, *i-v fuzzy set*) A defined on X is given by

$$A = \{(x, [\mu_A^L(x), \mu_A^U(x)])\}, \quad \forall x \in X \text{ (briefly, denoted by } A = [\mu_A^L, \mu_A^U]), \quad (2.2)$$

where μ_A^L and μ_A^U are two fuzzy sets in X such that $\mu_A^L(x) \leq \mu_A^U(x)$ for all $x \in X$.

Let $\bar{\mu}_A(x) = [\mu_A^L(x), \mu_A^U(x)]$, $\forall x \in X$ and let $D[0, 1]$ denotes the family of all closed subintervals of $[0, 1]$. If $\mu_A^L(x) = \mu_A^U(x) = c$, say, where $0 \leq c \leq 1$, then we have $\bar{\mu}_A(x) = [c, c]$ which we also assume, for the sake of convenience, to belong to $D[0, 1]$. Thus $\bar{\mu}_A(x) \in D[0, 1]$, $\forall x \in X$, and therefore the i-v fuzzy set A is given by

$$A = \{(x, \bar{\mu}_A(x))\}, \quad \forall x \in X, \text{ where } \bar{\mu}_A : X \rightarrow D[0, 1]. \quad (2.3)$$

Now let us define what is known as *refined minimum* (briefly, *rmin*) of two elements in $D[0, 1]$. We also define the symbols “ $\geq$ ”, “ $\leq$ ”, and “ $=$ ” in case of two elements in $D[0, 1]$. Consider two elements $D_1 := [a_1, b_1]$ and $D_2 := [a_2, b_2] \in D[0, 1]$. Then

$$\begin{aligned} \text{rmin}(D_1, D_2) &= [\min\{a_1, a_2\}, \min\{b_1, b_2\}]; \\ D_1 \geq D_2 &\text{ if and only if } a_1 \geq a_2, b_1 \geq b_2; \end{aligned} \quad (2.4)$$

and similarly we may have $D_1 \leq D_2$ and $D_1 = D_2$.

DEFINITION 2.1. A fuzzy set μ in a BCI-algebra X is called a *fuzzy BCI-subalgebra* of X if $\mu(x * y) \geq \min\{\mu(x), \mu(y)\}$ for all $x, y \in X$.

3. Interval-valued fuzzy BCI-subalgebras. In what follows, let X denote a BCI-algebra unless otherwise specified. We begin with the following two propositions.

PROPOSITION 3.1. *Let f be a homomorphism from a BCI-algebra X into a BCI-algebra Y . If ν is a fuzzy BCI-subalgebra of Y , then the inverse image $f^{-1}[\nu]$ of ν is a fuzzy BCI-subalgebra of X .*

PROOF. For any $x, y \in X$, we have

$$\begin{aligned} f^{-1}[\nu](x * y) &= \nu(f(x * y)) = \nu(f(x) * f(y)) \\ &\geq \min\{\nu(f(x)), \nu(f(y))\} \\ &= \min\{f^{-1}[\nu](x), f^{-1}[\nu](y)\}. \end{aligned} \quad (3.1)$$

Hence $f^{-1}[\nu]$ is a fuzzy BCI-subalgebra of X . $\square$

PROPOSITION 3.2. *Let $f : X \rightarrow Y$ be a homomorphism between BCI-algebras X and Y . For every fuzzy BCI-subalgebra μ of X , the image $f[\mu]$ of μ is a fuzzy BCI-subalgebra of Y .*

PROOF. We first prove that

$$f^{-1}(y_1) * f^{-1}(y_2) \subseteq f^{-1}(y_1 * y_2) \quad (3.2)$$

for all $y_1, y_2 \in Y$. For, if $x \in f^{-1}(y_1) * f^{-1}(y_2)$, then $x = x_1 * x_2$ for some $x_1 \in f^{-1}(y_1)$ and $x_2 \in f^{-1}(y_2)$. Since f is a homomorphism, it follows that $f(x) = f(x_1 * x_2) = f(x_1) * f(x_2) = y_1 * y_2$ so that $x \in f^{-1}(y_1 * y_2)$. Hence (3.2) holds. Now let $y_1, y_2 \in Y$ be arbitrarily given. Assume that $y_1 * y_2 \notin \text{Im}(f)$. Then $f[\mu](y_1 * y_2) = 0$. But if $y_1 * y_2 \in \text{Im}(f)$, that is, $f^{-1}(y_1 * y_2) \neq \emptyset$, then $f^{-1}(y_1) \neq \emptyset$ or $f^{-1}(y_2) \neq \emptyset$ by (3.2). Thus $f[\mu](y_1) = 0$ or $f[\mu](y_2) = 0$, and so

$$f[\mu](y_1 * y_2) = 0 = \min\{f[\mu](y_1), f[\mu](y_2)\}. \quad (3.3)$$

Suppose that $f^{-1}(y_1 * y_2) \neq \emptyset$. Then we should consider the two cases:

$$f^{-1}(y_1) = \emptyset \quad \text{or} \quad f^{-1}(y_2) = \emptyset, \quad (3.4)$$

$$f^{-1}(y_1) \neq \emptyset \quad \text{and} \quad f^{-1}(y_2) \neq \emptyset. \quad (3.5)$$

For the case (3.4), we have $f[\mu](y_1) = 0$ or $f[\mu](y_2) = 0$, and so

$$f[\mu](y_1 * y_2) \geq 0 = \min\{f[\mu](y_1), f[\mu](y_2)\}. \quad (3.6)$$

Case (3.5) implies, from (3.2), that

$$\begin{aligned} f[\mu](y_1 * y_2) &= \sup_{z \in f^{-1}(y_1 * y_2)} \mu(z) \geq \sup_{z \in f^{-1}(y_1) * f^{-1}(y_2)} \mu(z) \\ &= \sup_{x_1 \in f^{-1}(y_1), x_2 \in f^{-1}(y_2)} \mu(x_1 * x_2). \end{aligned} \quad (3.7)$$

Since μ is a fuzzy BCI-subalgebra of X , it follows from the definition of a fuzzy BCI-subalgebra that

$$\begin{aligned}
 f[\mu](y_1 * y_2) &\geq \sup_{x_1 \in f^{-1}(y_1), x_2 \in f^{-1}(y_2)} \min \{ \mu(x_1), \mu(x_2) \} \\
 &= \sup_{x_1 \in f^{-1}(y_1)} \left(\min \left\{ \sup_{x_2 \in f^{-1}(y_2)} \mu(x_1), \mu(x_2) \right\} \right) \\
 &= \sup_{x_1 \in f^{-1}(y_1)} \left(\min \left\{ \mu(x_1), \sup_{x_2 \in f^{-1}(y_2)} \mu(x_2) \right\} \right) \\
 &= \sup_{x_1 \in f^{-1}(y_1)} (\min \{ \mu(x_1), f[\mu](y_2) \}) \\
 &= \min \left\{ \sup_{x_1 \in f^{-1}(y_1)} \mu(x_1), f[\mu](y_2) \right\} \\
 &= \min \{ f[\mu](y_1), f[\mu](y_2) \}.
 \end{aligned} \tag{3.8}$$

Hence $f[\mu](y_1 * y_2) \geq \min \{ f[\mu](y_1), f[\mu](y_2) \}$ for all $y_1, y_2 \in Y$. This completes the proof. $\square$

DEFINITION 3.3. An i-v fuzzy set A in X is called an *interval-valued fuzzy BCI-subalgebra* (briefly, *i-v fuzzy BCI-subalgebra*) of X if

$$\bar{\mu}_A(x * y) \geq \min \{ \bar{\mu}_A(x), \bar{\mu}_A(y) \} \quad \forall x, y \in X. \tag{3.9}$$

EXAMPLE 3.4. Let $X = \{0, a, b, c\}$ be a BCI-algebra with the following Cayley table:

TABLE 3.1.

$*$	0	a	b	c
0	0	c	0	a
a	a	0	a	c
b	b	c	0	a
c	c	a	c	0

let an i-v fuzzy set A defined on X be given by

$$\bar{\mu}_A(x) = \begin{cases} [0.2, 0.8] & \text{if } x \in \{0, b\}, \\ [0.1, 0.7] & \text{otherwise.} \end{cases} \tag{3.10}$$

It is easy to check that A is an i-v fuzzy BCI-subalgebra of X .

LEMMA 3.5. If A is an i-v fuzzy BCI-subalgebra of X , then $\bar{\mu}_A(0) \geq \bar{\mu}_A(x)$ for all $x \in X$.

PROOF. For every $x \in X$, we have

$$\begin{aligned}\bar{\mu}_A(0) &= \bar{\mu}_A(x * x) \geq \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(x)\} \\ &= \text{rmin}\{[\mu_A^L(x), \mu_A^U(x)], [\mu_A^L(x), \mu_A^U(x)]\} \\ &= [\mu_A^L(x), \mu_A^U(x)] = \bar{\mu}_A(x),\end{aligned}\tag{3.11}$$

this completes the proof. $\square$

THEOREM 3.6. *Let A be an i -v fuzzy BCI-subalgebra of X . If there is a sequence $\{x_n\}$ in X such that*

$$\lim_{n \rightarrow \infty} \bar{\mu}_A(x_n) = [1, 1],\tag{3.12}$$

then $\bar{\mu}_A(0) = [1, 1]$.

PROOF. Since $\bar{\mu}_A(0) \geq \bar{\mu}_A(x)$ for all $x \in X$, we have $\bar{\mu}_A(0) \geq \bar{\mu}_A(x_n)$ for every positive integer n . Note that

$$[1, 1] \geq \bar{\mu}_A(0) \geq \lim_{n \rightarrow \infty} \bar{\mu}_A(x_n) = [1, 1].\tag{3.13}$$

Hence $\bar{\mu}_A(0) = [1, 1]$. $\square$

THEOREM 3.7. *An i -v fuzzy set $A = [\mu_A^L, \mu_A^U]$ in X is an i -v fuzzy BCI-subalgebra of X if and only if μ_A^L and μ_A^U are fuzzy BCI-subalgebras of X .*

PROOF. Suppose that μ_A^L and μ_A^U are fuzzy BCI-subalgebras of X . Let $x, y \in X$. Then

$$\begin{aligned}\bar{\mu}_A(x * y) &= [\mu_A^L(x * y), \mu_A^U(x * y)] \\ &\geq [\min\{\mu_A^L(x), \mu_A^L(y)\}, \min\{\mu_A^U(x), \mu_A^U(y)\}] \\ &= \text{rmin}\{[\mu_A^L(x), \mu_A^U(x)], [\mu_A^L(y), \mu_A^U(y)]\} \\ &= \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\}.\end{aligned}\tag{3.14}$$

Hence A is an i -v fuzzy BCI-subalgebra of X .

Conversely, assume that A is an i -v fuzzy BCI-subalgebra of X . For any $x, y \in X$, we have

$$\begin{aligned}[\mu_A^L(x * y), \mu_A^U(x * y)] &= \bar{\mu}_A(x * y) \geq \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\} \\ &= \text{rmin}\{[\mu_A^L(x), \mu_A^U(x)], [\mu_A^L(y), \mu_A^U(y)]\} \\ &= [\min\{\mu_A^L(x), \mu_A^L(y)\}, \min\{\mu_A^U(x), \mu_A^U(y)\}].\end{aligned}\tag{3.15}$$

It follows that $\mu_A^L(x * y) \geq \min\{\mu_A^L(x), \mu_A^L(y)\}$ and $\mu_A^U(x * y) \geq \min\{\mu_A^U(x), \mu_A^U(y)\}$. Hence μ_A^L and μ_A^U are fuzzy BCI-subalgebras of X . $\square$

THEOREM 3.8. *Let A be an i -v fuzzy set in X . Then A is an i -v fuzzy BCI-subalgebra of X if and only if the nonempty set*

$$U(A; [\delta_1, \delta_2]) := \{x \in X \mid \bar{\mu}_A(x) \geq [\delta_1, \delta_2]\}\tag{3.16}$$

is a BCI-subalgebra of X for every $[\delta_1, \delta_2] \in D[0, 1]$.

We then call $\tilde{U}(A; [\delta_1, \delta_2])$ the *i-v level BCI-subalgebra* of A .

PROOF. Assume that A is an i-v fuzzy BCI-subalgebra of X and let $[\delta_1, \delta_2] \in D[0, 1]$ be such that $x, y \in \tilde{U}(A; [\delta_1, \delta_2])$. Then

$$\bar{\mu}_A(x * y) \geq \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\} \geq \text{rmin}\{[\delta_1, \delta_2], [\delta_1, \delta_2]\} = [\delta_1, \delta_2], \quad (3.17)$$

and so $x * y \in \tilde{U}(A; [\delta_1, \delta_2])$. Thus $\tilde{U}(A; [\delta_1, \delta_2])$ is a BCI-subalgebra of X .

Conversely, assume that $\tilde{U}(A; [\delta_1, \delta_2]) (\neq \emptyset)$ is a BCI-subalgebra of X for every $[\delta_1, \delta_2] \in D[0, 1]$. Suppose there exist $x_0, y_0 \in X$ such that

$$\bar{\mu}_A(x_0 * y_0) < \text{rmin}\{\bar{\mu}_A(x_0), \bar{\mu}_A(y_0)\}. \quad (3.18)$$

Let $\bar{\mu}_A(x_0) = [y_1, y_2]$, $\bar{\mu}_A(y_0) = [y_3, y_4]$, and $\bar{\mu}_A(x_0 * y_0) = [\delta_1, \delta_2]$. Then

$$[\delta_1, \delta_2] < \text{rmin}\{[y_1, y_2], [y_3, y_4]\} = [\min\{y_1, y_3\}, \min\{y_2, y_4\}]. \quad (3.19)$$

Hence $\delta_1 < \min\{y_1, y_3\}$ and $\delta_2 < \min\{y_2, y_4\}$. Taking

$$[\lambda_1, \lambda_2] = \frac{1}{2}(\bar{\mu}_A(x_0 * y_0) + \text{rmin}\{\bar{\mu}_A(x_0), \bar{\mu}_A(y_0)\}), \quad (3.20)$$

we obtain

$$\begin{aligned} [\lambda_1, \lambda_2] &= \frac{1}{2}([\delta_1, \delta_2] + [\min\{y_1, y_3\}, \min\{y_2, y_4\}]) \\ &= \left[\frac{1}{2}(\delta_1 + \min\{y_1, y_3\}), \frac{1}{2}(\delta_2 + \min\{y_2, y_4\}) \right]. \end{aligned} \quad (3.21)$$

It follows that

$$\begin{aligned} \min\{y_1, y_3\} &> \lambda_1 = \frac{1}{2}(\delta_1 + \min\{y_1, y_3\}) > \delta_1, \\ \min\{y_2, y_4\} &> \lambda_2 = \frac{1}{2}(\delta_2 + \min\{y_2, y_4\}) > \delta_2 \end{aligned} \quad (3.22)$$

so that $[\min\{y_1, y_3\}, \min\{y_2, y_4\}] > [\lambda_1, \lambda_2] > [\delta_1, \delta_2] = \bar{\mu}_A(x_0 * y_0)$. Therefore, $x_0 * y_0 \notin \tilde{U}(A; [\lambda_1, \lambda_2])$. On the other hand,

$$\begin{aligned} \bar{\mu}_A(x_0) &= [y_1, y_2] \geq [\min\{y_1, y_3\}, \min\{y_2, y_4\}] > [\lambda_1, \lambda_2], \\ \bar{\mu}_A(y_0) &= [y_3, y_4] \geq [\min\{y_1, y_3\}, \min\{y_2, y_4\}] > [\lambda_1, \lambda_2], \end{aligned} \quad (3.23)$$

and so $x_0, y_0 \in \tilde{U}(A; [\lambda_1, \lambda_2])$. It contradicts that $\tilde{U}(A; [\lambda_1, \lambda_2])$ is a BCI-subalgebra of X . Hence $\bar{\mu}_A(x * y) \geq \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\}$ for all $x, y \in X$. This completes the proof. $\square$

THEOREM 3.9. Every BCI-subalgebra of X can be realized as an i-v level BCI-subalgebra of an i-v fuzzy BCI-subalgebra of X .

PROOF. Let Y be a BCI-subalgebra of X and let A be an i-v fuzzy set on X defined by

$$\bar{\mu}_A(x) = \begin{cases} [\alpha_1, \alpha_2] & \text{if } x \in Y, \\ [0, 0] & \text{otherwise,} \end{cases} \quad (3.24)$$

where $\alpha_1, \alpha_2 \in (0, 1]$ with $\alpha_1 < \alpha_2$. It is clear that $\tilde{U}(A; [\alpha_1, \alpha_2]) = Y$. We show that A

is an i-v fuzzy BCI-subalgebra of X . Let $x, y \in X$. If $x, y \in Y$, then $x * y \in Y$ and so

$$\bar{\mu}_A(x * y) = [\alpha_1, \alpha_2] = \text{rmin}\{[\alpha_1, \alpha_2], [\alpha_1, \alpha_2]\} = \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\}. \quad (3.25)$$

If $x, y \notin Y$, then $\bar{\mu}_A(x) = [0, 0] = \bar{\mu}_A(y)$ and thus

$$\bar{\mu}_A(x * y) \geq [0, 0] = \text{rmin}\{[0, 0], [0, 0]\} = \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\}. \quad (3.26)$$

If $x \in Y$ and $y \notin Y$, then $\bar{\mu}_A(x) = [\alpha_1, \alpha_2]$ and $\bar{\mu}_A(y) = [0, 0]$. It follows that

$$\bar{\mu}_A(x * y) \geq [0, 0] = \text{rmin}\{[\alpha_1, \alpha_2], [0, 0]\} = \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\}. \quad (3.27)$$

Similarly for the case $x \notin Y$ and $y \in Y$, we get $\bar{\mu}_A(x * y) \geq \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\}$. Therefore A is an i-v fuzzy BCI-subalgebra of X , and the proof is complete. $\square$

THEOREM 3.10. *Let Y be a subset of X and let A be an i-v fuzzy set on X which is given in the proof of Theorem 3.9. If A is an i-v fuzzy BCI-subalgebra of X , then Y is a BCI-subalgebra of X .*

PROOF. Assume that A is an i-v fuzzy BCI-subalgebra of X . Let $x, y \in Y$. Then $\bar{\mu}_A(x) = [\alpha_1, \alpha_2] = \bar{\mu}_A(y)$, and so

$$\bar{\mu}_A(x * y) \geq \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\} = \text{rmin}\{[\alpha_1, \alpha_2], [\alpha_1, \alpha_2]\} = [\alpha_1, \alpha_2]. \quad (3.28)$$

This implies that $x * y \in Y$. Hence Y is a BCI-subalgebra of X . $\square$

THEOREM 3.11. *If A is an i-v fuzzy BCI-subalgebra of X , then the set*

$$X_{\bar{\mu}_A} := \{x \in X \mid \bar{\mu}_A(x) = \bar{\mu}_A(0)\} \quad (3.29)$$

is a BCI-subalgebra of X .

PROOF. Let $x, y \in X_{\bar{\mu}_A}$. Then $\bar{\mu}_A(x) = \bar{\mu}_A(0) = \bar{\mu}_A(y)$, and so

$$\bar{\mu}_A(x * y) \geq \text{rmin}\{\bar{\mu}_A(x), \bar{\mu}_A(y)\} = \text{rmin}\{\bar{\mu}_A(0), \bar{\mu}_A(0)\} = \bar{\mu}_A(0). \quad (3.30)$$

Combining this and Lemma 3.5, we get $\bar{\mu}_A(x * y) = \bar{\mu}_A(0)$, that is, $x * y \in X_{\bar{\mu}_A}$. Hence $X_{\bar{\mu}_A}$ is a BCI-subalgebra of X . $\square$

The following is a way to make a new i-v fuzzy BCI-subalgebra from old one.

THEOREM 3.12. *For an i-v fuzzy BCI-subalgebra A of X , the i-v fuzzy set A^* in X defined by $\bar{\mu}_{A^*}(x) = \bar{\mu}_A(0 * x)$ for all $x \in X$ is an i-v fuzzy BCI-subalgebra of X .*

PROOF. Since the equality $0 * (x * y) = (0 * x) * (0 * y)$ holds for all $x, y \in X$, we have

$$\begin{aligned} \bar{\mu}_{A^*}(x * y) &= \bar{\mu}_A(0 * (x * y)) = \bar{\mu}_A((0 * x) * (0 * y)) \\ &\geq \text{rmin}\{\bar{\mu}_A(0 * x), \bar{\mu}_A(0 * y)\} \\ &= \text{rmin}\{\bar{\mu}_{A^*}(x), \bar{\mu}_{A^*}(y)\} \end{aligned} \quad (3.31)$$

for all $x, y \in X$. Therefore A^* is an i-v fuzzy BCI-subalgebra of X . $\square$

DEFINITION 3.13 (Biswas [1]). Let f be a mapping from a set X into a set Y . Let B be an i-v fuzzy set in Y . Then the *inverse image* of B , denoted by $f^{-1}[B]$, is the i-v fuzzy set in X with the membership function given by $\bar{\mu}_{f^{-1}[B]}(x) = \bar{\mu}_B(f(x))$ for all $x \in X$.

LEMMA 3.14 (Biswas [1]). Let f be a mapping from a set X into a set Y . Let $m = [m^L, m^U]$ and $n = [n^L, n^U]$ be i-v fuzzy sets in X and Y , respectively. Then

- (i) $f^{-1}(n) = [f^{-1}(n^L), f^{-1}(n^U)]$,
- (ii) $f(m) = [f(m^L), f(m^U)]$.

THEOREM 3.15. Let f be a homomorphism from a BCI-algebra X into a BCI-algebra Y . If B is an i-v fuzzy BCI-subalgebra of Y , then the inverse image $f^{-1}[B]$ of B is an i-v fuzzy BCI-subalgebra of X .

PROOF. Since $B = [\mu_B^L, \mu_B^U]$ is an i-v fuzzy BCI-subalgebra of Y , it follows from Theorem 3.7 that μ_B^L and μ_B^U are fuzzy BCI-subalgebras of Y . Using Proposition 3.1, we know that $f^{-1}[\mu_B^L]$ and $f^{-1}[\mu_B^U]$ are fuzzy BCI-subalgebras of X . Hence, by Lemma 3.14 and Theorem 3.7, we conclude that $f^{-1}[B] = [f^{-1}[\mu_B^L], f^{-1}[\mu_B^U]]$ is an i-v fuzzy BCI-subalgebra of X . $\square$

DEFINITION 3.16 (Biswas [1]). Let f be a mapping from a set X into a set Y . Let A be an i-v fuzzy set in X . Then the *image* of A , denoted by $f[A]$, is the i-v fuzzy set in Y with the membership function defined by

$$\bar{\mu}_{f[A]}(y) = \begin{cases} \text{rsup}_{z \in f^{-1}(y)} \bar{\mu}_A(z) & \text{if } f^{-1}(y) \neq \emptyset, \forall y \in Y, \\ [0, 0] & \text{otherwise,} \end{cases} \quad (3.32)$$

where $f^{-1}(y) = \{x \mid f(x) = y\}$.

THEOREM 3.17. Let f be a homomorphism from a BCI-algebra X into a BCI-algebra Y . If A is an i-v fuzzy BCI-subalgebra of X , then the image $f[A]$ of A is an i-v fuzzy BCI-subalgebra of Y .

PROOF. Assume that A is an i-v fuzzy BCI-subalgebra of X . Note that $A = [\mu_A^L, \mu_A^U]$ is an i-v fuzzy BCI-subalgebra of X if and only if μ_A^L and μ_A^U are fuzzy BCI-subalgebras of X . It follows from Proposition 3.2 that the images $f[\mu_A^L]$ and $f[\mu_A^U]$ are fuzzy BCI-subalgebras of Y . Combining Theorem 3.7 and Lemma 3.14, we conclude that $f[A] = [f[\mu_A^L], f[\mu_A^U]]$ is an i-v fuzzy BCI-subalgebra of Y . $\square$

ACKNOWLEDGEMENT. This work was supported by Korea Research Foundation Grant KRF-99-005-D00003.

REFERENCES

- [1] R. Biswas, *Rosenfeld's fuzzy subgroups with interval-valued membership functions*, Fuzzy Sets and Systems **63** (1994), no. 1, 87–90. CMP 1 273 001. Zbl 844.20060.
- [2] K. Iséki, *An algebra related with a propositional calculus*, Proc. Japan Acad. **42** (1966), 26–29. MR 34#2433. Zbl 207.29304.
- [3] L. A. Zadeh, *Fuzzy sets*, Information and Control **8** (1965), 338–353. MR 36#2509. Zbl 139.24606.

- [4] ———, *The concept of a linguistic variable and its application to approximate reasoning. I*, Information Sci. **8** (1975), 199–249. MR 52#7225a. Zbl 397.68071.

SUNG MIN HONG: DEPARTMENT OF MATHEMATICS, GYEONGSANG NATIONAL UNIVERSITY, CHINJU 660-701, KOREA

E-mail address: smhong@nongae.gsnu.ac.kr

YOUNG BAE JUN: DEPARTMENT OF MATHEMATICS EDUCATION, GYEONGSANG NATIONAL UNIVERSITY, CHINJU 660-701, KOREA

E-mail address: ybjun@nongae.gsnu.ac.kr

SEON JEONG KIM: DEPARTMENT OF MATHEMATICS, GYEONGSANG NATIONAL UNIVERSITY, CHINJU 660-701, KOREA

E-mail address: skim@nongae.gsnu.ac.kr

GWANG IL KIM: DEPARTMENT OF MATHEMATICS, GYEONGSANG NATIONAL UNIVERSITY, CHINJU 660-701, KOREA

E-mail address: gikim@nongae.gsnu.ac.kr