ON n -FOLD FUZZY POSITIVE IMPLICATIVE IDEALS OF BCK-ALGEBRAS

YOUNG BAE JUN and KYUNG HO KIM

(Received 11 March 2000)

ABSTRACT. We consider the fuzzification of the notion of an n -fold positive implicative ideal. We give characterizations of an n -fold fuzzy positive implicative ideal. We establish the extension property for n -fold fuzzy positive implicative ideals, and state a characterization of PI^n -Noetherian BCK-algebras. Finally we study the normalization of n -fold fuzzy positive implicative ideals.

2000 Mathematics Subject Classification. 06F35, 03G25, 03E72.

1. Introduction. For the general development of BCK-algebras, the ideal theory plays an important role. In 1999, Huang and Chen [1] introduced the notion of n -fold positive implicative ideals in BCK-algebras. In this paper, we consider the fuzzification of n -fold positive implicative ideals in BCK-algebras. We first define the notion of n -fold fuzzy positive implicative ideals of BCK-algebras, and then discuss the related properties. We give the relation between a fuzzy ideal and an n -fold fuzzy positive implicative ideal. We state a condition for a fuzzy ideal to be an n -fold fuzzy positive implicative ideal. Using level sets, we give a characterization of an n -fold fuzzy positive implicative ideal. We establish the extension property for an n -fold fuzzy positive implicative ideal. Using a family of n -fold fuzzy positive implicative ideals, we make a new n -fold fuzzy positive implicative ideal. We define the notion of PI^n -Noetherian BCK-algebras, and give its characterization. Furthermore, we study the normalization of an n -fold fuzzy positive implicative ideal.

2. Preliminaries. By a *BCK-algebra* we mean an algebra $(X; *, 0)$ of type $(2, 0)$ satisfying the axioms

- (I) $((x * y) * (x * z)) * (z * y) = 0$,
- (II) $(x * (x * y)) * y = 0$,
- (III) $x * x = 0$,
- (IV) $0 * x = 0$,
- (V) $x * y = 0$ and $y * x = 0$ imply $x = y$,

for all $x, y, z \in X$. We can define a partial ordering $\leq$ on X by $x \leq y$ if and only if $x * y = 0$. A BCK-algebra X is said to be *n -fold positive implicative* (see Huang and Chen [1]) if there exists a natural number n such that $x * y^{n+1} = x * y^n$ for all $x, y \in X$.

In any BCK-algebra X , the following hold:

- (P1) $x * 0 = x$,
- (P2) $x * y \leq x$,
- (P3) $(x * y) * z = (x * z) * y$,

$$(P4) \quad (x * z) * (y * z) \leq x * y,$$

$$(P5) \quad x \leq y \text{ implies } x * z \leq y * z \text{ and } z * y \leq z * x.$$

Throughout this paper X will always mean a BCK-algebra unless otherwise specified.

A nonempty subset I of X is called an *ideal* of X if it satisfies

$$(I1) \quad 0 \in I,$$

$$(I2) \quad x * y \in I \text{ and } y \in I \text{ imply } x \in I.$$

A nonempty subset I of X is said to be a *positive implicative ideal* if it satisfies

$$(I1) \quad 0 \in I,$$

$$(I3) \quad (x * y) * z \in I \text{ and } y * z \in I \text{ imply } x * z \in I.$$

THEOREM 2.1 (see [3, Theorem 3]). *A nonempty subset I of X is a positive implicative ideal of X if and only if it satisfies*

$$(I1) \quad 0 \in I,$$

$$(I4) \quad ((x * y) * y) * z \in I \text{ and } z \in I \text{ imply } x * y \in I.$$

We now review some fuzzy logic concepts. A fuzzy set in a set X is a function $\mu : X \rightarrow [0, 1]$. For a fuzzy set μ in X and $t \in [0, 1]$ define $U(\mu; t)$ to be the set $U(\mu; t) := \{x \in X \mid \mu(x) \geq t\}$.

A fuzzy set μ in X is said to be a *fuzzy ideal* of X if

$$(F1) \quad \mu(0) \geq \mu(x) \text{ for all } x \in X,$$

$$(F2) \quad \mu(x) \geq \min\{\mu(x * y), \mu(y)\} \text{ for all } x, y \in X.$$

Note that every fuzzy ideal μ of X is order reversing, that is, if $x \leq y$ then $\mu(x) \geq \mu(y)$.

A fuzzy set μ in X is called a *fuzzy positive implicative ideal* of X if it satisfies

$$(F1) \quad \mu(0) \geq \mu(x) \text{ for all } x \in X,$$

$$(F3) \quad \mu(x * z) \geq \min\{\mu((x * y) * z), \mu(y * z)\} \text{ for all } x, y, z \in X.$$

THEOREM 2.2 (see [2, Proposition 1]). *For any fuzzy ideal μ of X , we have*

$$\mu(x * y) \geq \mu((x * y) * y) \iff \mu((x * z) * (y * z)) \geq \mu((x * y) * z) \quad \forall x, y, z \in X. \quad (2.1)$$

3. n -fold fuzzy positive implicative ideals. For any elements x and y of a BCK-algebra, $x * y^n$ denotes

$$(\cdots((x * y) * y) * \cdots) * y \quad (3.1)$$

in which y occurs n times. Using Theorem 2.1, Huang and Chen [1] introduced the concept of an n -fold positive implicative ideal as follows.

DEFINITION 3.1. A subset A of X is called an *n -fold positive implicative ideal* of X if

$$(I1) \quad 0 \in A,$$

$$(I5) \quad x * y^n \in A \text{ whenever } (x * y^{n+1}) * z \in A \text{ and } z \in A \text{ for every } x, y, z \in X.$$

We try to fuzzify the concept of n -fold positive implicative ideal.

DEFINITION 3.2. Let n be a positive integer. A fuzzy set μ in X is called an *n -fold fuzzy positive implicative ideal* of X if

$$(F1) \quad \mu(0) \geq \mu(x) \text{ for all } x \in X,$$

(F4) $\mu(x * y^n) \geq \min\{\mu((x * y^{n+1}) * z), \mu(z)\}$ for all $x, y, z \in X$.

Notice that the 1-fold fuzzy positive implicative ideal is a fuzzy positive implicative ideal.

EXAMPLE 3.3. Let $X = \{0, a, b\}$ be a BCK-algebra with the following Cayley table:

$*$	0	a	b
0	0	0	0
a	a	0	0
b	b	b	0

Define a fuzzy set $\mu : X \rightarrow [0, 1]$ by $\mu(0) = t_0$, $\mu(a) = t_1$, and $\mu(b) = t_2$ where $t_0 > t_1 > t_2$ in $[0, 1]$. Then μ is an n -fold fuzzy positive implicative ideal of X for every natural number n .

PROPOSITION 3.4. Every n -fold fuzzy positive implicative ideal is a fuzzy ideal for every natural number n .

PROOF. Let μ be an n -fold fuzzy positive implicative ideal of X . Then

$$\begin{aligned} \mu(x) &= \mu(x * 0^n) \geq \min\{\mu((x * 0^{n+1}) * z), \mu(z)\} \\ &= \min\{\mu(x * z), \mu(z)\} \quad \forall x, z \in X. \end{aligned} \quad (3.2)$$

Hence μ is a fuzzy ideal of X . □

The following example shows that the converse of [Proposition 3.4](#) may not be true.

EXAMPLE 3.5. Let $X = \mathbb{N} \cup \{0\}$, where $\mathbb{N}$ is the set of natural numbers, in which the operation $*$ is defined by $x * y = \max\{0, x - y\}$ for all $x, y \in X$. Then X is a BCK-algebra [[1](#), Example 1.3]. Let μ be a fuzzy set in X given by $\mu(0) = t_0 > t_1 = \mu(x)$ for all $x (\neq 0) \in X$. Then μ is a fuzzy ideal of X . But μ is not a 2-fold fuzzy positive implicative ideal of X because $\mu(5 * 2^2) = \mu(1) = t_1$ and $\mu((5 * 2^3) * 0) = \mu(0) = t_0$, and so

$$\mu(5 * 2^2) \not\geq \min\{\mu((5 * 2^3) * 0), \mu(0)\}. \quad (3.3)$$

Let X be an n -fold positive implicative BCK-algebra and let μ be a fuzzy ideal of X . For any $x, y, z \in X$ we have

$$\mu(x * y^n) = \mu(x * y^{n+1}) \geq \min\{\mu((x * y^{n+1}) * z), \mu(z)\}. \quad (3.4)$$

Hence μ is an n -fold fuzzy positive implicative ideal of X . Combining this and [Proposition 3.4](#), we have the following theorem.

THEOREM 3.6. In an n -fold positive implicative BCK-algebra, the notion of n -fold fuzzy positive implicative ideals and fuzzy ideals coincide.

PROPOSITION 3.7. Let μ be a fuzzy ideal of X . Then μ is an n -fold fuzzy positive implicative ideal of X if and only if it satisfies the inequality $\mu(x * y^n) \geq \mu(x * y^{n+1})$ for all $x, y \in X$.

PROOF. Suppose that μ is an n -fold fuzzy positive implicative ideal of X and let $x, y \in X$. Then

$$\begin{aligned}\mu(x * y^n) &\geq \min\{\mu((x * y^{n+1}) * 0), \mu(0)\} \\ &= \min\{\mu(x * y^{n+1}), \mu(0)\} \\ &= \mu(x * y^{n+1}).\end{aligned}\tag{3.5}$$

Conversely, let μ be a fuzzy ideal of X satisfying the inequality

$$\mu(x * y^n) \geq \mu(x * y^{n+1}) \quad \forall x, y \in X.\tag{3.6}$$

Then

$$\mu(x * y^n) \geq \mu(x * y^{n+1}) \geq \min\{\mu((x * y^{n+1}) * z), \mu(z)\} \quad \forall x, y, z \in X.\tag{3.7}$$

Hence μ is an n -fold fuzzy positive implicative ideal of X . $\square$

COROLLARY 3.8. Every n -fold fuzzy positive implicative ideal μ of X satisfies the inequality $\mu(x * y^n) \geq \mu(x * y^{n+k})$ for all $x, y \in X$ and $k \in \mathbb{N}$.

PROOF. Using Proposition 3.7, the proof is straightforward by induction. $\square$

LEMMA 3.9. Let A be a nonempty subset of X and let μ be a fuzzy set in X defined by

$$\mu(x) := \begin{cases} t_1 & \text{if } x \in A, \\ t_2 & \text{otherwise,} \end{cases}\tag{3.8}$$

where $t_1 > t_2$ in $[0, 1]$. Then μ is a fuzzy ideal of X if and only if A is an ideal of X .

PROOF. Let A be an ideal of X . Since $0 \in A$, therefore $\mu(0) = t_1 \geq \mu(x)$ for all $x \in X$. Suppose that (F2) does not hold. Then there exist $a, b \in X$ such that $\mu(a) = t_2$ and $\min\{\mu(a * b), \mu(b)\} = t_1$. Thus $\mu(a * b) = t_1 = \mu(b)$, and so $a * b \in A$ and $b \in A$. It follows from (I2) that $a \in A$ so that $\mu(a) = t_1$. This is a contradiction. Suppose that μ is a fuzzy ideal of X . Since $\mu(0) \geq \mu(x)$ for all $x \in X$, we have $\mu(0) = t_1$ and hence $0 \in A$. Let $x, y \in X$ be such that $x * y \in A$ and $y \in A$. Using (F2), we get $\mu(x) \geq \min\{\mu(x * y), \mu(y)\} = t_1$ and so $\mu(x) = t_1$, that is, $x \in A$. Consequently, A is an ideal of X . $\square$

PROPOSITION 3.10. Let A be a nonempty subset of X , n a positive integer, and μ a fuzzy set in X defined as follows:

$$\mu(x) := \begin{cases} t_1 & \text{if } x \in A, \\ t_2 & \text{otherwise,} \end{cases}\tag{3.9}$$

where $t_1 > t_2$ in $[0, 1]$. Then μ is an n -fold fuzzy positive implicative ideal of X if and only if A is an n -fold positive implicative ideal of X .

PROOF. Assume that μ is an n -fold fuzzy positive implicative ideal of X . Then μ is a fuzzy ideal of X . It follows from Lemma 3.9 that A is an ideal of X . Let $x, y \in X$ be such that $x * y^{n+1} \in A$. Using Proposition 3.7, we get $\mu(x * y^n) \geq \mu(x * y^{n+1}) = t_1$ and so

$\mu(x * y^n) = t_1$, that is, $x * y^n \in A$. Hence by [1, Theorem 1.5], we conclude that A is an n -fold positive implicative ideal of X . Conversely, suppose that A is an n -fold positive implicative ideal of X . Then A is an ideal of X (see [1, Proposition 1.2]). It follows from Lemma 3.9 that μ is a fuzzy ideal of X . For any $x, y \in X$, either $x * y^n \in A$ or $x * y^n \notin A$. The former induces $\mu(x * y^n) = t_1 \geq \mu(x * y^{n+1})$. In the latter, we know that $x * y^{n+1} \notin A$ by [1, Theorem 1.5]. Hence $\mu(x * y^n) = t_2 = \mu(x * y^{n+1})$. From Proposition 3.7 it follows that μ is an n -fold fuzzy positive implicative ideal of X . $\square$

PROPOSITION 3.11. *A fuzzy set μ in X is an n -fold fuzzy positive implicative ideal of X if and only if it satisfies*

$$(F1) \quad \mu(0) \geq \mu(x),$$

$$(F5) \quad \mu(x * z^n) \geq \min\{\mu((x * y) * z^n), \mu(y * z^n)\}, \text{ for all } x, y, z \in X.$$

PROOF. Suppose that μ is an n -fold fuzzy positive implicative ideal of X and let $x, y, z \in X$. Then μ is a fuzzy ideal of X (see Proposition 3.4), and so μ is order reversing. It follows from (P3), (P4), and (P5) that

$$\mu((x * z^{2n}) * (y * z^n)) = \mu(((x * z^n) * (y * z^n)) * z^n) \geq \mu((x * y) * z^n). \quad (3.10)$$

Using (F2) and Corollary 3.8, we get

$$\begin{aligned} \mu(x * z^n) &\geq \mu(x * z^{2n}) \geq \min\{\mu((x * z^{2n}) * (y * z^n)), \mu(y * z^n)\} \\ &\geq \min\{\mu((x * y) * z^n), \mu(y * z^n)\}, \end{aligned} \quad (3.11)$$

which proves (F5). Conversely, assume that μ satisfies conditions (F1) and (F5). Taking $z = 0$ in (F5) and using (P1), we conclude that

$$\begin{aligned} \mu(x) &= \mu(x * 0) \geq \min\{\mu((x * y) * 0^n), \mu(y * 0^n)\} \\ &= \min\{\mu(x * y), \mu(y)\}. \end{aligned} \quad (3.12)$$

Hence μ is a fuzzy ideal of X . Putting $z = y$ in (F5) and applying (III), (IV), and (F1), we have

$$\begin{aligned} \mu(x * y^n) &\geq \min\{\mu((x * y) * y^n), \mu(y * y^n)\} \\ &= \min\{\mu(x * y^{n+1}), \mu(0)\} = \mu(x * y^{n+1}). \end{aligned} \quad (3.13)$$

By Proposition 3.7, we know that μ is an n -fold fuzzy positive implicative ideal of X . $\square$

Now we give a condition for a fuzzy ideal to be an n -fold fuzzy positive implicative ideal.

THEOREM 3.12. *A fuzzy set μ in X is an n -fold fuzzy positive implicative ideal of X if and only if μ is a fuzzy ideal of X in which the following inequality holds:*

$$(F6) \quad \mu((x * z^n) * (y * z^n)) \geq \mu((x * y) * z^n) \text{ for all } x, y, z \in X.$$

PROOF. Assume that μ is an n -fold fuzzy positive implicative ideal of X . By Proposition 3.4, it follows that μ is a fuzzy ideal of X . Let $a = x * (y * z^n)$ and $b = x * y$. Then

$$\begin{aligned} \mu((a * b) * z^n) &= \mu(((x * (y * z^n)) * (x * y)) * z^n) \\ &\geq \mu((y * (y * z^n)) * z^n) = \mu(0), \end{aligned} \quad (3.14)$$

and so $\mu((a * b) * z^n) = \mu(0)$. Using (F5) we obtain

$$\begin{aligned}\mu((x * z^n) * (y * z^n)) &= \mu((x * (y * z^n)) * z^n) = \mu(a * z^n) \\ &\geq \min\{\mu((a * b) * z^n), \mu(b * z^n)\} \\ &= \min\{\mu(0), \mu(b * z^n)\} \\ &= \mu(b * z^n) = \mu((x * y) * z^n),\end{aligned}\tag{3.15}$$

which is condition (F6). Conversely, let μ be a fuzzy ideal of X satisfying condition (F6). It is sufficient to show that μ satisfies condition (F5). For any $x, y, z \in X$ we have

$$\begin{aligned}\mu(x * z^n) &\geq \min\{\mu((x * z^n) * (y * z^n)), \mu(y * z^n)\} \\ &\geq \min\{\mu((x * y) * z^n), \mu(y * z^n)\},\end{aligned}\tag{3.16}$$

which is precisely (F5). Hence μ is an n -fold fuzzy positive implicative ideal of X . $\square$

THEOREM 3.13. *Let μ be a fuzzy set in X and let n be a positive integer. Then μ is an n -fold fuzzy positive implicative ideal of X if and only if the nonempty level set $U(\mu; t)$ of μ is an n -fold positive implicative ideal of X for every $t \in [0, 1]$.*

PROOF. Assume that μ is an n -fold fuzzy positive implicative ideal of X and $U(\mu; t) \neq \emptyset$ for every $t \in [0, 1]$. Then there exists $x \in U(\mu; t)$. It follows from (F1) that $\mu(0) \geq \mu(x) \geq t$ so that $0 \in U(\mu; t)$. Let $x, y, z \in X$ be such that $(x * y^{n+1}) * z \in U(\mu; t)$ and $z \in U(\mu; t)$. Then $\mu((x * y^{n+1}) * z) \geq t$ and $\mu(z) \geq t$, which imply from (F4) that

$$\mu(x * y^n) \geq \min\{\mu((x * y^{n+1}) * z), \mu(z)\} \geq t,\tag{3.17}$$

so that $x * y^n \in U(\mu; t)$. Therefore $U(\mu; t)$ is an n -fold positive implicative ideal of X . Conversely, suppose that $U(\mu; t) (\neq \emptyset)$ is an n -fold positive implicative ideal of X for every $t \in [0, 1]$. For any $x \in X$, let $\mu(x) = t$. Then $x \in U(\mu; t)$. Since $0 \in U(\mu; t)$, we get $\mu(0) \geq t = \mu(x)$ and so $\mu(0) \geq \mu(x)$ for all $x \in X$. Now assume that there exist $a, b, c \in X$ such that $\mu(a * b^n) < \min\{\mu((a * b^{n+1}) * c), \mu(c)\}$. Selecting $s_0 = (1/2)(\mu(a * b^n) + \min\{\mu((a * b^{n+1}) * c), \mu(c)\})$, then

$$\mu(a * b^n) < s_0 < \min\{\mu((a * b^{n+1}) * c), \mu(c)\}.\tag{3.18}$$

It follows that $(a * b^{n+1}) * c \in U(\mu; s_0)$, $c \in U(\mu; s_0)$, and $a * b^n \notin U(\mu; s_0)$. This is a contradiction. Hence μ is an n -fold fuzzy positive implicative ideal of X . $\square$

THEOREM 3.14. *If μ is an n -fold fuzzy positive implicative ideal of X , then the set*

$$X_\mu := \{x \in X \mid \mu(x) = \mu(0)\}\tag{3.19}$$

is an n -fold positive implicative ideal of X .

PROOF. Let μ be an n -fold fuzzy positive implicative ideal of X . Clearly $0 \in X_\mu$. Let $x, y, z \in X$ be such that $(x * y^{n+1}) * z \in X_\mu$ and $z \in X_\mu$. Then

$$\mu(x * y^n) \geq \min\{\mu((x * y^{n+1}) * z), \mu(z)\} = \mu(0).\tag{3.20}$$

It follows from (F1) that $\mu(x * y^n) = \mu(0)$ so that $x * y^n \in X_\mu$. Hence X_μ is an n -fold positive implicative ideal of X . $\square$

THEOREM 3.15 (extension property for n -fold fuzzy positive implicative ideals). *Let μ and ν be fuzzy ideals of X such that $\mu(0) = \nu(0)$ and $\mu \subseteq \nu$, that is, $\mu(x) \leq \nu(x)$ for all $x \in X$. If μ is an n -fold fuzzy positive implicative ideal of X , then so is ν .*

PROOF. Using Proposition 3.7, it is sufficient to show that ν satisfies the inequality $\nu(x * y^n) \geq \nu(x * y^{n+1})$ for all $x, y \in X$. Let $x, y \in X$. Then

$$\begin{aligned} \nu(0) &= \mu(0) = \mu((x * (x * y^{n+1})) * y^{n+1}) \leq \mu((x * (x * y^{n+1})) * y^n) \\ &= \mu((x * y^n) * (x * y^{n+1})) \leq \nu((x * y^n) * (x * y^{n+1})). \end{aligned} \quad (3.21)$$

Since ν is a fuzzy ideal, it follows from (F1) and (F2) that

$$\begin{aligned} \nu(x * y^n) &\geq \min \{ \nu((x * y^n) * (x * y^{n+1})), \nu(x * y^{n+1}) \} \\ &\geq \min \{ \nu(0), \nu(x * y^{n+1}) \} = \nu(x * y^{n+1}). \end{aligned} \quad (3.22)$$

This completes the proof. $\square$

4. PI^n -Noetherian BCK-algebras

DEFINITION 4.1. A BCK-algebra X is said to satisfy the PI^n -ascending (resp., PI^n -descending) chain condition (briefly, PI^n -ACC (resp., PI^n -DCC)) if for every ascending (resp., descending) sequence $A_1 \subseteq A_2 \subseteq \dots$ (resp., $A_1 \supseteq A_2 \supseteq \dots$) of n -fold positive implicative ideals of X there exists a natural number r such that $A_r = A_k$ for all $r \geq k$. If X satisfies the PI^n -ACC, we say that X is a PI^n -Noetherian BCK-algebra.

THEOREM 4.2. *Let $\{A_k \mid k \in \mathbb{N}\}$ be a family of n -fold positive implicative ideals of X which is nested, that is, $A_1 \supseteq A_2 \supseteq \dots$. Let μ be a fuzzy set in X defined by*

$$\mu(x) = \begin{cases} \frac{k}{k+1} & \text{if } x \in A_k \setminus A_{k+1}, \quad k = 0, 1, 2, \dots, \\ 1 & \text{if } x \in \bigcap_{k=0}^{\infty} A_k, \end{cases} \quad (4.1)$$

for all $x \in X$, where A_0 stands for X . Then μ is an n -fold fuzzy positive implicative ideal of X .

PROOF. Clearly $\mu(0) \geq \mu(x)$ for all $x \in X$. Let $x, y, z \in X$. Suppose that

$$(x * y^{n+1}) * z \in A_k \setminus A_{k+1}, \quad z \in A_r \setminus A_{r+1} \quad (4.2)$$

for $k = 0, 1, 2, \dots; r = 0, 1, 2, \dots$. Without loss of generality, we may assume that $k \leq r$. Then obviously $z \in A_k$. Since A_k is an n -fold positive implicative ideal, it follows that $x * y^n \in A_k$ so that

$$\mu(x * y^n) \geq \frac{k}{k+1} = \min \{ \mu((x * y^{n+1}) * z), \mu(z) \}. \quad (4.3)$$

If $(x * y^{n+1}) * z \in \bigcap_{k=0}^{\infty} A_k$ and $z \in \bigcap_{k=0}^{\infty} A_k$, then $x * y^n \in \bigcap_{k=0}^{\infty} A_k$. Hence

$$\mu(x * y^n) = 1 = \min \{ \mu((x * y^{n+1}) * z), \mu(z) \}. \quad (4.4)$$

If $(x * y^{n+1}) * z \notin \cap_{k=0}^{\infty} A_k$ and $z \in \cap_{k=0}^{\infty} A_k$, then there exists $i \in \mathbb{N}$ such that $(x * y^{n+1}) * z \in A_i \setminus A_{i+1}$. It follows that $x * y^n \in A_i$ so that

$$\mu(x * y^n) \geq \frac{i}{i+1} = \min \{\mu((x * y^{n+1}) * z), \mu(z)\}. \quad (4.5)$$

Finally, assume that $(x * y^{n+1}) * z \in \cap_{k=0}^{\infty} A_k$ and $z \notin \cap_{k=0}^{\infty} A_k$. Then $z \in A_j \setminus A_{j+1}$ for some $j \in \mathbb{N}$. Hence $x * y^n \in A_j$, and thus

$$\mu(x * y^n) \geq \frac{j}{j+1} = \min \{\mu((x * y^{n+1}) * z), \mu(z)\}. \quad (4.6)$$

Consequently, μ is an n -fold fuzzy positive implicative ideal of X . $\square$

Theorem 4.2 tells that if every n -fold fuzzy positive implicative ideal of X has a finite number of values, then X satisfies the PI^n -DCC.

Now we consider the converse of **Theorem 4.2**.

THEOREM 4.3. *Let X be a BCK-algebra satisfying PI^n -DCC and let μ be an n -fold fuzzy positive implicative ideal of X . If a sequence of elements of $\text{Im}(\mu)$ is strictly increasing, then μ has a finite number of values.*

PROOF. Let $\{t_k\}$ be a strictly increasing sequence of elements of $\text{Im}(\mu)$. Hence $0 \leq t_1 < t_2 < \dots \leq 1$. Then $U(\mu; r) := \{x \in X \mid \mu(x) \geq t_r\}$ is an n -fold positive implicative ideal of X for all $r = 2, 3, \dots$. Let $x \in U(\mu; r)$. Then $\mu(x) \geq t_r \geq t_{r-1}$, and so $x \in U(\mu; r-1)$. Hence $U(\mu; r) \subseteq U(\mu; r-1)$. Since $t_{r-1} \in \text{Im}(\mu)$, there exists $x_{r-1} \in X$ such that $\mu(x_{r-1}) = t_{r-1}$. It follows that $x_{r-1} \in U(\mu; r-1)$, but $x_{r-1} \notin U(\mu; r)$. Thus $U(\mu; r) \subsetneq U(\mu; r-1)$, and so we obtain a strictly descending sequence

$$U(\mu; 1) \supsetneq U(\mu; 2) \supsetneq U(\mu; 3) \supsetneq \dots \quad (4.7)$$

of n -fold positive implicative ideals of X which is not terminating. This contradicts the assumption that X satisfies the PI^n -DCC. Consequently, μ has a finite number of values. $\square$

THEOREM 4.4. *The following are equivalent.*

- (i) X is a PI^n -Noetherian BCK-algebra.
- (ii) The set of values of any n -fold fuzzy positive implicative ideal of X is a well-ordered subset of $[0, 1]$.

PROOF. (i) $\Rightarrow$ (ii). Let μ be an n -fold fuzzy positive implicative ideal of X . Assume that the set of values of μ is not a well-ordered subset of $[0, 1]$. Then there exists a strictly decreasing sequence $\{t_k\}$ such that $\mu(x_k) = t_k$. It follows that

$$U(\mu; 1) \subsetneq U(\mu; 2) \subsetneq U(\mu; 3) \subsetneq \dots \quad (4.8)$$

is a strictly ascending chain of n -fold positive implicative ideals of X , where $U(\mu; r) = \{x \in X \mid \mu(x) \geq t_r\}$ for every $r = 1, 2, \dots$. This contradicts the assumption that X is PI^n -Noetherian.

(ii) $\Rightarrow$ (i). Assume that condition (i) is satisfied and X is not PI^n -Noetherian. Then there exists a strictly ascending chain

$$A_1 \subsetneq A_2 \subsetneq A_3 \subsetneq \dots \quad (4.9)$$

of n -fold positive implicative ideals of X . Let $A = \cup_{k \in \mathbb{N}} A_k$. Then A is an n -fold positive implicative ideal of X . Define a fuzzy set ν in X by

$$\nu(x) := \begin{cases} 0 & \text{if } x \notin A_k, \\ \frac{1}{r} & \text{where } r = \min \{k \in \mathbb{N} \mid x \in A_k\}. \end{cases} \quad (4.10)$$

We claim that ν is an n -fold fuzzy positive implicative ideal of X . Since $0 \in A_k$ for all $k = 1, 2, \dots$, we have $\nu(0) = 1 \geq \nu(x)$ for all $x \in X$. Let $x, y, z \in X$. If $(x * y^{n+1}) * z \in A_k \setminus A_{k-1}$ and $z \in A_k \setminus A_{k-1}$ for $k = 2, 3, \dots$, then $x * y^n \in A_k$. It follows that

$$\nu(x * y^n) \geq \frac{1}{k} = \min \{\nu((x * y^{n+1}) * z), \nu(z)\}. \quad (4.11)$$

Suppose that $(x * y^{n+1}) * z \in A_k$ and $z \in A_k \setminus A_r$ for all $r < k$. Since A_k is an n -fold positive implicative ideal, it follows that $x * y^n \in A_k$. Hence

$$\nu(x * y^n) \geq \frac{1}{k} \geq \frac{1}{r+1} \geq \nu(z), \quad \nu(x * y^n) \geq \min \{\nu((x * y^{n+1}) * z), \nu(z)\}. \quad (4.12)$$

Similarly for the case $(x * y^{n+1}) * z \in A_k \setminus A_r$ and $z \in A_k$, we have

$$\nu(x * y^n) \geq \min \{\nu((x * y^{n+1}) * z), \nu(z)\}. \quad (4.13)$$

Thus ν is an n -fold fuzzy positive implicative ideal of X . Since the chain (4.9) is not terminating, ν has a strictly descending sequence of values. This contradicts the assumption that the value set of any n -fold fuzzy positive implicative ideal is well ordered. Therefore X is PI^n -Noetherian. This completes the proof. $\square$

We note that a set is well ordered if and only if it does not contain any infinite descending sequence.

THEOREM 4.5. *Let $S = \{t_k \mid k = 1, 2, \dots\} \cup \{0\}$ where $\{t_k\}$ is a strictly descending sequence in $(0, 1)$. Then a BCK-algebra X is PI^n -Noetherian if and only if for each n -fold fuzzy positive implicative ideal μ of X , $\text{Im}(\mu) \subseteq S$ implies that there exists a natural number k such that $\text{Im}(\mu) \subseteq \{t_1, t_2, \dots, t_k\} \cup \{0\}$.*

PROOF. Assume that X is a PI^n -Noetherian BCK-algebra and let μ be an n -fold fuzzy positive implicative ideal of X . Then by Theorem 4.4 we know that $\text{Im}(\mu)$ is a well-ordered subset of $[0, 1]$ and so the condition is necessary.

Conversely, suppose that the condition is satisfied. Assume that X is not PI^n -Noetherian. Then there exists a strictly ascending chain of n -fold positive implicative ideals

$$A_1 \subsetneq A_2 \subsetneq A_3 \subsetneq \dots \quad (4.14)$$

Define a fuzzy set μ in X by

$$\mu(x) = \begin{cases} t_1 & \text{if } x \in A_1, \\ t_k & \text{if } x \in A_k \setminus A_{k-1}, \quad k = 2, 3, \dots, \\ 0 & \text{if } x \in X \setminus \cup_{k=1}^{\infty} A_k. \end{cases} \quad (4.15)$$

Since $0 \in A_1$, we have $\mu(0) = t_1 \geq \mu(x)$ for all $x \in X$. If either $(x * y^{n+1}) * z$ or z belongs to $X \setminus \cup_{k=1}^{\infty} A_k$, then either $\mu((x * y^{n+1}) * z)$ or $\mu(z)$ is equal to 0 and hence

$$\mu(x * y^n) \geq 0 = \min \{ \mu((x * y^{n+1}) * z), \mu(z) \}. \quad (4.16)$$

If $(x * y^{n+1}) * z \in A_1$ and $z \in A_1$, then $x * y^n \in A_1$ and thus

$$\mu(x * y^n) = t_1 = \min \{ \mu((x * y^{n+1}) * z), \mu(z) \}. \quad (4.17)$$

If $(x * y^{n+1}) * z \in A_k \setminus A_{k-1}$ and $z \in A_k \setminus A_{k-1}$, then $x * y^n \in A_k$. Hence

$$\mu(x * y^n) \geq t_k = \min \{ \mu((x * y^{n+1}) * z), \mu(z) \}. \quad (4.18)$$

Assume that $(x * y^{n+1}) * z \in A_1$ and $z \in A_k \setminus A_{k-1}$ for $k = 2, 3, \dots$. Then $x * y^n \in A_k$ and therefore

$$\mu(x * y^n) \geq t_k = \min \{ \mu((x * y^{n+1}) * z), \mu(z) \}. \quad (4.19)$$

Similarly for $(x * y^{n+1}) * z \in A_k \setminus A_{k-1}$ and $z \in A_1$, $k = 2, 3, \dots$, we obtain

$$\mu(x * y^n) \geq t_k = \min \{ \mu((x * y^{n+1}) * z), \mu(z) \}. \quad (4.20)$$

Consequently, μ is an n -fold fuzzy positive implicative ideal of X . This contradicts our assumption. $\square$

5. Normalizations of n -fold fuzzy positive implicative ideals

DEFINITION 5.1. An n -fold fuzzy positive implicative ideal μ of X is said to be *normal* if there exists $x \in X$ such that $\mu(x) = 1$.

EXAMPLE 5.2. Let $= \{0, a, b\}$ be a BCK-algebra in [Example 3.3](#). Then the fuzzy set μ in X defined by $\mu(0) = 1$, $\mu(a) = 0.8$, and $\mu(b) = 0.5$ is a normal n -fold fuzzy positive implicative ideal of X .

Note that if μ is a normal n -fold fuzzy positive implicative ideal of X , then clearly $\mu(0) = 1$, and hence μ is normal if and only if $\mu(0) = 1$.

PROPOSITION 5.3. Given an n -fold fuzzy positive implicative ideal μ of X let μ^+ be a fuzzy set in X defined by $\mu^+(x) = \mu(x) + 1 - \mu(0)$ for all $x \in X$. Then μ^+ is a normal n -fold fuzzy positive implicative ideal of X which contains μ .

PROOF. We have $\mu^+(0) = \mu(0) + 1 - \mu(0) = 1 \geq \mu(x)$ for all $x \in X$. For any $x, y, z \in X$, we have

$$\begin{aligned} & \min \{ \mu^+((x * y^{n+1}) * z), \mu^+(z) \} \\ &= \min \{ \mu((x * y^{n+1}) * z) + 1 - \mu(0), \mu(z) + 1 - \mu(0) \} \\ &= \min \{ \mu((x * y^{n+1}) * z), \mu(z) \} + 1 - \mu(0) \\ &\leq \mu(x * y^n) + 1 - \mu(0) = \mu^+(x * y^n). \end{aligned} \quad (5.1)$$

Hence μ^+ is a normal n -fold fuzzy positive implicative ideal of X , and obviously $\mu \subseteq \mu^+$. $\square$

Noticing that $\mu \subseteq \mu^+$, we have the following corollary.

COROLLARY 5.4. *If there is $x \in X$ such that $\mu^+(x) = 0$, then $\mu(x) = 0$.*

Using Proposition 3.10, we know that for any n -fold positive implicative ideal A of X , the characteristic function χ_A of A is a normal n -fold fuzzy positive implicative ideal of X . It is clear that μ is a normal n -fold fuzzy positive implicative ideal of X if and only if $\mu^+ = \mu$.

PROPOSITION 5.5. *If μ is an n -fold fuzzy positive implicative ideal of X , then $(\mu^+)^+ = \mu^+$.*

PROOF. The proof is straightforward. $\square$

COROLLARY 5.6. *If μ is a normal n -fold fuzzy positive implicative ideal of X , then $(\mu^+)^+ = \mu$.*

PROPOSITION 5.7. *Let μ and ν be n -fold fuzzy positive implicative ideals of X . If $\mu \subseteq \nu$ and $\mu(0) = \nu(0)$, then $X_\mu \subseteq X_\nu$.*

PROOF. If $x \in X_\mu$, then $\nu(x) \geq \mu(x) = \mu(0) = \nu(0)$ and so $\nu(x) = \nu(0)$, that is, $x \in X_\nu$. Therefore $X_\mu \subseteq X_\nu$. $\square$

PROPOSITION 5.8. *Let μ be an n -fold fuzzy positive implicative ideal of X . If there is an n -fold fuzzy positive implicative ideal ν of X satisfying $\nu^+ \subseteq \mu$, then μ is normal.*

PROOF. Assume that there is an n -fold fuzzy positive implicative ideal ν of X such that $\nu^+ \subseteq \mu$. Then $1 = \nu^+(0) \leq \mu(0)$, and so $\mu(0) = 1$. Hence μ is normal. $\square$

Given an n -fold fuzzy positive implicative ideal, we construct a new normal n -fold fuzzy positive implicative ideal.

THEOREM 5.9. *Let μ be an n -fold fuzzy positive implicative ideal of X and let $f : [0, \mu(0)] \rightarrow [0, 1]$ be an increasing function. Let $\mu_f : X \rightarrow [0, 1]$ be a fuzzy set in X defined by $\mu_f(x) = f(\mu(x))$ for all $x \in X$. Then μ_f is an n -fold fuzzy positive implicative ideal of X . In particular, if $f(\mu(0)) = 1$ then μ_f is normal; and if $f(t) \geq t$ for all $t \in [0, \mu(0)]$, then $\mu \subseteq \mu_f$.*

PROOF. Since $\mu(0) \geq \mu(x)$ for all $x \in X$ and since f is increasing, we have $\mu_f(0) = f(\mu(0)) \geq f(\mu(x)) = \mu_f(x)$ for all $x \in X$. For any $x, y, z \in X$ we get

$$\begin{aligned} \min \{ \mu_f((x * y^{n+1}) * z), \mu_f(z) \} &= \min \{ f(\mu((x * y^{n+1}) * z)), f(\mu(z)) \} \\ &= f(\min \{ \mu((x * y^{n+1}) * z), \mu(z) \}) \leq f(\mu(x * y^n)) = \mu_f(x * y^n). \end{aligned} \quad (5.2)$$

Hence μ_f is an n -fold fuzzy positive implicative ideal of X . If $f(\mu(0)) = 1$, then clearly μ_f is normal. Assume that $f(t) \geq t$ for all $t \in [0, \mu(0)]$. Then $\mu_f(x) = f(\mu(x)) \geq \mu(x)$ for all $x \in X$, which proves $\mu \subseteq \mu_f$. $\square$

Let $\mathcal{N}(X)$ denote the set of all normal n -fold fuzzy positive implicative ideals of X .

THEOREM 5.10. *Let $\mu \in \mathcal{N}(X)$ be nonconstant such that it is a maximal element of the poset $(\mathcal{N}(X), \subseteq)$. Then μ takes only the values 0 and 1.*

PROOF. Since μ is normal, we have $\mu(0) = 1$. Let $x \in X$ be such that $\mu(x) \neq 1$. It is sufficient to show that $\mu(x) = 0$. If not, then there exists $a \in X$ such that $0 < \mu(a) < 1$. Define a fuzzy set v in X by $v(x) = (1/2)\{\mu(x) + \mu(a)\}$ for all $x \in X$. Clearly, v is well defined, and we get

$$v(0) = \frac{1}{2}\{\mu(0) + \mu(a)\} = \frac{1}{2}\{1 + \mu(a)\} \geq \frac{1}{2}\{\mu(x) + \mu(a)\} = v(x) \quad \forall x \in X. \quad (5.3)$$

Let $x, y, z \in X$. Then

$$\begin{aligned} v(x * y^n) &= \frac{1}{2}\{\mu(x * y^n) + \mu(a)\} \geq \frac{1}{2}\{\min\{\mu((x * y^{n+1}) * z), \mu(z)\} + \mu(a)\} \\ &= \min\left\{\frac{1}{2}\{\mu((x * y^{n+1}) * z) + \mu(a)\}, \frac{1}{2}\{\mu(z) + \mu(a)\}\right\} \\ &= \min\{v((x * y^{n+1}) * z), v(z)\}. \end{aligned} \quad (5.4)$$

Hence v is an n -fold fuzzy positive implicative ideal of X . By Proposition 5.3, v^+ is a maximal n -fold fuzzy positive implicative ideal of X , where v^+ is defined by $v^+(x) = v(x) + 1 - v(0)$ for all $x \in X$. Note that

$$\begin{aligned} v^+(a) &= v(a) + 1 - v(0) = \frac{1}{2}\{\mu(a) + \mu(a)\} + 1 - \frac{1}{2}\{\mu(0) + \mu(a)\} \\ &= \frac{1}{2}\{\mu(a) + 1\} > \mu(a) \end{aligned} \quad (5.5)$$

and $v^+(a) < 1 = v^+(0)$. It follows that v^+ is nonconstant, and μ is not a maximal element of $(\mathcal{N}(X), \subseteq)$. This is a contradiction. $\square$

DEFINITION 5.11. An n -fold fuzzy positive implicative ideal μ of X is said to be *fuzzy maximal* if μ is nonconstant and μ^+ is a maximal element of the poset $(\mathcal{N}(X), \subseteq)$.

For any positive implicative ideal I of X let μ_I be a fuzzy set in X defined by

$$\mu_I(x) = \begin{cases} 1 & \text{if } x \in I, \\ 0 & \text{otherwise.} \end{cases} \quad (5.6)$$

THEOREM 5.12. Let μ be an n -fold fuzzy positive implicative ideal of X . If μ is fuzzy maximal, then

- (i) μ is normal,
- (ii) μ takes only the values 0 and 1,
- (ii) $\mu_{x_\mu} = \mu$,
- (iv) X_μ is a maximal n -fold positive implicative ideal of X .

PROOF. Let μ be an n -fold fuzzy positive implicative ideal of X which is fuzzy maximal. Then μ^+ is a nonconstant maximal element of the poset $(\mathcal{N}(X), \subseteq)$. It follows from Theorem 5.10 that μ^+ takes only the values 0 and 1. Note that $\mu^+(x) = 1$ if and only if $\mu(x) = \mu(0)$, and $\mu^+(x) = 0$ if and only if $\mu(x) = \mu(0) - 1$. By Corollary 5.4, we have $\mu(x) = 0$, and so $\mu(0) = 1$. Hence μ is normal and $\mu^+ = \mu$. This proves (i) and (ii).

(iii) Obviously $\mu_{x_\mu} \subset \mu$ and μ_{x_μ} takes only the values 0 and 1. Let $x \in X$. If $\mu(x) = 0$, then $\mu \subseteq \mu_{x_\mu}$. If $\mu(x) = 1$, then $x \in X_\mu$ and so $\mu_{x_\mu}(x) = 1$. This shows that $\mu \subseteq \mu_{x_\mu}$.

(iv) Since μ is nonconstant, X_μ is a proper n -fold positive implicative ideal of X . Let J be an n -fold positive implicative ideal of X containing X_μ . Then $\mu = \mu_{X_\mu} \subseteq \mu_J$. Since μ and μ_J are normal n -fold fuzzy positive implicative ideals of X and since $\mu = \mu^+$ is a maximal element of $\mathcal{N}(X)$, we have that either $\mu = \mu_J$ or $\mu_J = \mathbf{1}$ where $\mathbf{1} : X \rightarrow [0, 1]$ is a fuzzy set defined by $\mathbf{1}(x) = 1$ for all $x \in X$. The later case implies that $J = X$. If $\mu = \mu_J$, then $X_\mu = X_{\mu_J} = J$. This shows that X_μ is a maximal n -fold positive implicative ideal of X . This completes the proof. $\square$

ACKNOWLEDGEMENT. The first author was supported by Korea Research Foundation Grant (KRF-2000-005-D00003).

REFERENCES

- [1] Y. Huang and Z. Chen, *On ideals in BCK-algebras*, Math. Japon. **50** (1999), no. 2, 211–226. [CMP 1 718 851](#). [Zbl 938.06018](#).
- [2] Y. B. Jun, S. M. Hong, J. Meng, and X. L. Xin, *Characterizations of fuzzy positive implicative ideals in BCK-algebras*, Math. Japon. **40** (1994), no. 3, 503–507. [CMP 1 305 546](#). [Zbl 814.06012](#).
- [3] J. Meng, *On ideals in BCK-algebras*, Math. Japon. **40** (1994), no. 1, 143–154. [MR 95e:06049](#). [Zbl 807.06012](#).

YOUNG BAE JUN: DEPARTMENT OF MATHEMATICS EDUCATION, GYEONGSANG NATIONAL UNIVERSITY, JINJU 660-701, KOREA

E-mail address: ybjun@nongae.gsnu.ac.kr

KYUNG HO KIM: DEPARTMENT OF MATHEMATICS, CHUNGJU NATIONAL UNIVERSITY, CHUNGJU 380-702, KOREA

E-mail address: ghkim@gukwon.chungju.ac.kr