ON THE DIOPHANTINE EQUATION $x^3 = dy^2 \pm q^6$

FADWA S. ABU MURIEFAH

(Received 21 December 2000)

ABSTRACT. Let $q > 3$ denote an odd prime and d a positive integer without any prime factor $p \equiv 1 \pmod{3}$. In this paper, we have proved that if $(x, q) = 1$, then $x^3 = dy^2 \pm q^6$ has exactly two solutions provided $q \not\equiv \pm 1 \pmod{24}$.

2000 Mathematics Subject Classification. 11D41.

Cohn [1] and recently Zhang [2, 3] have solved the Diophantine equation

$$x^3 = dy^2 \pm q^6 \tag{1}$$

when $q = 1, 3, 4$, under some conditions on d . In this paper, we consider the general case of (1) where $q \neq 3$ is any odd prime by using arguments similar to those used by Cohn [1].

Let $(x, y) \in \mathbb{Z} \times \mathbb{Z}$ be a solution of (1) with $x, y > 0$, then the solution is trivial if $x = 0, \pm q^2$ or $y = \pm 1$. We need the following lemma.

LEMMA 1. *The equation $p^2 = a^4 - 3b^2$, where p denotes an odd prime and $(p, a) = 1$, may have a solution in positive integers a and b only if $p \equiv \pm 1 \pmod{24}$.*

PROOF. Suppose $3b^2 = a^4 - p^2$. Then clearly a is odd and b is even. Since $a^4 \equiv 3b^2 \pmod{p}$, and $(p, a) = 1$ therefore the Legendre symbol $(3/p) = 1$ and so $p \equiv \pm 1 \pmod{12}$. Now $(a^2 + p, a^2 - p) = 2$ implies that

$$a^2 \pm p = 3.2c^2, \tag{2}$$

$$a^2 \mp p = 2d^2, \tag{3}$$

where $2cd = b$ and $(c, d) = 1$. Whence

$$a^2 = 3c^2 + d^2. \tag{4}$$

Here d is odd, otherwise we get a contradiction modulo 4. Then considering (3) modulo 8, we get $p \equiv \pm 1 \pmod{8}$. This completes the proof. $\square$

Now we consider the upper sign in (1), our main result is laid down in the following.

THEOREM 2. *Let d be a positive integer without prime factor $p \equiv 1 \pmod{3}$ and let $q \neq 3$ be an odd prime. If $q \not\equiv \pm 1 \pmod{24}$ and $(x, q) = 1$, then the Diophantine equation*

$$x^3 = dy^2 + q^6 \tag{5}$$

has exactly two solutions given by

$$\begin{aligned} x_1 &= \frac{3q^4 - 2q^2 - 1}{4}, & y &= ab, & \text{where } a &= \frac{3q^4 + 1}{4}, & db^2 &= \frac{3q^4 - 6q^2 - 1}{4}, \\ x_2 &= \frac{q^4 - 2q^2 - 3}{4}, & y &= 9ab, & \text{where } a &= \frac{q^4 + 3}{4}, & db^2 &= \frac{q^4 - 6q^2 - 3}{4}. \end{aligned} \quad (6)$$

PROOF. If d has a square factor, then it can be absorbed into y^2 , so there is no loss of generality in supposing d a square free integer. Now

$$dy^2 = x^3 - q^6 = (x - q^2)(x^2 + q^2x + q^4). \quad (7)$$

If any prime r divides both d and $(x^2 + q^2x + q^4)$, then by hypothesis $r \equiv 2 \pmod{3}$ or $r = 3$. But $r \mid (x^2 + q^2x + q^4)$ implies that $(2x + q^2)^2 + 3q^4 \equiv 0 \pmod{r}$ so the Legendre symbol $(-3/r) = 1$, which is a contradiction, whence $r = 1$ or 3 . Also since $(x, q) = 1$, therefore $(x - q^2, x^2 + q^2x + q^4) = 1$ or 3 . So for (7) we have only two possibilities: either

$$x^2 + q^2x + q^4 = a^2, \quad x - q^2 = db^2, \quad (8)$$

or

$$x^2 + q^2x + q^4 = 3a^2, \quad x - q^2 = 3db^2, \quad (9)$$

where $(q, a) = 1$ and $(q, b) = 1$. Consider the first possibility when $(2x + q^2)^2 + 3q^4 = (2a)^2$ and $y = ab$. This equation is known to have a finite number of solutions. It can be written as

$$3q^4 = (2a + 2x + q^2)(2a - (2x + q^2)). \quad (10)$$

Then for the nontrivial solution of this equation we have only two cases:

CASE 1.

$$3q^4 = 2a \pm (2x + q^2), \quad 1 = 2a \mp (2x + q^2), \quad (11)$$

by subtracting and adding these two equations we get

$$x = \frac{3q^4 - 2q^2 - 1}{4}, \quad a = \frac{3q^4 + 1}{4}. \quad (12)$$

Here $a > 1$, so $y > 1$, and $x - q^2 = db^2$ implies that

$$db^2 = \frac{3q^4 - 6q^2 - 1}{4}. \quad (13)$$

CASE 2.

$$3 = 2a \pm (2x + q^2), \quad q^4 = 2a \mp (2x + q^2). \quad (14)$$

As in [Case 1](#) we get the nontrivial solution

$$x = \frac{3q^4 - 2q^2 - 1}{4}, \quad a = \frac{3q^4 + 1}{4}, \quad db^2 = \frac{3q^4 - 6q^2 - 1}{4}. \quad (15)$$

Now suppose the second possibility. Obviously a is odd and $x^2 \equiv 3a^2 \pmod{q}$, and since $(q, a) = 1$, so the Legendre symbol $(3/q) = 1$, hence $q \equiv \pm 1 \pmod{12}$. Eliminating x and dividing by 3, we get

$$a^2 = q^4 + 3db^2(q^2 + db^2). \quad (16)$$

Considering (16) modulo 8 we get either $db^2 \equiv -1 \pmod{8}$ or $db^2 \equiv 0 \pmod{8}$.

(1) $db^2 \equiv -1 \pmod{8}$. Then from (16) we get

$$3d^2b^4 = (2a + 2q^2 + 3db^2)(2a - 2q^2 - 3db^2). \quad (17)$$

Let S be a common prime divisor of the two factors in the right-hand side of (17), then S is odd, $S \mid 4a$ and $S \mid 2(2q^2 + 3db^2)$. But S^2 divides the left-hand side implies that $S \mid 3db^2$, so $S \mid q^2$. Here $S = 1$, otherwise $x - q^2 = 3db^2$ implies that $q \mid x$ which is not true. Thus from (17) we get

$$2a \pm (2q^2 + 3db^2) = d_1^2 b_1^4, \quad 2a \mp (2q^2 + 3db^2) = 3d_2^2 b_2^4, \quad (18)$$

where $d = d_1 d_2$ and $b = b_1 b_2$. Whence

$$\pm 2(2q^2 + 3db^2) = d_1^2 b_1^4 - 3d_2^2 b_2^4. \quad (19)$$

Considering this equation modulo 3, we get

$$4q^2 = d_1^2 b_1^4 - 3d_2^2 b_2^4 - 6db^2. \quad (20)$$

Now we prove that $d_1 = 1$. Since d is odd, therefore d_1 must be odd. Let t be any odd prime dividing d_1 then by hypothesis $t \equiv 2 \pmod{3}$ but then from (20) we get

$$4q^2 \equiv -3d_2^2 b_2^4 \pmod{t}, \quad (21)$$

so $(-3/t) = 1$, which is not true. Thus $d_1 = 1$ and (20) becomes

$$q^2 = b_1^4 - 3 \left(\frac{b_1^2 + db_2^4}{2} \right)^2, \quad (22)$$

since $(q, b_1) = 1$, therefore by Lemma 1, $q \equiv \pm 1 \pmod{24}$.

(2) $db^2 \equiv 0 \pmod{8}$. Now we prove that if (16) has a solution, then $q \equiv \pm 1 \pmod{24}$. Since d is a square free, b should be even. Suppose $b = 2m$, then (16) can be written as

$$12d^2 m^4 = (a + q^2 + 6dm^2)(a - q^2 - 6dm^2). \quad (23)$$

As before we can prove that the common divisor of the two factors in the right-hand side of (23) is 2, so

$$a \pm (q^2 + 6dm^2) = 2d_1^2 m_1^4, \quad a \mp (q^2 + 6dm^2) = 6d_2^2 m_2^4, \quad (24)$$

where $d = d_1 d_2$ and $m = m_1 m_2$. It is clear that $(a, q) = 1$ implies that $(m_1, q) = 1$.

Subtracting the two equations in (24) we get

$$\pm(q^2 + 6dm^2) = d_1^2 m_1^4 - 3d_2^2 m_2^4, \quad (25)$$

again considering this equation modulo 3, we get $q^2 = d_1^2 m_1^4 - 3d_2^2 m_2^4 - 6dm^2$. As before d_1 cannot have any odd prime divisor, so $d_1 = 1$ or 2.

If $d_1 = 1$, then

$$q^2 = 4m_1^4 - 3(m_1^2 + dm_2^2). \quad (26)$$

Here m_1 is odd, otherwise we get a contradiction modulo 8. Since $(m_1, q) = 1$, then from (26) we get

$$2m_1^2 \pm q = 3s^2, \quad 2m_1^2 \mp q = n^2, \quad (27)$$

where $sn = m_1^2 + dm_2^2$, so s and n are both odd. Hence $q \equiv \pm 1 \pmod{8}$, combining this result with $q \equiv \pm 1 \pmod{12}$, we get $q \equiv \pm 1 \pmod{24}$.

If $d_1 = 2$, then

$$q^2 = 16b_1^4 - 3(b_1^2 + db_2^2)^2 \quad (28)$$

which is impossible modulo 8. $\square$

Using the same argument as in Theorem 2 we can prove the following theorem.

THEOREM 3. *Let d be a positive integer without prime factor $p \equiv 1 \pmod{3}$ and $q \neq 3$ an odd prime. If $q \not\equiv \pm 1 \pmod{24}$ and $(x, q) = 1$, then the Diophantine equation $x^3 = dy^2 - q^6$ has exactly two solutions given by*

$$\begin{aligned} x_1 &= \frac{3q^4 + 2q^2 - 1}{4}, & y &= ab, & \text{where } a &= \frac{3q^4 + 1}{4}, & db^2 &= \frac{3q^4 + 6q^2 - 1}{4}, \\ x_2 &= \frac{q^4 + 2q^2 - 3}{4}, & y &= 9ab, & \text{where } a &= \frac{q^4 + 3}{4}, & db^2 &= \frac{q^4 + 6q^2 - 3}{4}. \end{aligned} \quad (29)$$

Sometimes, combining our results with Cohn's result [1] we can solve the title equation completely when d has no prime factor $\equiv 1 \pmod{3}$, as we show in the following example.

EXAMPLE 4. Consider the Diophantine equation $x^3 = dy^2 \pm 5^6$ where d has no prime factor $\equiv 1 \pmod{3}$ and $(5, d) = 1$.

Here $q = 5$, when $(x, 5) = 1$, using Theorem 2 for the positive sign this equation has only two solutions given by $x_1 = 456$, $db^2 = 431$, and $x_2 = 143$, $db^2 = 118$. So $d = 431, 118$. Now let $5 \mid x$, then because $(5, d) = 1$, the equation reduces to the form $x^3 = 5dy^2 + 1$, which by [1, Theorem 1] has no solution in positive integers.

So the equation $x^3 = dy^2 + 5^6$ has a solution only if $d = 431, 118$.

For the negative sign this equation has two solutions when $(x, 5) = 1$ given by

$$x_1 = 481, \quad db^2 = 506, \quad x_2 = 168, \quad db^2 = 193, \quad (30)$$

that is, when $d = 506, 193$. If $5 \mid x$, then the equation reduces to the form $x^3 = 5dy^2 - 1$, which by [1, Theorem 2] has no solution in positive integers.

REFERENCES

- [1] J. H. E. Cohn, *The Diophantine equations $x^3 = Ny^2 \pm 1$* , Quart. J. Math. Oxford Ser. (2) **42** (1991), no. 165, 27–30. [MR 92h:11020](#). [Zbl 0722.11019](#).
- [2] F. Z. Li, *On the Diophantine equation $x^3 \pm 4096 = Dy^2$* , Heilongjiang Daxue Ziran Kexue Xuebao **11** (1994), no. 1, 27–31 (Chinese). [MR 95c:11033](#). [Zbl 0836.11012](#).
- [3] ———, *The Diophantine equation $x^3 \pm 729 = Dy^2$* , Dongbei Shida Xuebao (1995), no. 1, 19–20 (Chinese). [MR 97b:11041](#).

FADWA S. ABU MURIEFAH: MATHEMATICS DEPARTMENT, GIRLS COLLEGE OF EDUCATION,
RIYADH, SAUDI ARABIA