RANDOM SUBGRAPHS OF CERTAIN GRAPH POWERS

LANE CLARK

Received 4 March 2002

We determine the limiting probability that a random subgraph of the Cartesian power K_a^n or of $K_{a,a}^n$ is connected.

2000 Mathematics Subject Classification: 05C80.

1. Introduction. A finite, simple, undirected graph G has vertex set $V(G)$ and edge set $E(G)$. The order of G is $|V(G)|$ and the size $e(G)$ of G is $|E(G)|$. For $S \subseteq V(G)$, let $G[S]$ denote the subgraph of G induced by S and $G[S, V(G) - S]$ denote the spanning subgraph of G with edges xy where $x \in S$ and $y \in V(G) - S$. For $U \subseteq V(G)$, let $N_G(U) = \{y \in V(G) : \exists xy \in E(G) \text{ with } x \in U\}$ and $\tilde{N}_G(U) = N_G(U) \cup U$. Of course, $N_G(v) = N_G(\{v\})$ and the degree $d_G(v)$ of v in G is $|N_G(v)|$ for $v \in V(G)$. For $S \subseteq V(G)$, let $b_G(S) = |\{xy \in E(G) : x \in S, y \in V(G) - S\}|$ and $b_G(s) = \min\{b_G(S) : S \subseteq V(G), |S| = s\}$ ($0 \leq s \leq |V(G)|$).

The Cartesian product $G \square H$ of graphs G and H is the graph with vertex set $V(G) \times V(H)$ where vertices (g_1, h_1) and (g_2, h_2) are adjacent if and only if $g_1 = g_2$ and $h_1 h_2 \in E(H)$, or, $h_1 = h_2$ and $g_1 g_2 \in E(G)$. For a graph G , define $G^1 = G$ and $G^n = G^{n-1} \square G$ for $n \geq 2$. We use the following recent isoperimetric result of Tillich [6]. Here K_a denotes the complete graph of order a and $K_{a,a}$ denotes the complete bipartite graph with parts of order a .

LEMMA 1.1 (see [6]). For $G = K_a^n$ with $a \geq 2$ and $n \geq 1$,

$$b_G(s) \geq (a-1)s(n - \log_a s) \quad \text{for } 1 \leq s \leq a^n \quad (1.1)$$

and, for $G = K_{a,a}^n$ with $a \geq 1$ and $n \geq 1$,

$$b_G(s) \geq as(n - \log_{2a} s) \quad \text{for } 1 \leq s \leq (2a)^n. \quad (1.2)$$

Let G be a graph of order n and size N . The probability space $\mathcal{G}(G, p)$ consists of all spanning subgraphs H of G where edges of G are chosen for H independently with probability $0 \leq p = p(n) \leq 1$, so that, $\Pr(H) = p^{e(H)} q^{N-e(H)}$ where $q = q(n) = 1 - p(n)$. (We denote the random graphs in $\mathcal{G}(G, p)$ generally by G_p .)

In this paper, we determine the limiting probability that G_p is connected for $G = K_a^n$ and $K_{a,a}^n$. Specifically, we show that

$$\lim_{n \rightarrow \infty} \Pr(G_p \in \mathcal{G}(K_a^n, p) \text{ is connected}) = e^{-\lambda} \quad (1.3)$$

for fixed $a \geq 2$ where $p = p(n) = 1 - q(n)$ with $q(n) = [(\lambda(n))^{1/n}/a]^{1/(a-1)}$ and $\lim_{n \rightarrow \infty} \lambda(n) = \lambda \in (0, \infty)$. In addition, we show that

$$\lim_{n \rightarrow \infty} \Pr(G_p \in \mathcal{G}(K_{a,a}^n, p) \text{ is connected}) = e^{-\lambda} \quad (1.4)$$

for fixed $a \geq 1$ where $p = p(n) = 1 - q(n)$ with $q(n) = [(\lambda(n))^{1/n}/2a]^{1/a}$ and $\lim_{n \rightarrow \infty} \lambda(n) = \lambda \in (0, \infty)$. Our first result includes those of Burtin [3], Erdős and Spencer [5], and Bollobás [1] as a special case ($a = 2$). Our approach is similar to [1].

The r th factorial moment of a random variable (r.v.) X is denoted by $E_r(X)$. We write $X_n \xrightarrow{d} X$ when the sequence X_n of r.v.s converges in distribution to the r.v. X . Also, we write P_λ for a r.v. having Poisson distribution with mean λ .

Let $[n] = \{1, \dots, n\}$ when n is a positive integer. For a real number x and a positive integer n , let $(x)_0 = 1$ and $(x)_n = (x) \cdots (x - n + 1)$. The cardinality of a set S is denoted by $|S|$. The greatest (least) integer at most (least) the real number x is denoted by $\lfloor x \rfloor$ ($\lceil x \rceil$). We write $\leq$ for an inequality that holds absolutely for the parameters considered and $\leq^*$ for an inequality that holds for the parameters considered and all sufficiently large n . We refer the reader to Bollobás [2] for random graph theory and to Durrett [4] for probability.

2. Results. We use the following result from [1].

LEMMA 2.1 (see [1]). *If G is a simple graph having order $n \geq 1$, maximum degree $\Delta(G) \leq \Delta$, average degree $d = d(G) = 2e(G)/n$, and $\Delta + 1 < u < n - \Delta - 1$, then there exists a u -set $U \subseteq V(G)$ with*

$$|\tilde{N}_G(U)| \geq n \frac{d}{\Delta} \left\{ 1 - \exp\left(-\frac{u(\Delta+1)}{n}\right) \right\}. \quad (2.1)$$

Assume $n \geq 2\Delta + 4$, since the result is vacuously true otherwise, and $\Delta > 0$ (the right-hand side is defined to be 0 for $\Delta = 0$).

We first consider $G = K_a^n$ with $V(G) = [a]^n$ for fixed $a \geq 2$ and for $n \geq 3$. Note that $V(G)$ is totally ordered lexicographically which naturally extends to u -subsets of $V(G)$. In Lemma 2.2 and Theorem 2.5, $\lambda(n) > 0$ for all n .

LEMMA 2.2. *For fixed $a \geq 2$, $q = q(n) = [(\lambda(n))^{1/n}/a]^{1/(a-1)}$ where $\lim_{n \rightarrow \infty} \lambda(n) = \lambda \in (0, \infty)$, and $p = p(n) = 1 - q(n)$, we have*

$$\lim_{n \rightarrow \infty} \Pr(G_p \in \mathcal{G}(K_a^n, p) \text{ has no isolated vertices}) = e^{-\lambda}. \quad (2.2)$$

PROOF. Recall that $G = K_a^n$. Let $X_n(G_p)$ denote the number of isolated vertices in G_p . Fix $r \in \mathbb{P}$ and let $\mathcal{A}_r$ denote the set of r -tuples of V with distinct coordinates; $\mathcal{B}_r = \{(v_1, \dots, v_r) \in \mathcal{A}_r : e(G[\{v_1, \dots, v_r\}]) \neq 0\}$ and $\mathcal{C}_r = \mathcal{A}_r - \mathcal{B}_r = \{(v_1, \dots, v_r) \in \mathcal{A}_r : e(G[\{v_1, \dots, v_r\}]) = 0\}$. Then $|\mathcal{B}_r| \leq (a^n)_{r-1} r a n \leq a^{n(r-1)} r a n$ and $|\mathcal{C}_r| = (a^n)_r - |\mathcal{B}_r| \geq a^{nr} e^{-r^2/a^n} - a^{n(r-1)} r a n$. Observe that the number of edges in G incident with $\{v_1, \dots, v_r\}$ is at least $(a-1)r(n-r)$ for all $(v_1, \dots, v_r) \in \mathcal{A}_r$.

First,

$$\begin{aligned} 0 &\leq \sum_{(v_1, \dots, v_r) \in \mathcal{B}_r} \Pr(d_{G_p}(v_1) = \dots = d_{G_p}(v_r) = 0) \leq |\mathcal{B}_r| q^{(a-1)r(n-r)} \\ &\leq a^{n(r-1)} r a n \frac{(\lambda(n))^{r-r^2/n}}{a^{r(n-r)}} = \frac{(\lambda(n))^{r-r^2/n} r a n}{a^{n-r^2}}. \end{aligned} \quad (2.3)$$

Next,

$$\begin{aligned} \sum_{(v_1, \dots, v_r) \in \mathcal{C}_r} \Pr(d_{G_p}(v_1) = \dots = d_{G_p}(v_r) = 0) &= |\mathcal{C}_r| q^{(a-1)nr} \\ &\stackrel{*}{\geq} [a^{nr} e^{-r^2/a^n} - a^{n(r-1)} r a n] \frac{\lambda^r(n)}{a^{nr}} \\ &= \lambda^r(n) e^{-r^2/a^n} - \frac{\lambda^r(n) r a n}{a^n} \end{aligned} \quad (2.4)$$

while,

$$\sum_{(v_1, \dots, v_r) \in \mathcal{C}_r} \Pr(d_{G_p}(v_1) = \dots = d_{G_p}(v_r) = 0) \leq a^{nr} q^{(a-1)nr} = \lambda^r(n). \quad (2.5)$$

Hence,

$$\lambda^r(n) e^{-r^2/a^n} - \frac{\lambda^r(n) r a n}{a^n} \stackrel{*}{\leq} E_r(X_n) \leq \lambda^r(n) + \frac{(\lambda(n))^{r-r^2/n} r a n}{a^{n-r^2}} \quad (2.6)$$

so that,

$$\lim_{n \rightarrow \infty} E_r(X_n) = \lambda^r \quad (2.7)$$

and $X_n \xrightarrow{d} P_\lambda$ (see [4]). $\square$

LEMMA 2.3. For fixed $a \geq 2$, $q = q(n) = [(\ln n)^{1/n}/a]^{1/(a-1)}$, and $p = p(n) = 1 - q(n)$, we have

$$\Pr(G_p \in \mathcal{G}(K_a^n, p) \text{ has a component of order } s \text{ with } 2 \leq s \leq a^n/2) = o(1) \quad \text{as } n \rightarrow \infty. \quad (2.8)$$

PROOF. Recall that $G = K_a^n$. Let $\mathcal{A}_s = \{S \subseteq V(G) : |S| = s\}$ ($1 \leq s \leq a^n$). We consider four cases.

CASE 1 ($2 \leq s \leq s_1 = \lfloor a^{n/2}/n \rfloor$). We have

$$\begin{aligned} |\{S \in \mathcal{A}_s : G[S] \text{ is connected}\}| &\leq a^n \cdot (a-1)n \cdot 2(a-1)n \cdots (s-1)(a-1)n \\ &\leq a^{n+s} n^{s-1} s^s, \end{aligned} \quad (2.9)$$

so that (Lemma 1.1)

$$\begin{aligned} \sum_{S \in \mathcal{A}_s} \Pr(G_p[S] \text{ is a component}) &\leq a^{n+s} n^{s-1} s^s q^{b_G(s)} \\ &= a^{n+s} n^{s-1} s^s \left[\frac{(\ln n)^{1/n}}{a} \right]^{s(n - \log_a s)} \\ &= \frac{1}{n} \left[\frac{a n s^2 \ln n}{a^{n(1-1/s)}} \right]^s. \end{aligned} \quad (2.10)$$

By examining the derivative $f(s) \ln(ce^2 s^2/a^n)$ with respect to s of $f(s) = c^s s^{2s}/a^{n(s-1)}$ with $c = an \ln n$, we see that $f(s)$ is decreasing for $s \in [2, a^{n/2}/ec^{1/2}]$. Here $f(s) \leq^* f(2) = 16a^2 n^2 \ln^2 n/a^n$. Hence,

$$\sum_{s=2}^{s_1} \sum_{S \in \mathcal{A}_s} \Pr(G_p[S] \text{ is a component}) \leq^* \sum_{s=2}^{s_1} \frac{16a^2 n \ln^2 n}{a^n} = o(1) \quad \text{as } n \rightarrow \infty. \quad (2.11)$$

CASE 2 ($s_1 + 1 \leq s \leq s_3 = \lfloor a^n/2 \rfloor$). Let $\mathcal{B}_s = \{S \in \mathcal{A}_s : b_G(S) \geq (a-1)s(n - \log_a(s/n))\}$ and $\mathcal{C}_s = \mathcal{A}_s - \mathcal{B}_s = \{S \in \mathcal{A}_s : b_G(S) < (a-1)s(n - \log_a(s/n))\}$.

First,

$$\begin{aligned} \sum_{S \in \mathcal{B}_s} \Pr(G_p[S] \text{ is a component}) &\leq \binom{a^n}{s} q^{(a-1)s(n - \log_a(s/n))} \\ &\leq \left(\frac{ea^n}{s}\right)^s \left[\frac{(\ln n)^{1/n}}{a}\right]^{s(n - \log_a(s/n))} \\ &= \left[\frac{e(\ln n)^{1 - (1/n)\log_a(s/n)}}{n}\right]^s \\ &\leq^* \left(\frac{e \ln n}{n}\right)^s. \end{aligned} \quad (2.12)$$

Hence,

$$\sum_{s=s_1+1}^{s_3} \sum_{S \in \mathcal{B}_s} \Pr(G_p[S] \text{ is a component}) \leq^* \sum_{s=s_1+1}^{s_3} \left(\frac{e \ln n}{n}\right)^s = o(1) \quad \text{as } n \rightarrow \infty. \quad (2.13)$$

Next, for $S \in \mathcal{C}_s$, let $H = G[S]$. Then

$$(a-1)sn = \sum_{v \in S} d_G(v) = 2e(H) + b_G(S) < 2e(H) + (a-1)s\left(n - \log_a \frac{s}{n}\right), \quad (2.14)$$

so that

$$2e(H) \geq (a-1)s \log_a \frac{s}{n} \quad (2.15)$$

and the average degree d in H satisfies

$$d > (a-1) \log_a \frac{s}{n}. \quad (2.16)$$

CASE 3 ($s_1 + 1 \leq s \leq s_2 = \lfloor a^n/\ln^2 n \rfloor$). Let $u = \lfloor s/n \rfloor$, so that $(a-1)n + 1 \leq^* u \leq^* s - (a-1)n - 1$, and by [Lemma 2.1](#), for sufficiently large n , there exists $U \subseteq S$, $|U| = u$, and

$$\begin{aligned} |\tilde{N}_H(U)| &\geq^* \frac{s}{n} \log_a \frac{s}{n} \left\{ 1 - \exp\left(-\frac{u[(a-1)n+1]}{s}\right) \right\} \\ &\geq \frac{\delta s}{n} \log_a \frac{s}{n} \quad \text{with } \delta = 1 - e^{-1} = 0.631 \dots \end{aligned} \quad (2.17)$$

Let $t = \lceil (\delta s/n) \log_a(s/n) \rceil$, so that $u \leq^* t \leq^* s$, and let $w = s - t = s(1-x) - \tau$ with $x = (\delta/n) \log_a(s/n)$ and $0 \leq \tau < 1$. Observe that $\delta/4 \leq^* x \leq^* \delta$ here. For sufficiently

large n , take the smallest such u -set $U = \{d_1, \dots, d_u\}$ in $S (\subseteq V(G))$ which is totally ordered; take the (uniquely determined) first $t - u$ vertices of $(N_G(d_1) \cap (S - U)) \cup \dots \cup (N_G(d_u) \cap (S - U)) (\subseteq V(G))$; and add the remaining w vertices W of S . Then

$$S \mapsto (\{d_1, \dots, d_u\}; N_G(d_1) \cap (S - U), \dots, N_G(d_u) \cap (S - U); W) \quad (2.18)$$

is an injection. Hence,

$$\begin{aligned} |\mathcal{C}_S| &\leq \binom{a^n}{u} 2^{(a-1)nu} \binom{a^n}{w} \\ &\leq \left(\frac{ea^n}{u}\right)^u 2^{(a-1)nu} \left(\frac{ea^n}{w}\right)^w \\ &\leq \left(\frac{ena^n}{s}\right)^{s/n} 2^{(a-1)s} \left(\frac{ea^n}{s(1-x)}\right)^{s(1-x)}. \end{aligned} \quad (2.19)$$

Then (where $x - 1/n \stackrel{*}{>} 0$, [Lemma 1.1](#))

$$\begin{aligned} \sum_{S \in \mathcal{C}_S} \Pr(G_p[S] \text{ is a component}) &\leq |\mathcal{C}_S| q^{b_G(S)} \\ &\stackrel{*}{\leq} \left(\frac{ena^n}{s}\right)^{s/n} 2^{(a-1)s} \left(\frac{ea^n}{s(1-x)}\right)^{s(1-x)} \left[\frac{(\ln n)^{1/n}}{a}\right]^{s(n - \log_a s)} \\ &= \left[(en)^{1/n} 2^{a-1} \left(\frac{e}{1-x}\right)^{1-x} \left(\frac{s}{a^n}\right)^{x-1/n} (\ln n)^{1-(1/n)\log_a s}\right]^s \\ &\stackrel{*}{\leq} \left[(en)^{1/n} 2^{a-1} \left(\frac{e}{1-x}\right)^{1-x} (\ln n)^{1+(2/n)-2x-(1/n)\log_a s}\right]^s. \end{aligned} \quad (2.20)$$

Here

$$2x + \frac{1}{n} \log_a s - 1 - \frac{2}{n} \geq \delta - \frac{1}{2} - \frac{4}{n} \log_a n - \frac{2}{n} \stackrel{*}{\geq} \frac{1}{10}, \quad (2.21)$$

so that

$$\sum_{S \in \mathcal{C}_S} \Pr(G_p[S] \text{ is a component}) \stackrel{*}{\leq} \left[(en)^{1/n} 2^{a-1} \left(\frac{e}{1-x}\right)^{1-x} (\ln n)^{-0.1}\right]^s. \quad (2.22)$$

Hence,

$$\begin{aligned} \sum_{s=s_1+1}^{s_2} \sum_{S \in \mathcal{C}_S} \Pr(G_p[S] \text{ is a component}) &\stackrel{*}{\leq} \sum_{s=s_1+1}^{s_2} \left[(en)^{1/n} 2^{a-1} \left(\frac{e}{1-x}\right)^{1-x} (\ln n)^{-0.1}\right]^s \\ &= o(1) \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (2.23)$$

CASE 4 ($s_2 + 1 \leq s \leq s_3$). For $S \in \mathcal{C}_s$ and $H = G[S]$, let $T = \{v \in S : d_H(v) \geq (a - 1)n - \log_a^2 n\}$, $t = |T|$ and $H_1 = H[T] = G[T]$. Then

$$\begin{aligned} 2e(H_1) &= 2e(H) - 2e(H[S - T, T]) - 2e(H[S - T]) \\ &> (a - 1)s \log_a \frac{s}{n} - 2(a - 1)n(s - t) \\ &= (a - 1)s \left[\log_a \frac{s}{n} - \frac{2n}{s}(s - t) \right]. \end{aligned} \quad (2.24)$$

Here

$$\log_a \frac{s}{n} \stackrel{*}{\geq} n - 2 \log_a n, \quad (2.25)$$

so that

$$\begin{aligned} s(a - 1)n - (s - t) \log_a^2 n &\geq \sum_{v \in T} d_H(v) + \sum_{v \in S - T} d_H(v) > (a - 1)s \log_a \frac{s}{n} \\ &\stackrel{*}{\geq} (a - 1)s(n - 2 \log_a n), \end{aligned} \quad (2.26)$$

hence,

$$t \stackrel{*}{\geq} s \left(1 - \frac{2(a - 1)}{\log_a n} \right). \quad (2.27)$$

We take the first t vertices of T for H_1 where $t = s(1 - \epsilon)$ with $s\epsilon = \lfloor 2(a - 1)s / \log_a n \rfloor$ so that $0 < (a - 1) / \log_a n \stackrel{*}{<} \epsilon \stackrel{*}{<} 2(a - 1) / \log_a n \stackrel{*}{<} 1/5$. Then

$$2e(H_1) \stackrel{*}{>} (a - 1)s[(1 - 2\epsilon)n - 2 \log_a n] \quad (2.28)$$

and the average degree d_1 in H_1 satisfies

$$d_1 \stackrel{*}{>} (a - 1) \left[n - \frac{\epsilon}{1 - \epsilon} n - \frac{2}{1 - \epsilon} \log_a n \right] \stackrel{*}{\geq} (a - 1)(1 - 3\epsilon)n. \quad (2.29)$$

Let $u = \lceil a^n / \ln^6 n \rceil$, so that $(a - 1)n + 1 \stackrel{*}{<} u \stackrel{*}{<} t - (a - 1)n - 1$, and by [Lemma 2.1](#), for all sufficiently large n , there exists $U \subseteq T$, $|U| = u$, and

$$\begin{aligned} |\tilde{N}_H(U)| &\geq |\tilde{N}_{H_1}(U)| \stackrel{*}{\geq} s(1 - \epsilon)(1 - 3\epsilon) \left\{ 1 - \exp \left(- \frac{u[(a - 1)n + 1]}{t} \right) \right\} \\ &\stackrel{*}{\geq} s(1 - \epsilon)^2(1 - 3\epsilon) \geq s(1 - 4\epsilon). \end{aligned} \quad (2.30)$$

Let $t = s - \lfloor 4\epsilon s \rfloor$, so that $u \stackrel{*}{<} t \stackrel{*}{<} s$, and $w = \lfloor 4\epsilon s \rfloor$. For sufficiently large n , take the smallest such u -set $U = \{d_1, \dots, d_u\}$ in $S (\subseteq V(G))$; take the (uniquely determined) first $t - u$ vertices of $(N_G(d_1) \cap (S - U)) \cup \dots \cup (N_G(d_u) \cap (S - U)) (\subseteq V(G))$; and add the remaining w vertices W of S . Then

$$S \mapsto (\{d_1, \dots, d_u\}; N_G(d_1) - S, \dots, N_G(d_u) - S; W) \quad (2.31)$$

is an injection with $|N_G(d_i) - S| \leq \lfloor \log_a^2 n \rfloor$ ($1 \leq i \leq u$). Hence, with $y = \lfloor \log_a^2 n \rfloor$,

$$\begin{aligned} |\mathcal{C}_S| &\leq \binom{a^n}{u} \sum_{(k_1, \dots, k_u) \in \{0, \dots, y\}^u} \prod_{i=1}^u \binom{(a-1)n}{k_i} \binom{a^n}{w} \\ &\leq \binom{a^n}{u} (y+1)^u \binom{(a-1)n}{y+1}^u \binom{a^n}{w}, \end{aligned} \quad (2.32)$$

since

$$\binom{(a-1)n}{k} \leq \binom{(a-1)n}{y+1}, \quad \forall k \in \{0, \dots, y\}. \quad (2.33)$$

Then

$$\begin{aligned} |\mathcal{C}_S| &\leq \left(\frac{ea^n}{u} \right)^u (y+1)^u \left(\frac{ean}{y+1} \right)^{u(y+1)} \left(\frac{ea^n}{w} \right)^w \\ &\leq (e^2 an \ln^6 n)^u \left(\frac{ean}{\log_a^2 n} \right)^{uy} \left(\frac{ea^n}{4\epsilon s} \right)^{4\epsilon s}. \end{aligned} \quad (2.34)$$

Hence, (Lemma 1.1)

$$\begin{aligned} \sum_{S \in \mathcal{C}_S} \Pr(G_p[S] \text{ is a component}) &\leq |\mathcal{C}_S| q^{b_G(S)} \\ &\leq (e^2 an \ln^6 n)^u \left(\frac{ean}{\log_a^2 n} \right)^{uy} \left(\frac{ea^n}{4\epsilon s} \right)^{4\epsilon s} \left[\frac{(\ln n)^{1/n}}{a} \right]^{s(n - \log_a s)} \\ &= \left[(e^2 an \ln^6 n)^{u/s} (ean \ln^2 a)^{uy/s} \left(\frac{e}{4\epsilon} \right)^{4\epsilon} \left(\frac{s}{a^n} \right)^{1-4\epsilon} (\ln n)^{1-(1/n)\log_a s - 2uy/s} \right]^s. \end{aligned} \quad (2.35)$$

Here

$$\begin{aligned} 1 \leq ean \ln^2 a \leq e^2 an \ln^6 n, \quad 0 < \frac{u}{s} \leq \frac{uy}{s} \leq \frac{5}{\ln^2 n}, \\ 1 - \frac{1}{n} \log_a s - \frac{2uy}{s} \leq \frac{2}{n} \log_a \ln n - \frac{4}{\ln^4 n} \leq 0, \end{aligned} \quad (2.36)$$

so that

$$\begin{aligned} \sum_{S \in \mathcal{C}_S} \Pr(G_p[S] \text{ is a component}) &\leq \left[(e^3 a^2 n^2 \ln^2 a \ln^6 n)^{5/\ln^2 n} \left(\frac{e}{4\epsilon} \right)^{4\epsilon} 2^{4\epsilon-1} \right]^s \\ &\leq \left(\frac{2}{3} \right)^s, \end{aligned} \quad (2.37)$$

since $(e^3 a^2 n^2 \ln^2 a \ln^6 n)^{5/\ln^2 n} \rightarrow 1$, $(e/4\epsilon)^{4\epsilon} \rightarrow 1$ and $\epsilon \rightarrow 0$ as $n \rightarrow \infty$. Hence,

$$\sum_{s=s_2+1}^{s_3} \sum_{S \in \mathcal{C}_S} \Pr(G_p[S] \text{ is a component}) \leq \sum_{s=s_2+1}^{s_3} \left(\frac{2}{3} \right)^s = o(1) \quad \text{as } n \rightarrow \infty. \quad (2.38)$$

□

REMARK 2.4. For all $a \geq 2$ and $n \geq 2$, $b_G(s) \geq 2$ when $2 \leq s \leq a^n/2$. Hence, $0 < \tilde{q}(n) \leq q(n)$ implies $(\tilde{q}(n))^{b_G(s)} \leq (q(n))^{b_G(s)}$ when $2 \leq s \leq a^n/2$. Then (2.10), (2.12), (2.20), and (2.35) hold for $G_{\tilde{p}(n)}$ where $\tilde{p}(n) = 1 - \tilde{q}(n)$ (the exponent in (2.12) is larger than $b_G(s)$). Hence, Lemma 2.3 holds for $G_{\tilde{p}(n)}$ as well. The inequalities in the proof of Lemma 2.3 hold for all sufficiently large n which can be determined from nineteen appropriate inequalities there.

THEOREM 2.5. For fixed $a \geq 2$, $q = q(n) = [(\lambda(n))^{1/n}/a]^{1/(a-1)}$ where $\lim_{n \rightarrow \infty} \lambda(n) = \lambda \in (0, \infty)$, and $p = p(n) = 1 - q(n)$, we have

$$\lim_{n \rightarrow \infty} \Pr(G_p \in \mathcal{G}(K_a^n, p) \text{ is connected}) = e^{-\lambda}. \quad (2.39)$$

PROOF. We have

$$\begin{aligned} 0 &\leq \Pr(G_p \text{ is disconnected}) - \Pr(G_p \text{ has isolated vertices}) \\ &\leq \Pr(G_p \text{ has a component of order } s \text{ with } 2 \leq s \leq a^{n/2}) = o(1) \quad \text{as } n \rightarrow \infty, \end{aligned} \quad (2.40)$$

by Remark 2.4. Hence, Lemma 2.2 gives

$$\lim_{n \rightarrow \infty} \Pr(G_p \text{ is disconnected}) = \lim_{n \rightarrow \infty} \Pr(G_p \text{ has isolated vertices}) = 1 - e^{-\lambda}. \quad (2.41)$$

□

We state the result for $G = K_{a,a}^n$ since its proof is similar to the proof of Theorem 2.5.

THEOREM 2.6. For fixed $a \geq 1$, $q = q(n) = [(\lambda(n))^{1/n}/2a]^{1/a}$ where $\lim_{n \rightarrow \infty} \lambda(n) = \lambda \in (0, \infty)$, and $p = p(n) = 1 - q(n)$, we have

$$\lim_{n \rightarrow \infty} \Pr(G_p \in \mathcal{G}(K_{a,a}^n) \text{ is connected}) = e^{-\lambda}. \quad (2.42)$$

REFERENCES

- [1] B. Bollobás, *The evolution of the cube*, Combinatorial Mathematics (Marseille-Luminy, 1981) (C. Berge, D. Bresson, P. Camion, J. F. Maurras, and F. Sterboul, eds.), North-Holland Mathematics Studies, vol. 75, North-Holland, Amsterdam, 1983, pp. 91–97.
- [2] ———, *Random Graphs*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 73, Cambridge University Press, Cambridge, 2001.
- [3] J. D. Burton, *The probability of connectedness of a random subgraph of an n -dimensional cube*, Problemy Peredachi Informatsii 13 (1977), no. 2, 90–95.
- [4] R. Durrett, *Probability. Theory and Examples*, The Wadsworth and Brooks/Cole Statistics/Probability Series, Wadsworth and Brooks/Cole Advanced Books and Software, Pacific Grove, California, 1991.
- [5] P. Erdős and J. Spencer, *Evolution of the n -cube*, Comput. Math. Appl. 5 (1979), no. 1, 33–39.
- [6] J.-P. Tillich, *Edge isoperimetric inequalities for product graphs*, Discrete Math. 213 (2000), no. 1–3, 291–320.

LANE CLARK: DEPARTMENT OF MATHEMATICS, SOUTHERN ILLINOIS UNIVERSITY CARBONDALE, CARBONDALE, IL 62901-4408, USA

E-mail address: lcclark@math.siu.edu