SUBMODULES OF SECONDARY MODULES

SHAHABADDIN EBRAHIMI ATANI

Received 31 July 2001 and in revised form 25 January 2002

Let R be a commutative ring with nonzero identity. Our objective is to investigate representable modules and to examine in particular when submodules of such modules are representable. Moreover, we establish a connection between the secondary modules and the pure-injective, the Σ -pure-injective, and the prime modules.

2000 Mathematics Subject Classification: 13F05.

1. Introduction. In this paper, all rings are commutative rings with identity and all modules are unital. The notion of associated prime ideals and the related one of primary decomposition are classical. In a dual way, we define the attached prime ideals and the secondary representation. This theory is developed in the appendix to Section 6 in Matsumura [6] and in Macdonald [5]. Now we define the concepts that we will need.

Let R be a ring and let $0 \neq M$ be an R -module. Then M is called a secondary module (second module) provided that for every element r of R the homothety $M \xrightarrow{r} M$ is either surjective or nilpotent (either surjective or zero). This implies that $\text{nilrad}(M) = P$ ($\text{Ann}(M) = P'$) is a prime ideal of R , and M is said to be P -secondary (P' -second), so every second module is secondary (the concept of second module is introduced by Yassemi [14]). A secondary representation for an R -module M is an expression for M as a finite sum of secondary modules (see [5]). If such a representation exists, we will say that M is representable.

If R is a ring and N is a submodule of an R -module M , the ideal $\{r \in R : rM \subseteq N\}$ will be denoted by $(N : M)$. Then $(0 : M)$ is the annihilator of M , $\text{Ann}(M)$. A proper submodule N of a module M over a ring R is said to be prime submodule (primary submodule) if for each $r \in R$ the homothety $M/N \xrightarrow{r} M/N$ is either injective or zero (either injective or nilpotent), so $(0 : M/N) = P$ ($\text{nilrad}(M/N) = P'$) is a prime ideal of R , and N is said to be P -prime submodule (P' -primary submodule). So N is prime in M if and only if whenever $rm \in N$, for some $r \in R$, $m \in M$, then $m \in N$ or $rM \subseteq N$. We say that M is a prime module (primary module) if zero submodule of M is prime (primary) submodule of M , so N is a prime submodule of M if and only if M/N is a prime module. Moreover, every prime module is primary.

Let R be a ring, and let N be an R -submodule of M . Then N is pure in M if for any finite system of equations over N which is solvable in M , the system is also solvable in N . A module is said to be absolutely pure if every embedding of it into any other modules is pure embedding. A submodule N of an R -module M is called relatively divisible (or an RD-submodule) if $rN = N \cap rM$ for all $r \in R$. Every RD-submodule of a P -secondary module over a commutative ring R is P -secondary (see [2, Lemma 2.1]).

A module M is pure-injective if and only if any system of equations in M which is finitely solvable in M , has a global solution in M [7, Theorem 2.8]. The module N is a pure-essential extension of M if M is pure in N and for all nonzero submodules L of N , if $M \cap L = 0$, then $(M \oplus L)/L$ is not pure in N/L . A pure-injective hull $H(M)$ of a module M is a pure essential extension of M which is pure-injective. Every module M has a pure-injective hull which is unique to isomorphism over M [12].

Given an R -module M and index set I , the direct sum of the family $\{M_i : i \in I\}$ where $M_i = M$ for each $i \in I$ will be denoted by $M^{(I)}$. Given a module property $\mathcal{P}$, we will say that a module M is $\sum\text{-}\mathcal{P}$ if $M^{(I)}$ satisfies $\mathcal{P}$ for every index set I .

Let R be a commutative ring. An element $a \in R$ is said to be regular if there exists $b \in R$ such that $a = a^2b$, and R is said to be regular if each of its elements is regular. An important property of regular rings is that every module is absolutely pure (see [13, Theorem 37.6]).

Let R be a ring and M an R -module. A prime ideal P of R is called an associated prime ideal of M if P is the annihilator $\text{Ann}(x)$ of some $x \in M$. The set of associated primes of M is written $\text{Ass}(M)$. For undefined terms, we refer to [6, 7].

2. Secondary submodules. In general, a nonzero submodule of a representable (even secondary) R -module is not representable (secondary), but we have the following results.

LEMMA 2.1. *Let R be a commutative ring and let $0 \neq N$ be an RD-submodule of R -module M . Then M is P -secondary if and only if N and M/N are P -secondary.*

PROOF. If M is P -secondary, then N and M/N are P -secondary by [2, Lemma 2.1] and [5, Theorem 2.4], respectively. Conversely, suppose that $r \in R$. If $r \in P$, then $r^n(M/N) = 0$ and $r^nN = 0$ for some n , hence $r^nM \subseteq N$ and $0 = r^nN = r^nM \cap N = r^nM$. If $r \notin P$, then $rM + N = M$, $rN = N$, and $N = rN = rM \cap N$, so we have $rM = M$, as required. $\square$

COROLLARY 2.2. *Let R be a commutative regular ring, and let $0 \neq N$ be a submodule of R -module M . Then M is P -secondary if and only if N and M/N are P -secondary.*

PROOF. This follows from Lemma 2.1. $\square$

THEOREM 2.3. *Let R be a commutative regular ring. Then every nonzero submodule of a representable R -module is representable.*

PROOF. Let M be a representable R -module and let $M = \sum_{i=1}^n M_i$ be a minimal secondary representation with $\text{nilrad}(M_i) = P_i$. There is an element $r_1 \in P_1$ such that $r_1 \notin \cup_{i=2}^n P_i$. Otherwise $P_1 \subseteq \cup_{i=2}^n P_i$, so by [10, Theorem 3.61], $P_1 \subseteq P_j$ for some j , and hence $P_1 = P_j$, a contradiction. Thus there exists a positive integer m_1 such that $r_1^{m_1} \in \text{Ann}(M_1)$ and the module $r_1^{m_1}M = \sum_{i=2}^n r_1^{m_1}M_i$ is representable. By using this process for the ideals $P_2, \dots, P_{n-1}$, there are integers $m_2, \dots, m_{n-1}$ and elements $r_2 \in P_2, \dots, r_{n-1} \in P_{n-1}$ such that $s_nM = M_n$, where $0 \neq s_n = r_1^{m_1}r_2^{m_2} \dots r_{n-1}^{m_{n-1}}$, $s_n \in \cap_{i=1}^{n-1} P_i$ and $s_n \notin P_n$. Therefore by a similar argument, there are elements $s_1, \dots, s_{n-1}$

such that $M = \sum_{i=1}^n s_i M$, where for each i , where $i = 1, \dots, n$, $s_i \notin P_i$, $s_i M = M_i$, and $s_i \in \cap_{i=1, i \neq j}^n \text{Ann}(M_j)$.

Let N be a nonzero submodule of M and $0 \neq a \in N$. Then $a = s_1 b_1 + \dots + s_n b_n$ for some $b_i \in M$, $i = 1, \dots, n$. By assumption, there exists $t_1, \dots, t_n \in R$ such that for each i , $s_i = s_i^2 t_i$. As $0 \neq a$, $s_i b_i \neq 0$ for some i and $s_i t_i a = s_i^2 t_i b_i = s_i b_i$, so $s_i N \neq 0$. We can assume that $s_{i_1} N \neq 0, \dots, s_{i_k} N \neq 0$, where $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$. By a similar argument as above, if $a \in N$, then $a = \sum_{j=1}^k s_{i_j} t_{i_j} a \in \sum_{j=1}^k s_{i_j} N$, and hence $N = \sum_{j=1}^k s_{i_j} N$. Since for each j , where $j = 1, \dots, k$, $s_{i_j} N$ is pure in the P_{i_j} -secondary module M_{i_j} , it is P_{i_j} -secondary by [2, Lemma 2.1], as required. $\square$

THEOREM 2.4. *Let R be a commutative ring and let N be a prime submodule of secondary R -module of M . Then N is $(N : M)$ -secondary.*

PROOF. Suppose that M is a P -secondary module over R . Let $r \in R$. If $r \in P$, then $r^n N \subseteq r^n M = 0$ for some n . If $r \notin P$, then $rM = M$. Suppose that $n \in N$, so there is an element $m \in M$ such that $n = rm$. As N is a prime submodule of M and $N \neq rM = M$, $m \in N$, so $rN = N$, hence N is P -secondary.

By [4, Lemma 1], the ideal $P' = (N : M) = \{r \in R : rM \subseteq N\}$ is prime. Clearly, $P' \subseteq P$. Let $s \in P$. Then $s^n N = s^n M = 0$ for some n . There is an element $m \in M$ such that $m \notin N$ and $s^n m = 0 \in N$, so $s^n \in P'$, hence $s \in P'$. Thus $P = P'$, as required. $\square$

PROPOSITION 2.5. *Let R be a commutative ring and let N be a prime submodule of P -second R -module of M . Then N is an RD-submodule of M .*

PROOF. Let $r \in R$. If $r \in P$, then $rN \subseteq rM = 0$, so $rN = N \cap rM = 0$. If $r \notin P$, then $rM = M$, so the homothety $M/N \xrightarrow{r} M/N$ is not zero since N is prime. It follows that the above homothety is injective. If $a \in N \cap rM$, then there is $b \in M$ such that $a = rb$. Since $r(b+N) = 0$, so $b \in N$, hence $rN = N \cap rM$, as required. $\square$

THEOREM 2.6. *Let M be a P -second module over a commutative ring R , and let N be a prime submodule of M . Then every submodule of M properly containing N is an RD-submodule. In particular, it is P -second.*

PROOF. Let K be a submodule of M properly containing N . Then K/N is a prime submodule of prime and P -second module M/N , so by Proposition 2.5, K/N is an RD-submodule of M/N . Now the assertion follows from [3, Consequences 18-2.2(c)] and Proposition 2.5. $\square$

LEMMA 2.7. *Let M be a nonzero module over a commutative domain R . Then M is (0) -second if and only if M is (0) -secondary.*

PROOF. The proof is completely straightforward. $\square$

By [3, Proposition 11-3.11] and [11, Proposition 12, page 506] (see also [14]), and the definitions of secondary and primary modules, we obtain the following corollary.

COROLLARY 2.8. *Let R be a commutative ring.*

- (i) *Every Artinian primary module over R is secondary.*
- (ii) *Every Noetherian secondary module over R is primary.*
- (iii) *Every finitely generated secondary module is primary.*

LEMMA 2.9. *Let R be a commutative ring. Let K and N be submodules of an R -module M such that N is prime and K is P -secondary. Then $N \cap K$ is P -secondary.*

PROOF. Let $r \in R$. If $r \in P$, then $r^n(N \cap K) \subseteq r^n K = 0$ for some n . Suppose $r \notin P$ and $t \in N \cap K$. Then $t = rs$ for some $s \in K$ since K P -secondary. As N is prime, we have $s \in N$, and hence $t \in r(N \cap K)$. This gives, $N \cap K = r(N \cap K)$. $\square$

THEOREM 2.10. *Let M be a representable module over a commutative ring R , and let N be a prime submodule of M with $(N : M) = P$. Then the following hold:*

- (i) N is representable;
- (ii) M/N is P -secondary.

PROOF. (i) Let M be a representable R -module and let $M = \sum_{i=1}^m M_i$ be a minimal secondary representation with $\text{nilrad}(M_i) = P_i$. For each i , $i = 1, 2, \dots, m$, let $m_i \in M_i$ and $r_i \in P_i$. Then $r_i^{n_i} m_i = 0$ for some n_i , and we have $(r_i^{n_i} + P)(m_i + M_i) = 0$ and hence either $P_i \subseteq P$ or $M_i \subseteq N$ ($i = 1, 2, \dots, m$). It follows that $M_i \not\subseteq N$ for some i (otherwise $M = N$). If $M_i \not\subseteq N$ and $M_j \not\subseteq N$ for $i \neq j$, then $P = P_i = P_j$, a contradiction (for if $t \in P - P_i$ then $M_i = tM_i \subseteq tM \subseteq N$). Therefore, without loss of generality, we can assume that $M_1 \not\subseteq N$ and $M_i \subseteq N$, so $P_1 = P$ and $P_i \not\subseteq P$ ($i = 2, 3, \dots, m$). Then $M_2 + M_3 + \dots + M_m \subseteq N$ and

$$N = N \cap M = N \cap (M_1 + \dots + M_m) = M_2 + \dots + M_m + (N \cap M_1). \quad (2.1)$$

Now the assertion follows from [Lemma 2.9](#).

- (ii) Since $M = M_1 + N$, we have $M/N = (M_1 + N)/N \cong M_1/(M_1 \cap N)$, as required. $\square$

PROPOSITION 2.11. *Let R be a Dedekind domain, and let M be a $0 \neq P$ -secondary R -module. Then M is a P -primary module.*

PROOF. Let $r \in R$. If $r \in P$, then the homothety $M \xrightarrow{r} M$ is nilpotent since M is secondary. Suppose that $r \notin P$. If $ra = 0$ for some $0 \neq a \in M$, then by [6, Theorem 6.1], there exists $0 \neq b \in M$ and $Q \in \text{Ass}(M)$ such that $r \in Q$ and $Q = (0 :_R b)$. As $(0 : M) \subseteq (0 : b) = Q$, we have $P = Q$, a contradiction. So the homothety $M \xrightarrow{r} M$ is injective, as required. $\square$

REMARKS. (i) Let R be a domain which is not a field. Then R is a prime R -module (since R is torsion-free) but it is not secondary (even it is not pure-injective).

(ii) Let R be a local Dedekind domain with maximal ideal $P = Rp$. We show that the module $E(R/P)$ is not prime (but it is (0) -secondary). Set $E = E(R/P)$ and $A_n = (0 :_E P^n)$ ($n \geq 1$). Then by [2, Lemma 2.6], $PA_{n+1} = A_n$, $A_n \subseteq E$ is a cyclic R -module with $A_n = Ra_n$ such that $pa_{n+1} = a_n$, every nonzero proper submodule L of E is of the form $L = A_m$ for some m and E is Artinian module with a strictly increasing sequence of submodules

$$A_1 \subset A_2 \subset \dots \subset A_n \subset A_{n+1} \subset \dots \quad (2.2)$$

We claim that $(A_n :_R E) = 0$ for every n . Suppose that $r \in (A_n :_R E)$ with $r \neq 0$. Then $rE \subseteq A_n$ and for all $a \in M$, we have $a = rb$ for some $b \in M$ since E is injective (= divisible). Thus $a = rb \in A_n$, so $E = A_n$, a contradiction. Therefore $(A_n :_R E) = 0$ for

every integer $n \geq 1$. However no A_n is a prime submodule of E , for if m is any positive integer, then $p^m \notin (A_n :_R E) = 0$ and $a_{n+m} \notin A_n$, but $p^m a_{m+n} = a_n \in A_n$.

THEOREM 2.12. *Let R be a Dedekind domain, and let M be an R -module. Then M is $0 \neq P$ -second if and only if M is P -prime.*

PROOF. By Proposition 2.11, it is enough to show that if M is P -prime, then M is P -second. Since $(0 : M) = P$ is a maximal ideal in R , so M is a vector space over R/P , hence M is P -second. $\square$

PROPOSITION 2.13. *Let R be a Dedekind domain. Then any $0 \neq P$ -prime R -module is a direct sum of copies of $R_P/PR_P \cong R/P$.*

PROOF. By the proof of Proposition 2.11, every element of $R - P$ acts invertibly on M , so the R -module structure of M extends naturally to a structure of M as a module over the localisation R_P of R at P . Therefore, we can assume that R is a commutative local Dedekind domain with maximal ideal $P = Rp$. Let M_j denote the indecomposable summand of M , so M_j is P -prime. Let m_j be a nonzero element of M_j , hence $(0 : m_j) = (0 : M) = P$. Then $Rm_j \cong R/P$ is pure in M_j since m_j is not divisible by p in M_j , but by [1, Proposition 1.3], the module R/P is itself pure-injective, so Rm_j is a direct summand of M_j , and hence $M_j \cong Rm_j$, as required. $\square$

3. Pure-injective modules

PROPOSITION 3.1. *Let M be a P -secondary module over a commutative ring R . Then $H = H(M)$, the pure-injective hull, is P -secondary.*

PROOF. Let $r \in R$. If $r \notin P$, then $rM = M$, so M satisfies the sentence for all x there exists y ($x = ry$), and hence so does H (because any module and its pure-injective hull satisfy the same sentences [7, Chapter 4]). If $r \in R$, then $r^n M = 0$, so M satisfies the sentence for all x ($r^n x = 0$), hence so does in H , as required. $\square$

THEOREM 3.2. *The following conditions are equivalent for a Prufer domain R :*

- (i) *the ring R is a Dedekind domain;*
- (ii) *every secondary R -module is pure-injective.*

PROOF. Let R be a Dedekind domain and M a secondary R -module. If $\text{Ann}(M) = 0$, then M is divisible, hence injective. If $\text{Ann}(M) \neq 0$, then M is a torsion R -module of bounded order, so that M is Σ -pure-injective (see [15]). In both cases, M is Σ -pure-injective (so pure-injective).

Conversely, let R be a Prufer domain with the property that every secondary module is pure-injective. In order to prove that R is Dedekind domain, it suffices to show that every divisible R -module is injective. Let M be a divisible R -module. Then M is secondary, Hence pure-injective. Since R is Prufer, pure-injective modules are RD-injective (see [7]). The embedding of M in its injective envelope $E(M)$ is an RD-pure monomorphism, because for every nonzero $r \in R$ we have that $M = rM$, so that $rE(M) \cap M \subseteq M \subseteq rM$. Since M is the RD-injective, M is a direct summand of $E(M)$. Thus M is injective. This shows that R is a Dedekind domain. $\square$

REMARKS. (i) There is a module over a commutative regular ring which is injective but not secondary (see [9, Theorem 2.3]). The commutative regular ring $R = F \times F$, F a field, is an Artinian Gorenstein, that is, R is injective (so pure-injective) as an R -module. But R is not secondary, because multiplication by $(1, 0)$ is neither nilpotent nor surjective.

(ii) The above consideration thus leads us to the following question: are secondary modules pure-injective? The answer is yes because of the following reason. Every non-Noetherian Prufer domain has secondary modules that are not pure-injective. For instance, every non-Noetherian valuation domain has secondary modules that are not pure-injective.

PROPOSITION 3.3. *Let M be an R -module.*

- (i) *M is Σ -secondary if and only if M is secondary.*
- (ii) *Let M be a direct sum of modules M_i ($i \in I$) where for each i , M_i is secondary and $\text{Ann}(M_i) = \text{Ann}(M_j)$ for all $i, j \in I$. Then M is secondary.*

PROOF. (i) The necessity is immediate by the definition. Conversely, suppose that M is P -secondary. Given an index set J , and let $r \in R$. If $r \in P$, then $r^n M = 0$ for some n , so $r^n M^{(J)} = 0$. If $r \notin P$ then $rM = M$, so $rM^{(J)} = M^{(J)}$, as required.

(ii) Since the annihilators of all direct summands coincide, we can assume that M_i is P -secondary (say) for all $i \in I$. Now the proof of (ii) is similar to that (i) and we omit it. $\square$

COROLLARY 3.4. *Let M be an indecomposable Σ -pure-injective module over a commutative Prufer ring R . Then M is secondary.*

PROOF. Set $P = \{r \in R : \text{Ann}_M r \neq 0\}$ and $P' = \bigcap_n P^n$. Then P and P' are prime ideals in R by [8, Fact 3.1 and Lemma 2.1]. By [8, Fact 3.2], M is either P -secondary or P' -secondary, as required. $\square$

COROLLARY 3.5. *Every Σ -pure-injective module over a Prufer ring is representable.*

PROOF. Suppose M is a Σ -pure-injective module over a commutative Prufer ring R . By [8, page 967], we can write $M = M_1 \oplus \cdots \oplus M_m$ where M_i is secondary for all i by Proposition 3.3 and Corollary 3.4, as required. $\square$

ACKNOWLEDGMENT. The author thanks the referee for useful comments.

REFERENCES

- [1] S. E. Atani, *On pure-injective modules over pullback rings*, Comm. Algebra **28** (2000), no. 9, 4037–4069.
- [2] ———, *On secondary modules over Dedekind domains*, Southeast Asian Bull. Math. **25** (2001), no. 1, 1–6.
- [3] J. Dauns, *Modules and Rings*, Cambridge University Press, Cambridge, 1994.
- [4] S. M. George, R. L. McCasland, and P. F. Smith, *A principal ideal theorem analogue for modules over commutative rings*, Comm. Algebra **22** (1994), no. 6, 2083–2099.
- [5] I. G. Macdonald, *Secondary representation of modules over a commutative ring*, Symposia Mathematica, Vol. 11 (Convegno di Algebra Commutativa, INDAM, Rome, 1971), Academic Press, London, 1973, pp. 23–43.

- [6] H. Matsumura, *Commutative Ring Theory*, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1986.
- [7] M. Prest, *Model Theory and Modules*, London Mathematical Society Lecture Note Series, vol. 130, Cambridge University Press, Cambridge, 1988.
- [8] M. Prest and G. Puninski, Σ -pure-injective modules over a commutative Prüfer ring, *Comm. Algebra* **27** (1999), no. 2, 961–971.
- [9] R. Y. Sharp, *Secondary representations for injective modules over commutative Noetherian rings*, *Proc. Edinburgh Math. Soc. (2)* **20** (1976), no. 2, 143–151.
- [10] ———, *Steps in Commutative Algebra*, London Mathematical Society Student Texts, vol. 19, Cambridge University Press, Cambridge, 1990.
- [11] W. V. Vasconcelos, *On finitely generated flat modules*, *Trans. Amer. Math. Soc.* **138** (1969), 505–512.
- [12] R. B. Warfield Jr., *Purity and algebraic compactness for modules*, *Pacific J. Math.* **28** (1969), 699–719.
- [13] R. Wisbauer, *Foundations of Module and Ring Theory*, Algebra, Logic and Applications, vol. 3, Gordon and Breach Science Publishers, Pennsylvania, 1991.
- [14] S. Yassemi, *The dual notion of prime submodules*, to appear in *Arch. Math.*
- [15] W. Zimmermann, *Rein injektive direkte Summen von Moduln*, *Comm. Algebra* **5** (1977), no. 10, 1083–1117 (German).

SHAHABADDIN EBRAHIMI ATANI: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GUILAN,
P.O. BOX 1914, RASHT, IRAN

E-mail address: ebrahimi@cd.gu.ac.ir