THE UNION PROBLEM ON COMPLEX MANIFOLDS

PATRICK W. DARKO

Received 14 May 2001

Let Ω be a relatively compact subdomain of a complex manifold, exhaustable by Stein open sets. We give a necessary and sufficient condition for Ω to be Stein, in terms of L^2 -estimates for the $\bar{\partial}$ -operator, equivalent to the condition of Markoe (1977) and Silva (1978).

2000 Mathematics Subject Classification: 32E10, 32C35, 35N15.

1. Introduction. As indicated in [7], from the beginning of the theory of Stein spaces, the following question has held great interest: is a complex space, which is exhaustable by a sequence $X_1 \Subset X_2 \Subset \cdots$ of Stein subspaces, itself Stein?

In [1], the following is proved: every domain in $\mathbb{C}^m$ which is exhaustable by a sequence of Stein domains $B_1 \Subset B_2 \Subset \cdots$ is itself Stein, and this is shown to hold more generally for unramified Riemann domain $\mathcal{B}$ over $\mathbb{C}^m$ in [6]. In [11], the following is proved: let X be a reduced complex space and $X_1 \Subset X_2 \Subset \cdots$ be an exhaustion of X by Stein domains, if every pair (X_j, X_{j+1}) is Runge then $X = \bigcup X_j$ is Stein. Recently, Markoe [9] and Silva [10] proved the following: let X be reduced and $X_1 \Subset X_2 \Subset \cdots$ be an exhaustion of X by Stein domains. Then X is Stein if and only if $H^1(X, \mathcal{O}) = 0$ ($\mathcal{O}$ being the structure sheaf of X).

More recently the following has been proved in [12]: let $\Omega_1 \subset \Omega_2 \subset \cdots$ be a sequence of open Stein subsets of a Stein space X , $\Omega = \bigcup_{j=1}^{\infty} \Omega_j$, and $\dim H^1(\Omega, \mathcal{O}) < \infty$. Then Ω is Stein.

Fornæss [4] produced an example to show that if $X_1 \Subset X_2 \Subset \cdots$ is a sequence of Stein manifolds, the limit manifold $X = \bigcup X_j$, in which each X_j is an open submanifold, need not be Stein. But it is known that if the limit manifold is itself an open submanifold of a Stein manifold then the limit manifold is necessarily Stein.

This led Fornæss and Narasimhan to pose the following problem [5]: let X be a Stein space and $\Omega_1 \Subset \Omega_2 \Subset \cdots$ an increasing sequence of Stein open sets in X . Is $\bigcup \Omega_j$ Stein? As indicated above this is the case when X is a Stein manifold, but this question remains open in the general case.

In this paper, we consider the case where X is a general complex manifold and $\Omega_1 \Subset \Omega_2 \Subset \cdots$ an increasing sequence of open Stein manifolds in X such that $\Omega = \bigcup \Omega_j$ is relatively compact in X . We give a condition for Ω to be Stein, equivalent to Markoe's and Silva's condition and involving L^2 -estimates for the $\bar{\partial}$ operator.

2. Preliminaries. Let X be an n -dimensional complex manifold with a C^∞ Hermitian metric. The space $L^2_{(p,q)}(X)$ of square integrable differential forms of type (p, q) on X

is a Hilbert space under the scalar product,

$$(f, g) = \int_X f \wedge * \bar{g}, \quad (2.1)$$

where $*$ is the Hodge $*$ -operator associated with the metric and orientation of X .

Let $\Omega_1 \Subset \Omega_2 \Subset \cdots$ be an increasing sequence of Stein open sets in X such that their union $\Omega = \bigcup_{j=1}^{\infty} \Omega_j$ is relatively compact in X .

The following theorem is our main result.

THEOREM 2.1. *The union Ω is Stein if and only if given an $f \in L^2_{(p,q)}(\Omega)$, which is $\bar{\partial}$ -closed in the sense of distributions, there is a $u \in L^2_{(p,q-1)}(\Omega)$ such that $\bar{\partial}u = f$ in the sense of distributions and*

$$\|u\|_{L^2_{(p,q-1)}(\Omega)} \leq K \|f\|_{L^2_{(p,q)}(\Omega)}, \quad q > 0, \quad (2.2)$$

where K depends on Ω .

Let U be a bounded open set in $\mathbb{C}^n$, and $\mathcal{O}$ the structure sheaf of $\mathbb{C}^n$. A section $f = (f_1, \dots, f_p) \in \Gamma(U, \mathcal{O}^p)$, where $p > 0$ is an integer, is L^2 -bounded if

$$\|f\|_{L^2(U)} = \|f_1\|_{L^2(U)} + \cdots + \|f_p\|_{L^2(U)} < \infty. \quad (2.3)$$

We then denote all sections of $\mathcal{O}^p$ over U that are L^2 -bounded by $\Gamma_2(U, \mathcal{O}^p)$.

For the definition of L^2 -bounded sections of coherent analytic sheaves, we require the coherent analytic sheaf $\mathcal{F}$ to be defined on a simply connected polycylinder neighborhood V of the closure of U . Then by [8, Theorem 5, Section F, Chapter VI], there is an $\mathcal{O}$ -homographic in another simply connected polycylinder neighborhood V' of the closure of U ,

$$\mathcal{O}^p \xrightarrow{\lambda} \mathcal{F} \rightarrow 0, \quad (2.4)$$

where $p > 0$ is some integer; and $f \in \Gamma(U, \mathcal{F})$ is L^2 -bounded if $f \in \Gamma_2(U, \mathcal{F}) := \lambda(\Gamma_2(U, \mathcal{O}^p))$. It can be shown that $\Gamma_2(U, \mathcal{F})$ is independent of λ and p , so that $\Gamma_2(U, \mathcal{F})$ is well defined.

Now let Ω be a relatively compact subdomain of an n -dimensional complex manifold X . An open subset Y of Ω is said to be admissible for the coherent analytic sheaf $\mathcal{F}$ defined in the neighborhood of the closure of Ω in X , if Y is Stein. There is a coordinate neighborhood V in X of the closure, $\bar{Y}$ of Y such that V is biholomorphic to a simply connected polycylinder V' in $\mathbb{C}^n$, and $\bar{Y}$ is contained in the neighborhood of $\bar{\Omega}$ where $\mathcal{F}$ is defined as $f \in \Gamma(Y, \mathcal{F})$ which is L^2 -bounded if

$$f \in \Gamma_2(Y, \mathcal{F}) := \{g \in \Gamma(Y, \mathcal{F}) : \eta_*(g) \in \Gamma_2(\eta(Y), \eta_*(\mathcal{F}))\}, \quad (2.5)$$

where η is the restriction of the biholomorphic map $V \rightarrow V'$ to Y , and $\eta_*(\mathcal{F})$ is the zero direct image of $\mathcal{F}$ on Y .

Let Ω be as in Theorem 2.1 (then clearly Ω is locally Stein). Let $\mathcal{F}$ be a coherent analytic sheaf in a neighborhood of the closure of Ω . Then it is clear that Ω is a finite union, $\Omega = \bigcup_{j=1}^m U_j$, where each U_j is admissible for $\mathcal{F}$. If $\mathcal{V} = \{U_j\}_{j \in I}$, $I = \{1, \dots, m\}$,

where the U_j 's are as above, we say that $\mathcal{V}$ is a finite admissible cover of Ω for $\mathcal{F}$ and we define the L^2 (alternate) q -cochains of $\mathcal{V}$ with values in $\mathcal{F}$ as those cochains,

$$\begin{aligned} c &= (c_\alpha) \in C^q(\mathcal{V}, \mathcal{F}) = \prod_{\alpha \in I^{q+1}} \Gamma(U_\alpha, \mathcal{F}), \\ U_\alpha &= U_{i_0} \cap \cdots \cap U_{i_q}, \quad \alpha = (i_0, \dots, i_q), \end{aligned} \quad (2.6)$$

which are alternate and satisfy $c_\alpha \in \Gamma_2(U_\alpha, \mathcal{F})$ for all $\alpha \in I^{q+1}$. We denote by $C_2^q(\mathcal{V}, \mathcal{F})$ the space of L^2 -bounded cochains.

The coboundary operator,

$$\delta : C^q(\mathcal{V}, \mathcal{F}) \longrightarrow C^{q+1}(\mathcal{V}, \mathcal{F}), \quad (2.7)$$

maps $C_2^q(\mathcal{V}, \mathcal{F})$ into $C_2^{q+1}(\mathcal{V}, \mathcal{F})$. If $Z_2^q(\mathcal{V}, \mathcal{F}) = \{c \in C_2^q(\mathcal{V}, \mathcal{F}) : \delta c = 0\}$ and $B_2^q(\mathcal{V}, \mathcal{F}) = \delta C_2^{q-1}(\mathcal{V}, \mathcal{F})$, then as usual $B_2^q(\mathcal{V}, \mathcal{F}) \subseteq Z_2^q(\mathcal{V}, \mathcal{F})$ and we define $H_2^q(\mathcal{V}, \mathcal{F}) := Z_2^q(\mathcal{V}, \mathcal{F}) / B_2^q(\mathcal{V}, \mathcal{F})$ and call it the L^2 -bounded cohomology of $\mathcal{V}$ with values in $\mathcal{F}$. We then have the following theorem.

THEOREM 2.2. *For any $q > 0$, the natural map*

$$H_2^q(\mathcal{V}, \mathcal{F}) \longrightarrow H^q(\Omega, \mathcal{F}) \quad (2.8)$$

is an isomorphism.

We use [Theorem 2.2](#) as a pivot to prove [Theorem 2.1](#), but the proof of [Theorem 2.2](#) is not given here, since it is similar to that of [\[2, Theorem\]](#).

3. A triangle of isomorphisms. Let Ω be as in [Theorem 2.1](#). By the end of the section [Theorem 2.1](#) will be proved. If $U \neq \emptyset$ is an open set in $\tilde{\Omega}$, then $\mathcal{B}_\Omega^p(U)$ is the Hilbert space of holomorphic p -forms h on $\Omega \cap U$ such that

$$\|h\|_{L_{(p,0)}^2(\Omega \cap U)} < \infty. \quad (3.1)$$

If V is open in $\tilde{\Omega}$ with $\emptyset \neq V \subset U$, the restriction map $\gamma_V^U : \mathcal{B}_\Omega^p(U) \rightarrow \mathcal{B}_\Omega^p(V)$ is defined. Then $\mathcal{B}_\Omega^p = \{\mathcal{B}_\Omega^p(U), \gamma_V^U\}$ is the canonical presheaf of L^2 -holomorphic p -forms on $\tilde{\Omega}$. The associated sheaf $\mathcal{B}_2^p$ is the sheaf of germs of L^2 -holomorphic p -forms on $\tilde{\Omega}$. We then have the following lemma.

LEMMA 3.1. *Let $\mathcal{D}^p$ be the sheaf of germs of holomorphic p -forms on X , and $\mathcal{V}$ a finite admissible cover of Ω for $\mathcal{D}^p$. Then the following diagram is an isomorphism triangle of cohomology groups:*

$$\begin{array}{ccc} H_2^q(\mathcal{V}, \mathcal{D}^p) & \xrightarrow{\sim} & H^q(\Omega, \mathcal{D}^p) \\ & \searrow & \nearrow \\ & H^q(\tilde{\Omega}, \mathcal{B}_2^p) & \end{array} \quad (3.2)$$

for $q \geq 1$ and $p \geq 0$.

PROOF. From [Theorem 2.2](#) and the fact that any finite cover of $\bar{\Omega}$ has a refinement $\mathcal{U} = \{V_j\}_{j \in J}$ such that $\mathcal{U}_\Omega = \{V_j \cap \Omega\}_{j \in J}$ is a finite admissible cover of Ω for $\mathcal{D}^p$, the lemma follows. $\square$

Now, using Hörmander's L^2 -estimates locally we get the following lemma.

LEMMA 3.2. *The cohomology group $H^q(\bar{\Omega}, \mathcal{B}_2^p)$ is isomorphic to the quotient space*

$$\{g : g \in L_{(p,q)}^2(\Omega) \text{ and } \bar{\partial}g = 0\} / \{\bar{\partial}h : h \in L_{(p,q-1)}^2(\Omega) \text{ and } \bar{\partial}h \in L_{(p,q)}^2(\Omega)\}, \quad (3.3)$$

where Ω is as in [Theorem 2.1](#).

Also the following lemma is proved in [3].

LEMMA 3.3. *If $\Omega \Subset X$ is Stein, where X is a complex manifold, then given $f \in L_{(p,q)}^2(\Omega)$ with $\bar{\partial}f = 0$, there is $u \in L_{(p,q-1)}^2(\Omega)$ such that*

$$\bar{\partial}u = f, \quad \|u\|_{L_{(p,q-1)}^2(\Omega)} \leq K \|f\|_{L_{(p,q)}^2(\Omega)}, \quad (3.4)$$

where K depends on Ω .

To finish with the proof of [Theorem 2.1](#) we remark that $\mathcal{D}^0 = \mathcal{O}$ is the structure sheaf of X (X, Ω as in [Theorem 2.1](#)), therefore [Theorem 2.1](#) follows from [Lemmas 3.1, 3.2, and 3.3](#), and from Markoe's and Silva's condition.

REFERENCES

- [1] H. Behnke and K. Stein, *Konvergente folge von regularit ätsbereichen und die meromorphiekonvexität*, Math. Ann. **116** (1938), 204–216 (German).
- [2] P. W. Darko, *On cohomology with bounds on complex spaces*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) **60** (1976), no. 3, 189–194.
- [3] ———, *L^2 estimates for the $\bar{\partial}$ operator on Stein manifolds*, Math. Proc. Cambridge Philos. Soc. **129** (2000), no. 1, 73–76.
- [4] J. E. Fornæss, *An increasing sequence of Stein manifolds whose limit is not Stein*, Math. Ann. **223** (1976), no. 3, 275–277.
- [5] J. E. Fornæss and R. Narasimhan, *The Levi problem on complex spaces with singularities*, Math. Ann. **248** (1980), no. 1, 47–72.
- [6] H. Grauert and R. Remmert, *Konvexität in der komplexen Analysis. Nicht-holomorph-konvexe Holomorphiegebiete und Anwendungen auf die Abbildungstheorie*, Comment. Math. Helv. **31** (1956), 152–160, 161–183 (German).
- [7] ———, *Theory of Stein Spaces*, Springer-Verlag, Berlin, 1979.
- [8] R. C. Gunning and H. Rossi, *Analytic Functions of Several Complex Variables*, Prentice-Hall, New Jersey, 1965.
- [9] A. Markoe, *Runge families and inductive limits of Stein spaces*, Ann. Inst. Fourier (Grenoble) **27** (1977), no. 3, 117–127.
- [10] A. Silva, *Rungescher Satz and a condition for Steinness for the limit of an increasing sequence of Stein spaces*, Ann. Inst. Fourier (Grenoble) **28** (1978), no. 2, 187–200 (German).
- [11] K. Stein, *Überlagerungen holomorph-vollständiger komplexer Räume*, Arch. Math. **7** (1956), 354–361 (German).

- [12] L. M. Tovar, *Open Stein subsets and domains of holomorphy in complex spaces*, Topics in Several Complex Variables (Mexico, 1983), Pitman, Massachusetts, 1985, pp. 183–189.

PATRICK W. DARKO: DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, LINCOLN UNIVERSITY, LINCOLN UNIVERSITY, PA 19352, USA

E-mail address: pdarko@lu.lincoln.edu