A NOTE ON RUSCHEWEYH TYPE OF INTEGRAL OPERATORS FOR UNIFORMLY α -CONVEX FUNCTIONS

M. ANBU DURAI and R. PARVATHAM

Received 7 March 2001

We prove that the class of uniformly α -convex functions introduced by Kanas is closed under the generalized Ruscheweyh integral operator for $0 < \alpha \leq 1$.

2000 Mathematics Subject Classification: 30C45.

We denote by $\mathcal{A}$ the class of functions $f(z) = z + a_2 z^2 + \dots$ which are analytic in $\Delta = \{z \in \mathbb{C} : |z| < 1\}$. Let S denote the class of functions in $\mathcal{A}$ that are univalent in Δ . The subclasses of S containing functions which are uniformly convex and uniformly starlike, introduced by Goodman [1, 2], are denoted by UCV and UST , respectively.

The class of uniformly α -convex functions was introduced by Kanas [3] and she gave an analytic condition for such functions as follows: $f(z)$ is a uniformly α -convex function if and only if

$$\operatorname{Re} \left\{ (1 - \alpha) \frac{(z - \zeta) f'(z)}{f(z) - f(\zeta)} + \alpha \left(1 + \frac{(z - \zeta) f''(z)}{f'(z)} \right) \right\} > 0 \quad (1)$$

for all $z, \zeta \in \Delta$ and $0 \leq \alpha \leq 1$. For $\zeta = 0$, this class of functions reduces to Mocanu's class $M(\alpha)$ of α -convex functions [4].

In this note, for $\alpha > 0$, we consider the integral operator

$$F(z) = \frac{F_\alpha(z, \zeta) - F_\alpha(0, \zeta)}{F'_\alpha(0, \zeta)}, \quad (2)$$

where

$$F_\alpha(z, \zeta) = \left\{ \frac{c + 1/\alpha}{(z - \zeta)^c} \int_\zeta^z (t - \zeta)^{c-1} (f(t) - f(\zeta))^{1/\alpha} dt \right\}^\alpha \quad (3)$$

for all $z \in \Delta$ and for fixed $\zeta \in \Delta$ with $z \neq \zeta$. We prove that this normalized function $F(z)$ is a uniformly α -convex function when $f(z)$ is a uniformly α -convex function in the sense of Kanas [3].

For $\zeta = 0$ the operator $F(z)$ reduces to Ruscheweyh's integral operator [5]. It is well known that Mocanu's class $M(\alpha)$ of α -convex functions is closed under Ruscheweyh's integral operator for $\alpha > 0$.

THEOREM 1. Let $f(z) = z + a_2 z^2 + \dots$ be a uniformly α -convex function in Δ and let $c > 0$. Then, for $0 < \alpha \leq 1$, the function

$$F(z) = \frac{F_\alpha(z, \zeta) - F_\alpha(0, \zeta)}{F'_\alpha(0, \zeta)}, \quad z, \zeta \in \Delta, \quad (4)$$

is uniformly α -convex where $F_\alpha(z, \zeta)$ is defined as in (3).

PROOF. We have from (3) that

$$F_\alpha^{1/\alpha}(z, \zeta) = \frac{c+1/\alpha}{(z-\zeta)^c} \int_\zeta^z (t-\zeta)^{c-1} (f(t) - f(\zeta))^{1/\alpha} dt. \quad (5)$$

Differentiating with respect to z , we have

$$\begin{aligned} (z-\zeta)^c \frac{1}{\alpha} F_\alpha^{1/\alpha-1}(z, \zeta) F'_\alpha(z, \zeta) + c(z-\zeta)^{c-1} F_\alpha^{1/\alpha}(z, \zeta) \\ = \left(c + \frac{1}{\alpha}\right) (z-\zeta)^{c-1} (f(z) - f(\zeta))^{1/\alpha} \end{aligned} \quad (6)$$

and again differentiating with respect to z we get

$$\begin{aligned} \frac{1}{\alpha} \left\{ (z-\zeta) F_\alpha^{1/\alpha-1}(z, \zeta) F''_\alpha(z, \zeta) + (z-\zeta) \left(\frac{1}{\alpha} - 1 \right) F_\alpha^{1/\alpha-2}(z, \zeta) (F'_\alpha(z, \zeta))^2 \right. \\ \left. + F_\alpha^{1/\alpha-1}(z, \zeta) F'_\alpha(z, \zeta) \right\} + \frac{c}{\alpha} F_\alpha^{1/\alpha-1}(z, \zeta) F'_\alpha(z, \zeta) \\ = \left(c + \frac{1}{\alpha}\right) \frac{1}{\alpha} f'(z) (f(z) - f(\zeta))^{1/\alpha-1}; \\ F_\alpha^{1/\alpha-1}(z, \zeta) F'_\alpha(z, \zeta) \left\{ \alpha \frac{(z-\zeta) F''_\alpha(z, \zeta)}{F'_\alpha(z, \zeta)} + (1-\alpha)(z-\zeta) \frac{F'_\alpha(z, \zeta)}{F_\alpha(z, \zeta)} + \alpha(1+c) \right\} \\ = (\alpha c + 1) f'(z) (f(z) - f(\zeta))^{1/\alpha-1}. \end{aligned} \quad (7)$$

Thus we get

$$\begin{aligned} F_\alpha^{1/\alpha-1}(z, \zeta) F'_\alpha(z, \zeta) \left\{ \alpha \left[1 + \frac{(z-\zeta) F''_\alpha(z, \zeta)}{F'_\alpha(z, \zeta)} - \frac{(z-\zeta) F'_\alpha(z, \zeta)}{F_\alpha(z, \zeta)} \right] \right. \\ \left. + \frac{(z-\zeta) F'_\alpha(z, \zeta)}{F_\alpha(z, \zeta)} + c\alpha \right\} \\ = (c\alpha + 1) f'(z) (f(z) - f(\zeta))^{1/\alpha-1}. \end{aligned} \quad (8)$$

From (2) we have

$$F'(z) = \frac{F'_\alpha(z, \zeta)}{F'_\alpha(0, \zeta)}, \quad (9)$$

showing that $F(0) = 0$ and $F'(0) = 1$.

Considering

$$\frac{(z-\zeta) F'(z)}{F(z) - F(\zeta)} = \frac{(z-\zeta) F'_\alpha(z, \zeta)}{F_\alpha(z, \zeta)} \quad (10)$$

and differentiating with respect to z , we have

$$\frac{F''(z)}{F'(z)} + \frac{1}{z-\zeta} - \frac{F'(z)}{F(z)-F(\zeta)} = \frac{1}{z-\zeta} + \frac{F''_{\alpha}(z, \zeta)}{F'_{\alpha}(z, \zeta)} - \frac{F'_{\alpha}(z, \zeta)}{F_{\alpha}(z, \zeta)}; \quad (11)$$

$$\frac{(z-\zeta)F''(z)}{F'(z)} + 1 - \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} = 1 + \frac{(z-\zeta)F''_{\alpha}(z, \zeta)}{F'_{\alpha}(z, \zeta)} - \frac{F'_{\alpha}(z, \zeta)(z-\zeta)}{F_{\alpha}(z, \zeta)}. \quad (12)$$

Substituting (10) and (12) in (8), we obtain

$$\begin{aligned} & F_{\alpha}^{1/\alpha-1}(z, \zeta)F'_{\alpha}(z, \zeta) \left\{ \alpha \left[\frac{(z-\zeta)F''(z)}{F'(z)} + 1 - \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} \right] + \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} + c\alpha \right\} \\ &= (c\alpha+1)f'(z)(f(z)-f(\zeta))^{1/\alpha-1}; \\ & F_{\alpha}^{1/\alpha-1}(z, \zeta)F'_{\alpha}(z, \zeta) \left\{ (1-\alpha) \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} + \alpha \left(1 + \frac{(z-\zeta)F''(z)}{F'(z)} \right) + c\alpha \right\} \\ &= (c\alpha+1)f'(z)(f(z)-f(\zeta))^{1/\alpha-1}. \end{aligned} \quad (13)$$

Setting

$$P(z, \zeta) = (1-\alpha) \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} + \alpha \left(\frac{(z-\zeta)F''(z)}{F'(z)} + 1 \right), \quad (14)$$

equation (13) becomes

$$F_{\alpha}^{1/\alpha-1}(z, \zeta)F'_{\alpha}(z, \zeta) \{P(z, \zeta) + c\alpha\} = (c\alpha+1)f'(z)(f(z)-f(\zeta))^{1/\alpha-1}. \quad (15)$$

Taking logarithmic differentiation with respect to z , we get

$$\begin{aligned} & (1-\alpha)(z-\zeta) \frac{F'_{\alpha}(z, \zeta)}{F_{\alpha}(z, \zeta)} + \alpha(z-\zeta) \frac{F''_{\alpha}(z, \zeta)}{F'_{\alpha}(z, \zeta)} + \frac{\alpha(z-\zeta)P'(z, \zeta)}{P(z, \zeta) + c\alpha} + \alpha \\ &= \alpha + \alpha \frac{(z-\zeta)f''(z)}{f'(z)} + (1-\alpha) \frac{(z-\zeta)f'(z)}{f(z)-f(\zeta)}; \\ & \alpha \left[(z-\zeta) \frac{F''_{\alpha}(z, \zeta)}{F'_{\alpha}(z, \zeta)} + 1 - \frac{(z-\zeta)F'_{\alpha}(z, \zeta)}{F_{\alpha}(z, \zeta)} \right] + \frac{(z-\zeta)F'_{\alpha}(z, \zeta)}{F_{\alpha}(z, \zeta)} + \frac{\alpha(z-\zeta)P'(z, \zeta)}{P(z, \zeta) + c\alpha} \\ &= \alpha \left(1 + \frac{(z-\zeta)f''(z)}{f'(z)} \right) + (1-\alpha) \frac{(z-\zeta)f'(z)}{f(z)-f(\zeta)}. \end{aligned} \quad (16)$$

Equations (10) and (12) give

$$\begin{aligned} & \alpha \left[\frac{(z-\zeta)F''(z)}{F'(z)} + 1 - \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} \right] + \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} + \frac{\alpha(z-\zeta)P'(z, \zeta)}{P(z, \zeta) + c\alpha} \\ &= \alpha \left(1 + \frac{(z-\zeta)f''(z)}{f'(z)} \right) + (1-\alpha) \frac{(z-\zeta)f'(z)}{f(z)-f(\zeta)}. \end{aligned} \quad (17)$$

That is

$$\begin{aligned} & \left[\alpha \left(1 + \frac{(z-\zeta)F''(z)}{F'(z)} \right) + (1-\alpha) \frac{(z-\zeta)F'(z)}{F(z)-F(\zeta)} \right] + \frac{\alpha(z-\zeta)P'(z, \zeta)}{P(z, \zeta) + c\alpha} \\ &= \alpha \left(1 + \frac{(z-\zeta)f''(z)}{f'(z)} \right) + (1-\alpha) \frac{(z-\zeta)f'(z)}{f(z)-f(\zeta)}. \end{aligned} \quad (18)$$

Hence, we have

$$P(z, \zeta) + \frac{\alpha(z-\zeta)P'(z, \zeta)}{P(z, \zeta) + c\alpha} = \alpha \left(1 + \frac{(z-\zeta)f''(z)}{f'(z)} \right) + (1-\alpha) \frac{(z-\zeta)f'(z)}{f(z)-f(\zeta)}. \quad (19)$$

Since $f(z)$ is uniformly α -convex, we have

$$\operatorname{Re} \left\{ P(z, \zeta) + \frac{\alpha(z-\zeta)P'(z, \zeta)}{P(z, \zeta) + c\alpha} \right\} > 0 \quad (20)$$

for all $z, \zeta \in \Delta$, $0 \leq \alpha \leq 1$.

We show that $\operatorname{Re} P(z, \zeta) > 0$. Suppose that there exists a point $\zeta_0 \in \Delta$ such that the image of the arc $\Gamma : z(t) = \zeta_0 + re^{it}$ is tangent to the imaginary axis. Let w_0 be the point of contact and let $z_0 \in \Delta$ such that $w_0 = P(z_0, \zeta_0)$. Then $\operatorname{Re} P(z_0, \zeta_0) = 0$ and therefore $P(z_0, \zeta_0) = ix$, where $x \in R$. Hence the outer normal to $F(\Gamma)$ is

$$(z_0 - \zeta_0)P'(z_0, \zeta_0) = y < 0. \quad (21)$$

For such ζ_0 , we have

$$\begin{aligned} \operatorname{Re} \left\{ P(z_0, \zeta_0) + \frac{\alpha(z_0 - \zeta_0)P'(z_0, \zeta_0)}{P(z_0, \zeta_0) + c\alpha} \right\} &= \operatorname{Re} \left\{ ix + \frac{\alpha y}{c\alpha + ix} \right\} \\ &= \operatorname{Re} \left\{ ix + \frac{\alpha y(c\alpha - ix)}{c^2\alpha^2 + x^2} \right\} \\ &= \frac{c\alpha^2 y}{(c\alpha)^2 + x^2} < 0 \quad \text{for } c > 0 \end{aligned} \quad (22)$$

which contradicts (20) and hence $\operatorname{Re} P(z, \zeta) > 0$ in Δ showing that $F(z)$ is a uniformly α -convex function. $\square$

ACKNOWLEDGEMENT. This work was carried out when the first author was under the Faculty Improvement programme of University Grant and Commission of IX plan.

REFERENCES

- [1] A. W. Goodman, *On uniformly convex functions*, Ann. Polon. Math. **56** (1991), no. 1, 87-92.
- [2] ———, *On uniformly starlike functions*, J. Math. Anal. Appl. **155** (1991), no. 2, 364-370.
- [3] S. Kanas, *Uniformly alpha convex functions*, Int. J. Appl. Math. **1** (1999), no. 3, 305-310.
- [4] P. T. Mocanu, *Une propriété de convexité généralisée dans la théorie de la représentation conforme*, Mathematica (Cluj) **11** (34) (1969), 127-133 (French).
- [5] S. Ruscheweyh, *Eine Invarianzeigenschaft der Basileviv Funktionen*, Math. Z. **134** (1973), 215-219 (German).

M. ANBU DURAI: THE RAMANUJAN INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS,
UNIVERSITY OF MADRAS, CHENNAI-600 500, INDIA
E-mail address: anbuduraim@yahoo.co.in

R. PARVATHAM: THE RAMANUJAN INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS,
UNIVERSITY OF MADRAS, CHENNAI-600 500, INDIA