ON UNIFORMLY CLOSE-TO-CONVEX FUNCTIONS AND UNIFORMLY QUASICONVEX FUNCTIONS

K. G. SUBRAMANIAN, T. V. SUDHARSAN, and HERB SILVERMAN

Received 30 October 2002

Two new subclasses of uniformly convex and uniformly close-to-convex functions are introduced. We obtain inclusion relationships and coefficient bounds for these classes.

2000 Mathematics Subject Classification: 30C45.

1. The class $UCC(\alpha)$. Denote by S the family consisting of functions

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

that are analytic and univalent in $\Delta = \{z : |z| < 1\}$ and by C , S^* , and K the subfamilies of functions that are, respectively, convex, starlike, and close to convex in Δ . Noor and Thomas [7] introduced the class of functions known as quasiconvex functions. A normalized function of the form (1.1) is said to be quasiconvex in Δ if there exists a convex function g with $g(0) = 0$, $g'(0) = 1$ such that for $z \in \Delta$,

$$\operatorname{Re} \frac{(zf'(z))'}{g'(z)} > 0. \quad (1.2)$$

Let Q denote the class of quasiconvex functions defined in Δ . It was shown that $Q < K$, where $<$ denotes subordination, so that every quasiconvex function is close to convex. Goodman [2, 3] introduced the classes UCV and UST of uniformly convex and uniformly starlike functions. In [10], Rønning defined the class $UCV(\alpha)$, $-1 \leq \alpha < 1$, consisting of functions of the form (1.1) satisfying

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) - \alpha \geq \left| \frac{zf''(z)}{f'(z)} \right|, \quad z \in \Delta. \quad (1.3)$$

Geometrically, $UCV(\alpha)$ is the family of functions f for which $1 + zf''(z)/f'(z)$ takes values that lie inside the parabola $\Omega = \{\omega : \operatorname{Re}(\omega - \alpha) > |\omega - 1|\}$, which is symmetric about the real axis and whose vertex is $w = (1 + \alpha)/2$.

Since the function

$$q_{\alpha}(z) = 1 + \frac{2(1-\alpha)}{\pi^2} \left(\log \frac{1+\sqrt{z}}{1-\sqrt{z}} \right)^2 \quad (1.4)$$

maps Δ onto this parabolic region, $f \in \text{UCV}(\alpha)$ if and only if

$$1 + \frac{zf''(z)}{f'(z)} \prec q_{\alpha}(z). \quad (1.5)$$

Rønning [10] also defined the family $S_p(\alpha)$ consisting of functions $zf'(z)$ when f is in $\text{UCV}(\alpha)$. In particular, f is in $S_p(\alpha)$ if and only if $zf'(z)/f(z) \prec q_{\alpha}(z)$.

Note for $g(z) = zf'(z)/f(z)$ that $g(z) + zg'(z)/g(z) = 1 + zf''(z)/f'(z)$, and hence a result of Miller and Mocanu [6] shows that $\text{UCV}(\alpha) \subset S_p(\alpha)$.

Kumar and Ramesha [4] investigated the class UCC of uniformly close-to-convex functions consisting of normalized functions of the form (1.1) satisfying $f'(z)/g'(z) \prec q_0(z)$, where $g(z) \in C$ and $q_0(z)$ is given by (1.4) for $\alpha = 0$.

More generally, we give the following definition.

DEFINITION 1.1. A function f is said to be uniformly close to convex of order α , $-1 \leq \alpha < 1$, denoted by $\text{UCC}(\alpha)$, if $f'(z)/g'(z) \prec q_{\alpha}(z)$, where $q_{\alpha}(z)$ is as defined by (1.4) and $g(z)$ is convex.

Since $\text{Re } q_{\alpha}(z) > 0$, we see that $\text{UCC}(\alpha)$ is a subclass of K . To see that $\text{UCC}(\alpha)$ also contains the family $S_p(\alpha)$, we note for $f \in S_p(\alpha) \subset S^*$ that $f(z) = zg'(z)$ for some $g \in C$. Hence, $zf'(z)/f(z) = f'(z)/g'(z) \prec q_{\alpha}(z)$.

We have thus proved the following inclusion chain.

THEOREM 1.2. For $-1 \leq \alpha < 1$, $\text{UCV}(\alpha) \subset S_p(\alpha) \subset \text{UCC}(\alpha) \subset K$.

We next give a sufficient condition for a function to be in $\text{UCC}(\alpha)$.

THEOREM 1.3. If $\sum_{n=2}^{\infty} n|a_n| \leq (1-\alpha)/2$, then $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is in $\text{UCC}(\alpha)$, $-1 \leq \alpha < 1$.

PROOF. Setting $g(z) = z$, we have $f'(z)/g'(z) = f'(z) = 1 + \sum_{n=2}^{\infty} n a_n z^{n-1}$, so that for $z \in \Delta$,

$$\left| \frac{f'(z)}{g'(z)} - 1 \right| < \sum_{n=2}^{\infty} n|a_n| \leq 1 - \sum_{n=2}^{\infty} n|a_n| - \alpha \leq \text{Re } f'(z) - \alpha. \quad (1.6)$$

Thus $f'(z)/g'(z)$ lies in the parabolic region $\Omega = \{\omega : |\omega - 1| < \text{Re}(\omega - \alpha)\}$. That is, $f'(z)/g'(z) \prec q_{\alpha}(z)$, where $q_{\alpha}(z)$ is as defined by (1.4). $\square$

2. A convolution relation. We now prove a convolution result for the family $\text{UCC}(\alpha)$. But first we need the following lemma.

LEMMA 2.1 (see [8]). *Let $\phi(z) \in C$, $\psi \in S^*$. If $F(z)$ is analytic and $\text{Re}\{F(z)\} > \alpha$, $-1 \leq \alpha < 1$, then*

$$\text{Re} \left\{ \frac{\phi * F \psi}{\phi * \psi} \right\} > \alpha, \quad z \in \Delta. \quad (2.1)$$

The above result was proved in [11] for the case $\alpha = 0$.

THEOREM 2.2. *If $f \in \text{UCC}(\alpha)$, then to each $g \in S^*$, an $h \in S^*$ may be associated for which $\text{Re}(f * g)/h > (1 + \alpha)/2$, $z \in \Delta$.*

PROOF. If $f \in \text{UCC}(\alpha)$, then $f'(z)/g'_1(z) \prec q_\alpha(z)$, where $g_1(z) \in C$ and $q_\alpha(z)$ is defined by (1.4). Hence, $\text{Re}(f'(z)/g'_1(z)) > (1 + \alpha)/2$. Therefore, we can find an $\psi \in S^*$ for which

$$\text{Re} \frac{zf'(z)}{\psi(z)} > \frac{1 + \alpha}{2}. \quad (2.2)$$

Set $F(z) = zf'(z)/\psi(z)$. Then, for $g \in S^*$, there corresponds a $\phi \in C$ such that $z\phi' = g$. Also $f * g = zf' * \phi = \phi * F\psi$ and $h = \phi * \psi \in S^*$. By Lemma 2.1,

$$\text{Re} \frac{\Phi * F\Psi}{\Phi * \Psi} = \text{Re} \frac{f * g}{h} > \frac{1 + \alpha}{2}, \quad (2.3)$$

and this proves the result. $\square$

3. Coefficient estimates. We need the following result by Rogosinski [9] to obtain coefficient bounds for the class $\text{UCC}(\alpha)$.

LEMMA 3.1. *Let $h(z) = 1 + \sum_{k=1}^{\infty} c_k z^k$ be subordinate to $H(z) = 1 + \sum_{k=1}^{\infty} C_k z^k$. If $H(z)$ is univalent in Δ and $H(\Delta)$ is convex, then $|c_n| \leq |C_1|$.*

THEOREM 3.2. *If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in \text{UCC}(\alpha)$, then*

$$|a_n| \leq (n-1)c + 1, \quad n \geq 2, \quad (3.1)$$

where $c = 4(1 - \alpha)/\pi^2$.

PROOF. Set

$$\Phi(z) = \frac{f'(z)}{g'(z)} = 1 + \sum_{k=1}^{\infty} c_k z^k \quad (3.2)$$

so that $\Phi(z) \prec q_\alpha(z)$, where $q_\alpha(z)$ is defined in (1.4).

Since $q_\alpha(z)$ is univalent and maps Δ onto a convex region, we may apply [Lemma 3.1](#).

Now

$$q_\alpha(z) = 1 + \frac{8(1-\alpha)}{\pi^2}z + \cdots, \quad \text{so that } |c_n| \leq \frac{8(1-\alpha)}{\pi^2}. \quad (3.3)$$

With $g(z) = z + \sum_{k=2}^{\infty} b_k z^k$, we compare the coefficients of z^n for the expansion of $\phi(z)$ to obtain

$$(n+1)|a_{n+1}| = c_n + \sum_{k=1}^{n-1} (k+1)b_{k+1}c_{n-k} + (n+1)b_{n+1}. \quad (3.4)$$

Since $g(z)$ is convex, it is well known that $|b_n| \leq 1$, $n = 1, 2, \dots$. From (3.4), we get

$$(n+1)|a_{n+1}| \leq cn(n+1) + (n+1), \quad (3.5)$$

and the proof is complete. $\square$

4. The class $\text{UQC}(\alpha)$. We now introduce a natural analogue to the class $\text{UCV}(\alpha)$ in terms of Alexander's result on convex functions [[1](#), page 43].

DEFINITION 4.1. A normalized function of the form (1.1) is said to be uniformly quasiconvex of order α , $-1 \leq \alpha < 1$, in Δ , denoted by $\text{UQC}(\alpha)$, if there exists a convex function $g(z)$ with $g(0) = 0$, $g'(0) = 1$, such that

$$\frac{(zf'(z))'}{g'(z)} < q_\alpha(z), \quad (4.1)$$

where $q_\alpha(z)$ is as defined by (1.4).

REMARK 4.2. (1) By setting $f(z) = g(z)$, we see that $\text{UCV}(\alpha) \subset \text{UQC}(\alpha)$.

(2) We see that $f \in \text{UQC}(\alpha)$ if and only if $zf' \in \text{UCC}(\alpha)$.

In view of the above remark, we obtain from [Theorem 1.3](#) a sufficient coefficient bound for inclusion in the family $\text{UQC}(\alpha)$.

THEOREM 4.3. If $\sum_{n=2}^{\infty} n^2 |a_n| \leq (1-\alpha)/2$, then $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in \text{UQC}(\alpha)$.

We next prove a theorem which shows that every function in $\text{UQC}(\alpha)$ is close to convex and hence univalent. We need a result due to Miller and Mocanu [[5](#)].

LEMMA 4.4. Let $M(z)$ and $N(z)$ be regular in Δ with $M(z) = N(z) = 0$ and let α be real. If $N(z)$ maps Δ onto a possibly many-sheeted region which is starlike with respect to the origin, then for $z \in \Delta$,

$$\operatorname{Re} \frac{M'(z)}{N'(z)} > \alpha \implies \operatorname{Re} \frac{M(z)}{N(z)} > \alpha. \quad (4.2)$$

THEOREM 4.5. *If $F(z) \in \text{UQC}(\alpha)$, then $F(z) \in K$ and hence it is univalent in Δ .*

PROOF. Since

$$\frac{(zf'(z))'}{g'(z)} < q_\alpha(z) \Rightarrow \operatorname{Re} \left\{ \frac{(zf'(z))'}{g'(z)} \right\} > \frac{1+\alpha}{2}, \quad (4.3)$$

an application of [Lemma 4.4](#), with $M(z) = zf'(z)$, $N(z) = g(z)$, proves the result. $\square$

THEOREM 4.6. *If $f(z) \in \text{UQC}(\alpha)$, then $H(z) = \int_0^z (tf'(t))' dt$ is in $\text{UCC}(\alpha)$.*

PROOF. If $f(z) \in \text{UQC}(\alpha)$, then there exists a function $g(z) \in C$ such that $(zf'(z))'/g'(z) < q_\alpha(z)$, where $q_\alpha(z)$ is as given by (1.4). The result now follows on observing that $H'(z) = (zf'(z))'$. $\square$

We close with coefficient estimates for the class $\text{UQC}(\alpha)$.

THEOREM 4.7. *If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in \text{UQC}(\alpha)$, then*

$$|a_n| \leq \frac{(n-1)c+1}{n}, \quad n \geq 2, \quad (4.4)$$

where $c = 4(1-\alpha)/\pi^2$.

PROOF. Proceeding on the same lines as in the proof of [Theorem 3.2](#), we obtain the result. $\square$

REMARK 4.8. When $\alpha = 0$, $\text{UQC}(0) = Q$ [6] and we see that the bounds are lower than the corresponding bounds for Q in [6].

REFERENCES

- [1] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, vol. 259, Springer-Verlag, New York, 1983.
- [2] A. W. Goodman, *On uniformly convex functions*, Ann. Polon. Math. **56** (1991), no. 1, 87-92.
- [3] ———, *On uniformly starlike functions*, J. Math. Anal. Appl. **155** (1991), no. 2, 364-370.
- [4] S. Kumar and C. Ramesha, *Subordination properties of uniformly convex and uniformly close to convex functions*, J. Ramanujan Math. Soc. **9** (1994), no. 2, 203-214.
- [5] S. S. Miller and P. T. Mocanu, *Second-order differential inequalities in the complex plane*, J. Math. Anal. Appl. **65** (1978), no. 2, 289-305.
- [6] ———, *On some classes of first-order differential subordinations*, Michigan Math. J. **32** (1985), no. 2, 185-195.
- [7] K. I. Noor and D. K. Thomas, *Quasi-convex univalent functions*, Int. J. Math. Math. Sci. **3** (1980), no. 2, 255-266.
- [8] S. Owa, *On convolution theorems*, Proc. Japan Acad. Ser. A Math. Sci. **65** (1989), no. 6, 161-162.
- [9] W. Rogosinski, *On the coefficients of subordinate functions*, Proc. London Math. Soc. (2) **48** (1943), 48-82.

- [10] F. Rønning, *On starlike functions associated with parabolic regions*, Ann. Univ. Mariae Curie-Skłodowska Sect. A **45** (1991), 117–122.
- [11] St. Ruscheweyh and T. Sheil-Small, *Hadamard products of Schlicht functions and the Pólya-Schoenberg conjecture*, Comment. Math. Helv. **48** (1973), 119–135.

K. G. Subramanian: Department of Mathematics, Madras Christian College, Tambaram, Chennai 600 059, India

T. V. Sudharsan: Department of Mathematics, South India Vaniar Educational Trust (SIVET) College, Gowrivakkam, Chennai 601 302, India

Herb Silverman: Department of Mathematics, University of Charleston, Charleston, SC 29424, USA

E-mail address: silvermanh@cofc.edu