ON COMMON FIXED POINTS OF PAIRS OF A SINGLE AND A MULTIVALUED COINCIDENTALLY COMMUTING MAPPINGS IN D -METRIC SPACES

B. C. DHAGE, A. JENNIFER ASHA, and S. M. KANG

Received 10 December 2002 and in revised form 15 February 2003

The present paper studies some common fixed-point theorems for pairs of a single-valued and a multivalued coincidentally commuting mappings in D -metric spaces satisfying a certain generalized contraction condition. Our result generalizes more than a dozen known fixed-point theorems in D -metric spaces including those of Dhage (2000) and Rhoades (1996).

2000 Mathematics Subject Classification: 47H10, 54H25.

1. Introduction. The concept of a D -metric space introduced by the first author in [1] is as follows. A nonempty set, together with a function $\rho : X \times X \times X \rightarrow [0, \infty)$, is called a D -metric space and denoted by (X, ρ) if the function ρ , called a D -metric on X , satisfies the following properties:

- (i) $\rho(x, y, z) = 0 \Leftrightarrow x = y = z$ (coincidence),
- (ii) $\rho(x, y, z) = 0 = \rho(p\{x, y, z\})$ (symmetry), where p is a permutation,
- (iii) $\rho(x, y, z) \leq \rho(x, y, a) + \rho(x, a, z) + \rho(a, y, z)$ for all $x, y, z, a \in X$ (tetrahedral inequality).

It is known that the D -metric ρ in a continuous function on X^3 in the topology of D -metric convergence is Hausdorff. The details of a D -metric space and its topological properties appear in Dhage [8]. Some specific examples of a D -metric space are presented in Dhage [2].

A sequence $\{x_n\} \subset X$ is called *convergent* and *converges* to a point x if $\lim_{m,n} \rho(x_m, x_n, x) = 0$. Again a sequence $\{x_n\} \subset X$ is called *D -Cauchy* if $\lim_{m,n,p} \rho(x_m, x_n, x_p) = 0$. A complete D -metric space X is one in which every D -Cauchy sequence converges to a point in X . A subset S of a D -metric space X is called *bounded* if there exists a constant $k > 0$ such that $\rho(x, y, z) \leq k$ for all $x, y, z \in X$ and the constant k is called a D -bound of S . The smallest among all such D -bounds k of S is called the *diameter* of X and it is denoted by $\delta(S)$.

Let 2^X and $CB(X)$ denote the classes of nonempty closed and nonempty, closed, bounded subsets of X , respectively. A correspondence $F : X \rightarrow 2^X$ is called a *multivalued mapping* on a D -metric space X , and a point $u \in X$ is called a *fixed point* of F if $u \in Fu$.

In [3], the first author has defined a notion of the generalized or Kasusai D -metric on X . Let $\kappa : (\text{CB}(X))^3 \rightarrow [0, \infty)$ be a function defined by

$$\kappa(A, B, C) = \inf \{ \epsilon > 0 \mid A \cup B \subset N(c, \epsilon), B \cup C \subset N(a, \epsilon), C \cup A \subset N(b, \epsilon) \}, \quad (1.1)$$

where $N(A, \epsilon) = \cup_{a \in A} N(a, \epsilon)$, $N(a, \epsilon) = \{x \in N^*(a, \epsilon) \mid \rho(a, x, y) < \epsilon \text{ for all } y \in N^*(a, \epsilon)\}$, and $N^*(a, \epsilon) = \{x \in X \mid \rho(a, x, x) < \epsilon\}$.

The definition (1.1) is equivalent to

$$\kappa(A, B, C) = \max \left\{ \sup_{a \in A, b \in B} D(a, b, c), \sup_{b \in B, c \in C} D(b, c, a), \sup_{c \in C, a \in A} D(c, a, b) \right\}, \quad (1.2)$$

where $D(a, b, c) = \inf \{ \rho(a, b, c) \mid c \in C \}$.

Define

$$\begin{aligned} D(A, B, C) &= \inf \{ \rho(a, b, c) \mid a \in A, b \in B, c \in C \}, \\ \delta(A, B, C) &= \sup \{ \rho(a, b, c) \mid a \in A, b \in B, c \in C \}. \end{aligned} \quad (1.3)$$

Notice that D and δ are continuous functions on $(\text{CB}(X))^3$ and satisfy

$$D(A, B, C) \leq \kappa(A, B, C) \leq \delta(A, B, C). \quad (1.4)$$

A multivalued mapping $F : X \rightarrow \text{CB}(X)$ is called *continuous* if

$$\lim_{m, n} \rho(x_m, x_n, x) = 0 \implies \kappa(Fx_m, Fx_n, Fx) = 0. \quad (1.5)$$

In [3], the first author has proved some fixed-point theorem for multivalued contraction mappings in D -metric spaces, and in [5] he has proved some common fixed-point theorems for coincidentally commuting single-valued mappings in D -metric spaces satisfying a condition of generalized contraction.

In this paper, we prove some common fixed-point theorems for a pair of singlevalued and multivalued mappings in a D -metric space satisfying a contraction condition more general than that given in Dhage [1, 2, 3, 4, 5, 7] and Rhoades [12]. The results of this paper are new to the fixed-point theory in D -metric spaces and include nearly a dozen of known fixed-point theorems as special cases (see [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12]).

2. Preliminaries. Before going to the main results of this paper, we give some preliminaries needed in the sequel.

Let $F : X \rightarrow 2^X$. Then by an orbit of F at a point $x \in X$ we mean a set $O_F(x)$ in X defined by

$$O_F(x) = \{x_0 = x, x_{n+1} \in Fx_n, n \geq 0\}. \quad (2.1)$$

An orbit $O_F(x)$ is called *bounded* if $\delta(O_F(x)) < \infty$, and a D -metric space X is called *F-orbitally bounded* if $O_F(x)$ is bounded for each $x \in X$. Again an F -orbit $O_F(x)$ is called *complete* if every D -Cauchy sequence in $O_F(x)$ converges to a point in X . A D -metric space X is said to be *F-orbitally complete* if $O_F(x)$ is complete for each $x \in X$. Finally, F is called *F-orbitally continuous* if for any sequence $\{x_n\} \subseteq O_F(x)$, we have

$$\lim_{m,n} \rho(x_m, x_n, x^*) = 0 \Rightarrow \lim_{m,n} \kappa(Fx_m, Fx_n, Fx^*) = 0 \quad (2.2)$$

for each $x \in X$.

Let Φ denote the class of all functions $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfying the following properties:

- (i) ϕ is continuous,
- (ii) ϕ is nondecreasing,
- (iii) $\phi(t) < t$, $t > 0$,
- (iv) $\sum_{n=1}^{\infty} \phi^n(t) < \infty$ for all $t \in [0, \infty)$.

The function ϕ is called a *Lipschitz control function* or *Lipschitz growth function* and the usual growth function is $\phi(t) = \alpha t$, $0 \leq t < 1$. The following lemma concerning the function ϕ appears in [7].

LEMMA 2.1. *If $\phi \in \Phi$, then $\phi^n(t) = 0$ for each $n \in \mathbb{N}$ and $\lim_n \phi^n(t) = 0$ for each $t \in [0, \infty)$.*

We need the following D -Cauchy principle of Dhage [7] in the sequel.

LEMMA 2.2 (D -Cauchy principle). *Let $\{x_n\}$ be a bounded sequence in a D -metric space X with D -bound k satisfying, for some positive real number r ,*

$$\rho(x_n, x_{n+1}, x_m) \leq [\phi^n(k^r)]^{1/r} \quad (2.3)$$

for all $m > n \in \mathbb{N}$, where $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfies $\sum_{n=1}^{\infty} \phi^n(t) < \infty$ for each $t \in [0, \infty)$. Then $\{x_n\}$ is a D -Cauchy sequence in X .

PROOF. The proof appears in [7], but for the sake of completeness we give the details. Let $p, t \in \mathbb{N}$ be arbitrary but fixed. Then from (2.3) it follows that

$$\begin{aligned} \rho(x_n, x_{n+1}, x_{n+p}) &\leq [\phi^n(k^r)]^{1/r}, \\ \rho(x_n, x_{n+1}, x_{n+p+t}) &\leq [\phi^n(k^r)]^{1/r}, \end{aligned} \quad (2.4)$$

for all $n \in \mathbb{N}$.

Now by repeated application of the tetrahedral inequality, we obtain

$$\begin{aligned} &\rho(x_n, x_{n+p}, x_{n+p+t}) \\ &\leq \rho(x_n, x_{n+1}, x_{n+p}) + \rho(x_n, x_{n+1}, x_{n+p+t}) + \rho(x_{n+1}, x_{n+p}, x_{n+p+t}) \\ &\leq \rho(x_n, x_{n+1}, x_{n+p}) + \rho(x_n, x_{n+1}, x_{n+p+t}) + \rho(x_{n+1}, x_{n+2}, x_{n+p}) \\ &\quad + \rho(x_{n+1}, x_{n+2}, x_{n+p+t}) + \rho(x_{n+2}, x_{n+p}, x_{n+p+t}) \end{aligned}$$

$$\begin{aligned}
&\leq 2[\phi^n(k^r)]^{1/r} + 2[\phi^{n+1}(k^r)]^{1/r} + \rho(x_{n+2}, x_{n+p}, x_{n+p+t}) \\
&\leq 2\{[\phi^n(k^r)]^{1/r} + \dots + [\phi^{n+p-2}(k^r)]^{1/r}\} + \rho(x_{n+p-1}, x_{n+p}, x_{n+p+t}) \\
&\leq 2 \sum_{j=n}^{n+p-1} [\phi^j(k^r)]^{1/r}.
\end{aligned} \tag{2.5}$$

Since $\sum_{n=1}^{\infty} \phi^n(t) < \infty$ for each $t \in [0, \infty)$, we have $\sum_{j=1}^{\infty} [\phi^j(k^r)]^{1/r} < \infty$ and so $\lim_n \sum_{j=n}^{n+p-1} [\phi^j(k^r)]^{1/r} = 0$. Now from (2.5) it follows that

$$\lim_{n \rightarrow \infty} \rho(x_n, x_{n+p}, x_{n+p+t}) = 0. \tag{2.6}$$

This proves that $\{x_n\}$ is a D -Cauchy sequence in X and the proof of the lemma is complete. $\square$

As a direct application of Lemma 2.2, we obtain the following result proved in [5].

LEMMA 2.3. *Let $\{x_n\}$ be a bounded sequence in a D -metric space X with D -bound k satisfying*

$$\rho(x_n, x_{n+1}, x_m) \leq \lambda^n k \tag{2.7}$$

for all $m > n \in \mathbb{N}$, where $0 \leq \lambda < 1$. Then $\{x_n\}$ is D -Cauchy.

We use contractive conditions of the form

$$a^r \leq \phi(b^r) \tag{2.8}$$

for some positive real number r , where a and b are nonnegative real numbers and $\phi \in \Phi$, because sometimes inequality (2.8) holds, but for the same real numbers a and b , the inequality

$$a \leq \phi(b) \tag{2.9}$$

does not hold. To see this, let $\phi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a function defined by

$$\phi(t) = \frac{\alpha t}{1+t}, \quad 0 \leq \alpha < 1. \tag{2.10}$$

Obviously the function ϕ is continuous, nondecreasing and satisfies $\phi(t) = \alpha t / (1+t) < t$ for $t > 0$. Again since

$$\sum_{n=1}^{\infty} \phi^n(t) = \sum_{n=1}^{\infty} \frac{\alpha^n t}{1+t+\dots+\alpha^{n-1}t} < \sum_{n=1}^{\infty} \alpha^n < \infty, \tag{2.11}$$

we have that $\phi \in \Phi$.

Now for $a = 1/2$ and $b = 2/3$, we have, by (2.9),

$$\frac{1}{2} \leq \phi\left(\frac{2}{3}\right) = \frac{(2/3)\alpha}{1+2/3} = \frac{2}{5}\alpha, \quad (2.12)$$

which is not true since $0 \leq \alpha < 1$. But for the same values of a and b , we have a positive real number $r = 2$ such that

$$\left(\frac{1}{2}\right)^2 = \frac{1}{4} \leq \frac{4\alpha}{13} = \phi\left(\left(\frac{2}{3}\right)^2\right) \quad (2.13)$$

for $13/16 \leq \alpha < 1$. Hence inequality (2.8) holds. Thus inequality (2.9) does not imply inequality (2.8). Actually, inequalities (2.8) and (2.9) are independent. To show that inequality (2.8) does not imply inequality (2.9), let $a = 1/4$, $b = 4/9$, and $r = 1/2$. Clearly, inequality (2.8) does not hold, but for the same values of a , b , and r , one has

$$\frac{1}{4} \leq 4\frac{\alpha}{13} = \frac{\alpha(4/9)}{1+4/9} = \phi\left(\frac{4}{9}\right) \quad (2.14)$$

for $\alpha \geq 13/16$, and so inequality (2.9) holds. Thus inequalities (2.8) and (2.9) are independent.

In the following sections, we will prove the main results of this paper.

3. Weak commuting mappings in D -metric spaces. Let $F : X \rightarrow 2^X$ and $g : X \rightarrow X$. Then the pair $\{F, g\}$ of maps is called *limit coincident* if $\lim_n Fx_n = \{\lim_n gx_n\}$ for some sequence $\{x_n\}$ in X , and *coincident* if there exists a point $u \in X$ such that $Fu = \{gu\}$. Again two maps F and g are called *limit commuting* if $\lim_n Fgx_n = \{\lim_n gFx_n\}$, where $\{x_n\}$ is a sequence in X , and *commuting* if $Fgx = \{gFx\}$ for all $x \in X$. Two maps F and g are called *limit coincidentally commuting* if their limit coincidence implies the limit commutativity on X . Similarly, they are called *coincidentally commuting* if they are commuting at the coincidence points. Again two maps F and g are said to be *limit pseudocommuting* if $\lim_n Fgx_n \cap \lim_n gFx_n \neq \phi$, that is, $\lim_n D(Fgx_n, gFx_n, gFx_n) = 0$, where $\{x_n\}$ is a sequence in X , and *pseudocommuting* if $Fgx \cap gFx \neq \emptyset$ for each $x \in X$. Finally, the pair $\{F, g\}$ is called *limit coincidentally pseudocommuting* if its limit coincidence implies the limit pseudocommutativity on X , and *coincidentally pseudocommuting* if it is pseudocommuting at the coincidence points. It is known that a coincidentally commuting pair is limit coincidentally commuting and a coincidentally pseudocommuting pair is limit coincidentally pseudocommuting, but the converse implications need not hold. A pair of maps $\{F, g\}$ is *weak commuting* if it is either limit commuting, coincidentally commuting, limit pseudocommuting, or coincidentally pseudocommuting on X . Below, we will prove some common fixed-point theorems for each of these weak commuting mappings on D -metric spaces.

3.1. Limit coincidentally commuting maps in D -metric spaces. Let $F : X \rightarrow 2^X$ and $g : X \rightarrow X$. By an (F/g) -orbit of the pair $\{F, g\}$ of maps at a point $x \in X$, we mean a set $O_F(gx)$ in X defined by

$$O_F(gx) = \{y_n \mid y_0 = gx_0, y_n = gx_n \in Fx_{n-1}, n \in \mathbb{N}, \text{ where } x_0 = x\} \quad (3.1)$$

for some sequence $\{x_n\}$ in X . The orbit $O_F(gx)$ is well defined for each $x \in X$ if $F(X) \subseteq g(X)$. By $\overline{O_F(gx)}$ we denote the closure of the set $O_F(gx)$ in X .

A D -metric space X is called (F/g) -orbitally bounded if $\delta(O_F(gx)) < \infty$ for each $x \in X$. Further X is called (F/g) -orbitally complete if every D -Cauchy sequence $\{x_n\} \subset O_F(gx)$ converges to a point in X for each $x \in X$. Finally, a mapping $T : X \rightarrow CB(X)$ is called (F/g) -orbitally continuous if for any $\{x_n\} \subset O_F(gx)$, $x_n \rightarrow x^*$ implies that $Tx_n \rightarrow Tx^*$ for each $x \in X$.

THEOREM 3.1. Let $F : X \rightarrow CB(X)$ and $g : X \rightarrow X$ be two mappings satisfying, for some positive real number r ,

$$\begin{aligned} \delta^r(Fx, Fy, Fz) \\ \leq \phi(\max\{\rho^r(gx, gy, gz), \delta^r(Fx, Fy, gz), \delta^r(gx, Fx, gz), \\ \delta^r(gy, Fy, gz), \delta^r(gx, Fy, gz), \delta^r(gy, Fx, gz)\}) \end{aligned} \quad (3.2)$$

for all $x, y, z \in X$, where $\phi \in \Phi$. Suppose that

- (a) $F(X) \subseteq g(X)$ and $g(X)$ is bounded,
- (b) $\{F, g\}$ is limit coincidentally commuting,
- (c) F or g is (F/g) -orbitally continuous.

Further if X is (F/g) -orbitally complete D -metric space, then F and g have a unique common fixed point $u \in X$ such that $Fu = \{u\} = gu$. Moreover, if g is continuous at u , then F is also continuous at u in the Kasubai D -metric on X .

PROOF. Let $x \in X$ be arbitrary and define a sequence $\{y_n\}$ in X as follows. Take $x_0 = x$ and $y_0 = gx_0$. Choose a point $y_1 \in Fx_0 = X_1$. Since $F(X) \subseteq g(X)$, there is a point $x_1 \in X$ such that $y_1 = gx_1$. Again choose a point $y_2 \in Fx_1 = X_2$. By hypothesis (a), there is a point $x_2 \in X$ such that $y_2 = gx_2$. Proceeding in this way, by induction there is a sequence $\{x_n\}$ of points in X such that

$$y_0 = gx_0, \quad y_{n+1} = gx_{n+1} \in X_{n+1} = Fx_n, \quad n = 0, 1, 2, \dots \quad (3.3)$$

From hypothesis (a), it follows that

$$\delta(X_m, X_n, X_p) \leq \delta(g(X)) = k < \infty \quad (3.4)$$

for all $m, n, p \in \mathbb{N}$.

Now there are two cases.

CASE 1. Suppose that $y_r = y_{r+1}$ for some $r \in \mathbb{N}$. Then we have $gx_r = gx_{r+1} = u$ for some $u \in X$.

We will show that $Fx_r = \{u\}$. Suppose not. Then by (3.2),

$$\begin{aligned}
 \delta^r(Fx_r, Fx_r, u) &= \delta^r(Fx_r, Fx_r, gx_{r+1}) \\
 &\leq \delta^r(Fx_r, Fx_r, Fx_r) \\
 &\leq \phi(\max\{\rho^r(gx_r, gx_r, gx_r), \delta^r(gx_r, Fx_r, gx_r), \delta^r(Fx_r, Fx_r, gx_r)\}) \quad (3.5) \\
 &\leq \phi(\max\{0, \delta^r(gx_r, Fx_r, gx_r), \delta^r(Fx_r, Fx_r, u)\}) \\
 &= \phi(\max\{\delta^r(u, Fx_r, u), \delta^r(Fx_r, Fx_r, u)\}) \\
 &= \phi(\delta^r(u, Fx_r, u))
 \end{aligned}$$

because $\delta^r(Fx_r, Fx_r, u) \leq \phi(\delta^r(Fx_r, Fx_r, u))$ is not possible in view of $\phi \in \Phi$.

Again by (3.2),

$$\begin{aligned}
 \delta^r(Fx_r, u, u) &= \delta^r(Fx_r, gx_{r+1}, gx_{r+1}) \\
 &\leq \delta^r(Fx_r, Fx_r, Fx_r) \\
 &\leq \phi(\max\{\delta^r(u, Fx_r, u), \delta^r(Fx_r, Fx_r, u)\}) \quad (3.6) \\
 &= \phi(\delta^r(Fx_r, Fx_r, u)).
 \end{aligned}$$

Substituting (3.6) in (3.5), we obtain

$$\delta^r(Fx_r, Fx_r, u) \leq \phi^2(\delta^r(Fx_r, Fx_r, u)), \quad (3.7)$$

which is a contradiction since $\phi \in \Phi$. Hence $Fx_r = u$. Since F and g are limit coincidentally commuting, one has $Fgx_r = \{gFx_r\}$.

We will show that u is a common fixed point of F and g such that $Fu = \{u\} = gu$.

Now,

$$\begin{aligned}
 \delta^r(Fu, gu, u) &= \delta^r(FFx_r, Fgx_r, Fx_r) \\
 &\leq \phi(\max\{\rho^r(gFx_r, ggx_r, gx_r), \delta^r(FFx_r, Fgx_r, gx_r), \\
 &\quad \delta^r(gFx_r, FFx_r, gx_r), \delta^r(ggx_r, Fgx_r, gx_r), \\
 &\quad \delta^r(gFx_r, Fgx_r, gx_r), \delta^r(ggx_r, FFx_r, gx_r)\}) \quad (3.8) \\
 &= \phi(\max\{\rho^r(gFx_r, ggx_r, gx_r), \delta^r(ggx_r, FFx_r, gx_r)\}) \\
 &= \phi(\delta^r(Fu, gu, u)),
 \end{aligned}$$

which is possible only when $Fu = \{u\} = gu$ since $\phi \in \Phi$.

CASE 2. Assume that $y_n \neq y_{n+1}$ for each $n \in \mathbb{N}$. We will show that $\{y_n\}$ is a D -Cauchy sequence in X . Let $x = x_0$, $y = x_1$, and $z = x_{m-1}$, $m \geq 1$. Then by (3.2),

$$\begin{aligned}
& \rho^r(\mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_m) \\
& \leq \delta^r(Fx_0, Fx_1, Fx_{m-1}) \\
& \leq \phi(\max\{\rho^r(gx_0, gx_1, gx_{m-1}), \delta^r(Fx_0, Fx_1, gx_{m-1}), \delta^r(gx_0, Fx_0, gx_{m-1}), \\
& \quad \delta^r(gx_1, Fx_1, gx_{m-1}), \delta^r(gx_0, Fx_1, gx_{m-1}), \delta^r(gx_1, Fx_0, gx_{m-1})\}) \\
& \leq \phi(\max\{\delta^r(X_0, X_1, X_{m-1}), \delta^r(X_1, X_2, X_{m-1}), \delta^r(X_0, X_1, X_{m-1}), \\
& \quad \delta^r(X_1, X_2, X_{m-1}), \delta^r(X_0, X_2, X_{m-1}), \delta^r(X_1, X_1, X_{m-1})\}) \\
& \leq \phi\left(\max_{0 \leq a \leq 1, 1 \leq b \leq 2} \delta^r(X_a, X_b, X_{m-1})\right) \\
& \leq \phi(k^r),
\end{aligned} \tag{3.9}$$

that is,

$$\rho(\mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_m) \leq [\phi(k^r)]^{1/r}. \tag{3.10}$$

Similarly, letting $x = x_1$, $y = x_2$, and $z = z_{m-1}$, $m \geq 2$ in (3.2), we obtain

$$\begin{aligned}
& \rho^r(\mathcal{Y}_2, \mathcal{Y}_3, \mathcal{Y}_m) \\
& \leq \delta^r(Fx_1, Fx_2, Fx_{m-1}) \\
& \leq \phi(\max\{\rho^r(gx_1, gx_2, gx_{m-1}), \delta^r(Fx_1, Fx_2, gx_{m-1}), \\
& \quad \delta^r(gx_1, Fx_1, gx_{m-1}), \delta^r(gx_2, Fx_2, gx_{m-1}), \\
& \quad \delta^r(gx_1, Fx_2, gx_{m-1}), \delta^r(gx_2, Fx_1, gx_{m-1})\}) \\
& \leq \phi(\max\{\delta^r(Fx_0, Fx_1, Fx_{m-2}), \delta^r(Fx_1, Fx_2, Fx_{m-2}), \\
& \quad \delta^r(Fx_0, Fx_1, Fx_{m-2}), \delta^r(Fx_1, Fx_2, Fx_{m-2}), \\
& \quad \delta^r(Fx_0, Fx_2, Fx_{m-2}), \delta^r(Fx_1, Fx_1, Fx_{m-2})\}) \\
& \leq \phi\left(\phi\left(\max_{0 \leq a \leq 2, 1 \leq b \leq 3} \delta^r(X_a, X_b, X_{m-2})\right)\right) \\
& \leq \phi(\phi(k^r)) \\
& = \phi^2(k^r),
\end{aligned} \tag{3.11}$$

that is,

$$\rho(\mathcal{Y}_2, \mathcal{Y}_3, \mathcal{Y}_m) \leq [\phi^2(k^r)]^{1/r}. \tag{3.12}$$

In general, by induction,

$$\rho(\mathcal{Y}_n, \mathcal{Y}_{n+1}, \mathcal{Y}_m) \leq [\phi^n(k^r)]^{1/r} \tag{3.13}$$

for all $m > n \in \mathbb{N}$.

Hence, the application of [Lemma 2.2](#) yields that $\{\mathcal{Y}_n\}$ is a D -Cauchy sequence in X . The D -metric space X being complete, there is a point $u \in X$ such that $\lim_n \mathcal{Y}_n = u$. The definition of $\{\mathcal{Y}_n\}$ implies that $\lim_n gx_n = u$. We will show that $\lim_n Fx_n = \{u\}$.

Now,

$$\begin{aligned}
 \lim_n \delta^r(Fx_n, Fx_n, u) &= \lim_n \delta^r(Fx_n, Fx_n, y_{n+1}) \\
 &\leq \lim_n \delta^r(Fx_n, Fx_n, Fx_n) \\
 &\leq \lim_n \phi(\max\{\rho^r(gx_n, gx_n, gx_n), \delta^r(Fx_n, Fx_n, gx_n), \delta^r(gx_n, Fx_n, gx_n)\}) \\
 &= \lim_n \phi(\max\{\delta^r(Fx_n, Fx_n, u), 0\}) \\
 &= \phi\left(\lim_n \delta^r(Fx_n, Fx_n, u)\right),
 \end{aligned} \tag{3.14}$$

which implies that $\lim_n Fx_n = u$. Thus we have

$$\lim_n Fx_n = \{u\} = \lim_n gx_n. \tag{3.15}$$

Since F and g are limit coincidentally commuting, one has

$$\lim_n Fgx_n = \left\{ \lim_n gFx_n \right\}. \tag{3.16}$$

Suppose that g is (F/g) -orbitally continuous on X . Then we have

$$\lim_n Fgx_n = \lim_n gFx_n = \lim_n ggx_n = gu. \tag{3.17}$$

First, we will show that u is a common fixed point of F and g . Suppose not. Then we have

$$\begin{aligned}
 \delta^r(u, u, gu) &= \lim_n \delta^r(Fx_n, Fx_n, gFx_n) \\
 &= \lim_n \delta^r(Fx_n, Fx_n, Fgx_n) \\
 &\leq \lim_n \phi(\max\{\rho^r(gx_n, gx_n, ggx_n), \\
 &\quad \delta^r(Fx_n, Fx_n, ggx_n), \delta^r(gx_n, Fx_n, ggx_n)\}) \\
 &= \phi\left(\max\left\{\lim_n \delta^r(gx_n, gx_n, ggx_n), \lim_n \delta^r(Fx_n, Fx_n, ggx_n)\right\}\right) \\
 &= \phi(\delta^r(u, u, gu)),
 \end{aligned} \tag{3.18}$$

which is a contradiction and hence $gu = u$.

Again

$$\begin{aligned}
 \delta^r(Fu, gu, u) &= \lim_n \delta^r(Fu, Fx_n, Fgx_n) \\
 &\leq \lim_n \phi(\max\{\rho^r(gu, gx_n, ggx_n), \delta^r(Fu, Fx_n, ggx_n), \delta^r(gu, Fu, ggx_n), \\
 &\quad \delta^r(gx_n, Fx_n, ggx_n), \delta^r(gu, Fx_n, ggx_n), \delta^r(gx_n, Fu, ggx_n)\})
 \end{aligned}$$

$$\begin{aligned}
&= \phi(\max\{\rho^r(gu, u, gu), \delta^r(Fu, u, gu), \delta^r(gu, Fu, gu), \\
&\quad \delta^r(u, u, gu), \delta^r(gu, u, gu), \delta^r(u, Fu, gu)\}) \\
&= \phi(\delta^r(Fu, gu, u)),
\end{aligned} \tag{3.19}$$

which is possible only when $Fu = \{u\} = gu$ since $\phi \in \Phi$. Thus u is a common fixed point of F and g .

Next, suppose that F is (F/g) -orbitally continuous on X . Then we have

$$\lim_n Fgx_n = \lim_n gFx_n = \lim_n FFx_n = Fu = \{z\}. \tag{3.20}$$

We will show that z is a common fixed point of F and g . Since $F(X) \subseteq g(X)$, there is a point $v \in X$ such that $Fu = gv = z$. We will show that $Fv = gv = \{z\}$. By (3.2),

$$\begin{aligned}
&\delta^r(Fv, gv, Fv) \\
&= \lim_n \delta^r(Fv, Fv, FFx_n) \\
&\leq \lim_n \phi(\max\{\rho^r(gv, gv, gFx_n), \delta^r(Fv, gv, gFx_n), \delta^r(gv, Fv, gFx_n)\}) \\
&= \phi(\max\{\delta^r(gv, gv, gv), \delta^r(Fv, gv, z)\}),
\end{aligned} \tag{3.21}$$

that is,

$$\delta^r(Fv, gv, z) \leq \phi(\delta^r(Fv, gv, z)), \tag{3.22}$$

which implies that $Fv = gv = \{z\}$ since $\phi \in \Phi$.

Since F and g are limit coincidentally commuting, they are coincidentally commuting on X . Therefore, we have $Fgv = gFv$. Now, proceeding with the arguments as in Case 1, it is proved that z is a common fixed point of F and g .

To prove the uniqueness, let z^* ($\neq z$) be another common fixed point of F and g . Then by (3.2),

$$\begin{aligned}
\rho^r(z, z, z^*) &= \delta^r(Fz, Fz, Fz^*) \\
&\leq \phi(\max\{\rho^r(gz, gz, gz^*), \delta^r(Fz, Fz, gz^*), \\
&\quad \delta^r(gz, Fz, gz^*), \delta^r(gz, Fz, gz^*)\}) \\
&= \phi(\rho^r(z, z, z^*)),
\end{aligned} \tag{3.23}$$

which is a contradiction. Hence $z = z^*$. Then F and g have a unique common fixed point $z \in X$ with $Fz = \{z\} = gz$.

Finally, suppose that g is continuous at the common fixed point z of F and g . Then we will prove that F is also continuous at z . Let $\{z_n\}$ be any sequence

in X converging to the common fixed point z . Since g is continuous on X , we have

$$\lim_{m,n} \rho(z_m, z_n, z) = 0 \Rightarrow \lim_{m,n} \rho(gz_m, gz_n, gz) = 0. \quad (3.24)$$

From (1.2), it follows that

$$\kappa(Fz_m, Fz_n, Fz) \leq \delta(Fz_m, Fz_n, Fz). \quad (3.25)$$

Now,

$$\begin{aligned} & \delta^r(Fz_m, Fz_n, Fz) \\ & \leq \phi(\max\{\rho^r(gz_m, gz_n, gz), \delta^r(Fz_m, Fz_n, gz), \delta^r(gz_m, Fz_m, gz), \\ & \quad \delta^r(gz_n, Fz_n, gz), \delta^r(gz_m, Fz_n, gz), \delta^r(gz_n, Fz_m, gz)\}). \end{aligned} \quad (3.26)$$

Therefore,

$$\begin{aligned} & \lim_{m,n} \delta^r(Fz_m, Fz_n, Fz) \\ & \leq \lim_{m,n} \phi(\max\{\rho^r(gz_m, gz_n, gz), \delta^r(Fz_m, Fz_n, Fz), \delta^r(gz_m, Fz_m, z), \\ & \quad \delta^r(gz_n, Fz_n, z), \delta^r(gz_m, Fz_n, z), \delta^r(gz_n, Fz_m, z)\}) \\ & = \phi(\max\{0, \lim_{m,n} \delta^r(Fz_m, Fz_n, Fz), \lim_m \delta^r(z, Fz_m, z), \lim_n \delta^r(z, Fz_n, z)\}) \\ & = \phi(\max\{\lim_m \delta^r(z, Fz_m, z), \lim_n \delta^r(z, Fz_n, z)\}). \end{aligned} \quad (3.27)$$

But

$$\begin{aligned} & \lim_m \delta^r(z, Fz_m, z) \\ & = \lim_m \delta^r(Fz, Fz, Fz_m) \\ & \leq \lim_m \phi(\max\{\rho^r(gz, gz, gz_m), \delta^r(Fz, Fz, gz_m), \delta^r(gz, Fz, gz_m)\}) \\ & = \phi(\max\{0, 0, 0\}) \\ & = 0. \end{aligned} \quad (3.28)$$

Similarly, $\lim_n \delta^r(z, Fz_n, z) = 0$. Substituting these estimates in (3.27) yields that

$$\lim_{m,n} \delta^r(Fz_m, Fz_n, Fz) = 0 \quad (3.29)$$

or

$$\lim_{m,n} \delta(Fz_m, Fz_n, Fz) = 0. \quad (3.30)$$

Now from (3.25), it follows that

$$\lim_{m,n} \kappa(Fz_m, Fz_n, Fz) = 0, \quad (3.31)$$

and so F is continuous at the common fixed point z of F and g . This completes the proof. $\square$

Letting $g = I$, the identity map on X and $r = 1$, in [Theorem 3.1](#), we obtain the following corollary.

COROLLARY 3.2. *Let $F : X \rightarrow \text{CB}(X)$ be a multivalued mapping satisfying*

$$\begin{aligned} \delta(Fx, Fy, Fz) \leq \phi(\rho(x, y, z), \delta(Fx, Fy, z), \delta(x, Fx, z), \\ \delta(y, Fy, z), \delta(x, Fy, z), \delta(y, Fx, z)) \end{aligned} \quad (3.32)$$

for all $x, y, z \in X$, where $\phi \in \Phi$. Further if X is F -orbitally bounded and F -orbitally complete D -metric space, then F has a unique fixed point $u \in X$ such that $Fu = \{u\}$ and F is continuous at u .

COROLLARY 3.3. *Let $F : X \rightarrow \text{CB}(X)$ be a multivalued mapping satisfying*

$$\begin{aligned} \delta(Fx, Fy, Fz) \leq \lambda \max \{ \rho(x, y, z), \delta(Fx, Fy, z), \delta(x, Fx, z), \\ \delta(y, Fy, z), \delta(x, Fy, z), \delta(y, Fx, z) \} \end{aligned} \quad (3.33)$$

for all $x, y, z \in X$, where $0 \leq \lambda < 1$. Further if X is F -orbitally bounded and F -orbitally complete D -metric space, then F has a unique fixed point $u \in X$ such that $Fu = \{u\}$ and F is continuous at u .

[Corollary 3.3](#) includes the following fixed point of Dhage [\[3\]](#) as a special case.

COROLLARY 3.4 (see [\[3\]](#)). *Let X be a bounded and complete D -metric space and let $F : X \rightarrow \text{CB}(X)$ be a multivalued mapping satisfying*

$$\delta(Fx, Fy, Fz) \leq \lambda \rho(x, y, z) \quad (3.34)$$

for all $x, y, z \in X$, where $0 \leq \lambda < 1$. Then F has a unique fixed point $u \in X$ such that $Fu = \{u\}$ and F is continuous at u .

COROLLARY 3.5. *Let $f, g : X \rightarrow X$ be two mappings satisfying*

$$\begin{aligned} \rho^r(fx, fy, fz) \\ \leq \phi(\max \{ \rho^r(gx, gy, gz), \rho^r(fx, fy, gz), \rho^r(gx, fx, gz), \\ \rho^r(gy, fy, gz), \rho^r(gx, fy, gz), \rho^r(gy, fx, gz) \}) \end{aligned} \quad (3.35)$$

for all $x, y, z \in X$, where $\phi \in \Phi$. Suppose that

- (a) $f(X) \subseteq g(X)$,
- (b) $\{f, g\}$ is limit coincidentally commuting,
- (c) f or g is continuous.

Further if X is (f/g) -orbitally bounded and (f/g) -orbitally complete D -metric space, then f and g have a unique common fixed point $u \in X$. Moreover, if g is continuous at u , then f is also continuous at u .

REMARK 3.6. Note that [Corollary 3.5](#) includes the class of pairs of fixed-point mappings of Dhage [7] characterized by the inequality

$$\begin{aligned} & \rho^r(fx, fy, fz) \\ & \leq \phi(\max\{\rho^r(gx, gy, gz), \rho^r(gx, fx, gz), \\ & \quad \rho^r(gy, fy, gz), \rho^r(gx, fy, gz), \rho^r(gy, fx, gz)\}) \end{aligned} \quad (3.36)$$

for all $x, y, z \in X$ and $\phi \in \Phi$.

COROLLARY 3.7. Let $f, g : X \rightarrow X$ be two mappings satisfying for some positive real numbers p, q , and r ,

$$\begin{aligned} & \rho^r(f^p x, f^p y, f^p z) \\ & \leq \phi(\max\{\rho^r(g^q x, g^q y, g^q z), \rho^r(f^p x, f^p y, g^q z), \\ & \quad \rho^r(g^q x, f^p x, g^q z), \rho^r(g^q y, f^p y, g^q z), \\ & \quad \rho^r(g^q x, f^p y, g^q z), \rho^r(g^q y, f^p x, g^q z)\}) \end{aligned} \quad (3.37)$$

for all $x, y, z \in X$, where $\phi \in \Phi$. Suppose that

- (a) $f^p(X) \subseteq g^q(X)$,
- (b) $\{f, g\}$ is commuting,
- (c) f or g is continuous.

Further if X is an (f^p/g^q) -orbitally bounded and (f^p/g^q) -orbitally complete D -metric space, then f and g have a unique common fixed point $u \in X$. Moreover, if g is continuous at u , then f^p is also continuous at u .

PROOF. Let $S = f^p$ and $T = g^q$. Then by [Corollary 3.5](#), S and T have a unique common fixed point $u \in X$, that is, $Su = f^p u = u = g^q u = Tu$. Now by commutativity of f and g , we obtain

$$fu = f(f^p u) = f^p(fu), \quad fu = f(g^q u) = g^q(fu). \quad (3.38)$$

This shows that fu is again a common fixed point of f^p and g^q . By the uniqueness of u , we have $fu = u$. Similarly it is proved that $gu = u$. Thus f and g have a unique common fixed point $u \in X$. Further if g is continuous on X , g^q is continuous on X and by application of [Corollary 3.5](#) yields that f^p is continuous at u . This completes the proof. $\square$

[Corollary 3.7](#) includes the class of pairs of fixed-point mappings of Dhage [7] characterized by the inequality

$$\begin{aligned} & \rho^r(f^p x, f^p y, f^p z) \\ & \leq \phi(\max\{\rho^r(g^q x, g^q y, g^q z), \rho^r(g^q x, f^p x, g^q z), \\ & \quad \rho^r(g^q y, f^p y, g^q z), \rho^r(g^q x, f^p y, g^q z), \rho^r(g^q y, f^p x, g^q z)\}) \end{aligned} \quad (3.39)$$

for all $x, y, z \in X$ and $\phi \in \Phi$.

COROLLARY 3.8. *Let f be a self-map of a D -metric space X satisfying*

$$\rho(fx, fy, fz) \leq \lambda \max \{ \rho(x, y, z), \rho(fx, fy, z), \rho(x, fx, z), \rho(y, fy, z), \rho(x, fy, z), \rho(y, fx, z) \} \quad (3.40)$$

for all $x, y, z \in X$, where $0 \leq \lambda < 1$. Further if X is f -orbitally bounded and f -orbitally complete, then f has a unique fixed point $u \in X$ and f is continuous at u .

COROLLARY 3.9. *Let f be a self-map of a D -metric space X satisfying, for some positive real number p ,*

$$\begin{aligned} \rho(f^p x, f^p y, f^p z) \\ \leq \lambda \max \{ \rho(x, y, z), \rho(f^p x, f^p y, z), \rho(x, f^p x, z), \rho(y, f^p y, z), \rho(x, f^p y, z), \rho(y, f^p x, z) \} \end{aligned} \quad (3.41)$$

for all $x, y, z \in X$, where $0 \leq \lambda < 1$. Further if X is f -orbitally bounded and f -orbitally complete, then f has a unique fixed point $u \in X$, f^p is continuous, and f is f -orbitally continuous at u .

Note that Corollaries 3.8 and 3.9 include the fixed-point theorems of Rhoades [12] and Dhage [9] for the mappings characterized by the inequalities

$$\rho(fx, fy, fz) \leq \lambda \max \{ \rho(x, y, z), \rho(x, fx, z), \rho(y, fy, z), \rho(x, fy, z), \rho(y, fx, z) \}, \quad (3.42)$$

$$\rho(f^p x, f^p y, f^p z) \leq \lambda \max \{ \rho(x, y, z), \rho(x, f^p x, z), \rho(y, f^p y, z), \rho(x, f^p y, z), \rho(y, f^p x, z) \}, \quad (3.43)$$

for all $x, y, z \in X$ and $0 \leq \lambda < 1$.

3.2. Coincidentally commuting mappings. The coincidentally commuting mappings require some stronger condition than limit coincidentally commuting mappings and a good number of mathematicians have studied them on metric and D -metric spaces for the existence of their common fixed point. See [5, 11] and the references therein. The novelty of the fixed-point theorems for these coincidentally commuting mappings lies in the fact that here we do not require any of the maps under consideration to be continuous. Below, we prove a result in this direction and derive some interesting corollaries.

THEOREM 3.10. *Let X be a D -metric space and let $F : X \rightarrow CB(X)$ and $g : X \rightarrow X$ be two mappings satisfying (3.2). Further suppose that*

- (a) $F(X) \subseteq g(X)$,
- (b) $g(X)$ is bounded and complete,
- (c) $\{F, g\}$ is coincidentally commuting.

Then F and g have a unique common fixed point $u \in X$ such that $Fu = \{u\} = gu$. Moreover, if g is continuous at u , then F is also continuous at u in the Kasubai D -metric on X .

PROOF. Let $x \in X$ be arbitrary and define a sequence $\{y_n\} \subset X$ by (3.3). Clearly the sequence $\{y_n\}$ is well defined since $F(X) \subseteq g(X)$. Further we note that $\{y_n\} \subseteq g(X)$. We prove the conclusion of the theorem in two cases.

CASE 1. Suppose that $y_r = y_{r+1}$ for some $r \in \mathbb{N}$. Then proceeding with the arguments similar to [Case 1](#) of the proof of [Theorem 3.1](#), it is proved that $y_r = u$ is a common fixed point of F and g such that $Fu = \{u\} = gu$.

CASE 2. Assume that $y_n \neq y_{n+1}$ for each $n \in \mathbb{N}$. Then following [Case 2](#) of the proof of [Theorem 3.1](#), it is shown that $\{y_n\}$ is a D -Cauchy sequence. Since $g(X)$ is complete, there is a point $z \in g(X)$ such that $\lim_n y_n = z = \lim_n gx_n$. We will show that $\lim_n Fx_n = \{z\}$.

Now,

$$\begin{aligned}
 \lim_n \delta^r(Fx_n, Fx_n, z) &= \lim_n \delta^r(Fx_n, Fx_n, y_{n+1}) \\
 &\leq \lim_n \delta^r(Fx_n, Fx_n, Fx_n) \\
 &\leq \lim_n \phi(\max\{\rho^r(gx_n, gx_n, gx_n), \delta^r(Fx_n, Fx_n, gx_n), \delta^r(gx_n, Fx_n, gx_n)\}) \\
 &= \phi\left(\max\left\{0, \lim_n \delta^r(Fx_n, Fx_n, z)\right\}\right) \\
 &= \phi\left(\lim_n \delta^r(Fx_n, Fx_n, z)\right),
 \end{aligned} \tag{3.44}$$

which gives that $\lim_n Fx_n = \{z\}$.

Since $z \in g(X)$, there is a point $u \in X$ such that $gu = u$. We will show that $Fu = \{z\} = gu$. Now,

$$\begin{aligned}
 \delta^r(Fu, z, z) &= \lim_n \delta^r(Fu, Fx_n, Fx_n) \\
 &= \lim_n \delta^r(Fx_n, Fx_n, Fu) \\
 &\leq \lim_n \phi(\max\{\rho^r(gu, gx_n, gx_n), \delta^r(Fx_n, Fx_n, gu), \delta^r(gx_n, Fx_n, gu)\}) \\
 &= \phi(\max\{0, 0, 0\}) \\
 &= \phi(0) \\
 &= 0
 \end{aligned} \tag{3.45}$$

and so $Fu = gu = \{z\}$. Thus u is a coincidence point of F and g . The rest of the proof is similar to [Case 2](#) of the proof of [Theorem 3.1](#). We omitted the details. $\square$

As a consequence of [Theorem 3.10](#), we obtain the following corollaries.

COROLLARY 3.11. *Let $f, g : X \rightarrow X$ be two mappings satisfying (3.35). Suppose that*

- (a) $f(X) \subseteq g(X)$,
- (b) $g(X)$ is bounded and complete,
- (c) $\{f, g\}$ is coincidentally commuting.

Then f and g have a unique common fixed point u and if g is continuous at u , then f is also continuous at u .

COROLLARY 3.12. *Let X be a D -metric space and let $f, g : X \rightarrow X$ be two mappings satisfying*

$$\begin{aligned} \rho(fx, fy, fz) \\ \leq \lambda \max \{ \rho(gx, gy, gz), \rho(fx, fy, gz), \rho(gx, fx, gz), \\ \rho(gy, fy, gz), \rho(gx, fy, gz), \rho(gy, fx, gz) \} \end{aligned} \quad (3.46)$$

for all $x, y, z \in X$, where $0 \leq \lambda < 1$. Further suppose that hypotheses (a), (b), and (c) of [Corollary 3.11](#) hold. Then f and g have a unique common fixed point $u \in X$ and if g is continuous at u , then f is also continuous at u .

[Corollary 3.12](#) includes a common fixed-point theorem of Dhage [5] for the mappings f and g on a D -metric space characterized by the inequality

$$\begin{aligned} \rho(fx, fy, gz) \\ \leq \lambda \max \{ \rho(gx, gy, gz), \rho(gx, fx, gz), \\ \rho(gy, fy, gz), \rho(gx, fy, gz), \rho(gy, fx, gz) \} \end{aligned} \quad (3.47)$$

for all $x, y, z \in X$ and $0 \leq \lambda < 1$.

COROLLARY 3.13. *Let X be a D -metric space and let $f, g : X \rightarrow X$ be two mappings satisfying (3.37). Further suppose that*

- (a) $f^p(X) \subseteq g^q(X)$,
- (b) $g^p(X)$ is bounded and complete,
- (c) $\{f, g\}$ is commuting.

Then f and g have a unique common fixed point u and if g^q is continuous at u , then f^p is also continuous at u .

Notice that [Corollary 3.13](#) includes a class of common fixed-point mappings f and g on a D -metric space X characterized by the inequality

$$\begin{aligned} \rho(f^p x, f^p y, f^p z) \\ \leq \lambda \max \{ \rho(g^q x, g^q y, g^q z), \rho(g^q x, f^p x, g^q z), \\ \rho(g^q y, f^p y, g^q z), \rho(g^q x, f^p y, g^q z), \rho(g^q y, f^p x, g^q z) \} \end{aligned} \quad (3.48)$$

for all $x, y, z \in X$ and $0 \leq \lambda < 1$. See [5].

4. Weak commuting mappings in compact D -metric spaces. In this section, we prove some common fixed-point theorems for the pairs of singlevalued and multivalued coincidentally commuting mappings on a D -metric space satisfying a contraction condition more general than (4.3). But in this case the D -metric space under consideration is required to satisfy a stronger condition of compactness and the mappings under consideration are required to satisfy the continuity condition on the D -metric spaces. Our results of this section generalize some earlier known fixed-point theorems such as those of Dhage [9] and Rhoades [12] for single maps as well as for a pair of maps on D -metric spaces.

THEOREM 4.1. *Let X be a compact D -metric space and let $F : X \rightarrow \text{CB}(X)$ and $g : X \rightarrow X$ be two continuous mappings satisfying, for some positive real number r ,*

$$\begin{aligned} \delta^r(Fx, Fy, Fz) \\ < \max \{ \rho^r(gx, gy, gz), \delta^r(Fx, Fy, gz), \delta^r(gx, Fx, gz), \\ \delta^r(gy, Fy, gz), \delta^r(gx, Fy, gz), \delta^r(gy, Fx, gz) \} \end{aligned} \quad (4.1)$$

for all $x, y, z \in X$ for which the right-hand side is not zero. Further suppose that

- (a) $F(X) \subseteq g(X)$,
- (b) $\{F, g\}$ is limit coincidentally commuting.

Then F and g have a unique common fixed point $u \in X$ such that $Fu = \{u\} = gu$.

PROOF. From inequality (4.3), it follows that if F and g have a common fixed point $u \in X$, then it is unique and $Fu = \{u\} = gu$. Since X is compact and δ is continuous, both sides of inequality (4.1) are bounded on X . Now, there are two cases.

CASE 1. Suppose that the right-hand side of (4.1) is zero for some $x, y, z \in X$. Then, we have

$$Fx = gx = gz, \quad Fy = gy = gz. \quad (4.2)$$

Now, proceeding with the arguments similar to Case 1 of the proof of Theorem 3.1, it is proved that $u = Fx = gx$ is a common fixed point of F and g and so it is unique.

CASE 2. Suppose that the right-hand side of inequality (4.1) is not zero for all $x, y, z \in X$. Define a mapping $T : X \times X \times X \rightarrow (0, \infty)$ by

$$T(x, y, z) = \frac{\delta^r(Fx, Fy, Fz)}{M(x, y, z)}, \quad (4.3)$$

where

$$\begin{aligned} M(x, y, z) = \max \{ \rho^r(gx, gy, gz), \delta^r(Fx, Fy, gz), \delta^r(gx, Fx, gz), \\ \delta^r(gy, Fy, gz), \delta^r(gx, Fy, gz), \delta^r(gy, Fx, gz) \}. \end{aligned} \quad (4.4)$$

Clearly, the function T is well defined since $M(x, y, z) \neq 0$ for all $x, y, z \in X$. Since F and g are continuous, from the compactness of X it follows that the function T attains its maximum on X^3 at some point $u, v, w \in X$. Call the value c . It is clear from (4.1) that $0 < c < 1$. By the definition of c , we have $T(x, y, z) \leq c$ for all $x, y, z \in X$. This further, in view of (4.3), implies that

$$\begin{aligned} \delta^r(Fx, Fy, Fz) &\leq cM(x, y, z) \\ &= c \max \{ \rho^r(gx, gy, gz), \delta^r(Fx, Fy, Fz), \delta^r(gx, Fx, gx), \\ &\quad \delta^r(gy, Fy, gz), \delta^r(gx, Fy, gz), \delta^r(gy, Fx, gz) \} \end{aligned} \quad (4.5)$$

for all $x, y, z \in X$.

As X is compact, it is complete and $g(X)$ is bounded in view of the continuity of g on X . Now, the desired conclusion follows by an application of Theorem 3.1. This completes the proof. $\square$

Now we derive some interesting corollaries.

COROLLARY 4.2. *Let X be a compact D -metric space and let $F : X \rightarrow \text{CB}(X)$ be a continuous mapping satisfying*

$$\delta(Fx, Fy, Fz) < \max \{ \rho(x, y, z), \delta(Fx, Fy, z), \delta(x, Fx, z), \delta(y, Fy, z), \delta(x, Fy, z), \delta(y, Fx, z) \} \quad (4.6)$$

for all $x, y, z \in X$ for which the right-hand side is not zero. Then F has a unique fixed point $u \in X$ such that $Fu = \{u\}$.

PROOF. The proof follows by letting $g = I$ in Theorem 4.1, where I is the identity map on X . $\square$

COROLLARY 4.3 (see [3]). *Let X be a compact D -metric space and let $F : X \rightarrow \text{CB}(X)$ be a continuous mapping satisfying*

$$\delta(Fx, Fy, Fz) < \rho(x, y, z) \quad (4.7)$$

for all $x, y, z \in X$ for which $\rho(x, y, z) \neq 0$. Then F has a unique fixed point $u \in X$ such that $Fu = \{u\}$.

COROLLARY 4.4. *Let X be a compact D -metric space and let $f, g : X \rightarrow X$ be two continuous mappings satisfying*

$$\begin{aligned} \rho(fx, fy, fz) &< \max \{ \rho(gx, gy, gz), \rho(fx, fy, gz), \rho(gx, fx, gz), \\ &\quad \rho(gy, fy, gz), \rho(gx, fy, gz), \rho(gy, fx, gz) \} \end{aligned} \quad (4.8)$$

for all $x, y, z \in X$ for which the right-hand side is not zero. Further suppose that

- (a) $f(X) \subseteq g(X)$,
- (b) $\{f, g\}$ is limit coincidentally commuting.

Then f and g have a unique common fixed point.

PROOF. The proof follows by letting $F = \{f\}$, a single-valued mapping in [Theorem 4.1](#). □

COROLLARY 4.5. Let X be a compact D -metric space and let $f : X \rightarrow X$ be a continuous mapping satisfying

$$\rho(fx, fy, fz) < \max \{ \rho(x, y, z), \rho(fx, fy, z), \rho(x, fx, z), \rho(y, fy, z), \rho(x, fy, z), \rho(y, fx, z) \} \quad (4.9)$$

for all $x, y, z \in X$ for which the right-hand side is not zero. Then f has a unique fixed point.

PROOF. The conclusion follows by letting $g = I$ in [Corollary 4.4](#), where I is the identity map on X . □

Note that Corollaries [4.4](#) and [4.5](#) include the fixed-point theorems of Dhage [\[5\]](#) and Rhoades [\[12\]](#) for the mappings f and g on a D -metric space X characterized by the inequalities

$$\rho(fx, fy, fz) < \max \{ \rho(gx, gy, gz), \rho(gx, fx, gz), \rho(gy, fy, gz), \rho(gx, fy, gz), \rho(gy, fx, gz) \}, \quad (4.10)$$

$$\rho(fx, fy, fz) < \max \{ \rho(x, y, z), \rho(x, fx, z), \rho(y, fy, z), \rho(x, fy, z), \rho(y, fx, z) \}, \quad (4.11)$$

respectively.

THEOREM 4.6. Let X be a D -metric space and let $F : X \rightarrow \text{CB}(X)$, $g : X \rightarrow X$ be two continuous mappings satisfying [\(4.1\)](#). Suppose further that

- (a) $F(X) \subseteq g(X)$,
- (b) $g(X)$ is compact,
- (c) $\{f, g\}$ is coincidentally commuting.

Then F and g have a unique common fixed point $u \in X$ such that $Fu = \{u\} = gu$.

PROOF. Let $A = g(X)$. Then A is a compact D -metric space and F and g define the maps $F : A \rightarrow \text{CB}(A)$ and $g : A \rightarrow A$. Now, the desired conclusion follows by an application of [Theorem 4.1](#). □

COROLLARY 4.7. Let X be a D -metric space and let $f, g : X \rightarrow X$ be two continuous mappings satisfying [\(4.8\)](#). Further suppose that

- (a) $f(X) \subseteq g(X)$,
- (b) $g(X)$ is compact,
- (c) $\{f, g\}$ is coincidentally commuting.

Then f and g have a unique common fixed point.

5. Remarks and conclusion. It has been noted in [6, 10] that the fixed-point theorems for the limit coincidentally commuting mappings have some nice applications to approximation theory, and therefore it is of interest to discuss the fixed-point theorems for a wide class of coincidentally commuting mappings in a D -metric space. The terms “compatible” and “ δ -compatible” have been used by Jungck and Rhoades [11] for limit coincidentally commuting and coincidentally commuting mappings, respectively, but our terminologies are natural and more informative than the previous one patterned after [4]. Further we note that a similar study can be made for coincidentally pseudocommuting mappings on a D -metric space and analogously for limit coincidentally pseudocommuting mappings. But in order to prove fixed-point theorems for these classes of weakly pseudocommuting mappings, we require a stronger contraction condition for the mappings F and g under consideration:

$$\begin{aligned} \delta^r(Fx, Fy, Fz) \\ \leq \phi(\max\{\rho^r(gx, gy, gz), D^r(Fx, Fy, gz), D^r(gx, Fx, gz), \\ D^r(gy, Fy, gz), D^r(gx, Fy, gz), D^r(gy, Fx, gz)\}). \end{aligned} \quad (5.1)$$

Obviously, condition (5.1) implies condition (3.2) on a D -metric space X and hence the fixed-point theorems for weakly pseudocommuting mappings can be obtained very easily with appropriate modifications. Finally, we close this discussion with the following open question.

OPEN QUESTION. Can we prove fixed-point theorems for a class of multivalued mapping F on a D -metric space X satisfying the generalized contraction condition

$$\begin{aligned} \kappa(Fx, Fy, Fz) \leq \lambda \max\{\rho(x, y, z), D(Fx, Fy, z), D(x, Fx, z), \\ D(y, Fy, z), D(x, Fy, z), D(y, Fx, z)\} \end{aligned} \quad (5.2)$$

for all $x, y, z \in X$ and $0 \leq \lambda < 1$?

ACKNOWLEDGMENT. The authors are thankful to Professor B. E. Rhoades (USA) for giving some useful suggestions for the improvement of this paper.

REFERENCES

- [1] B. C. Dhage, *A study of some fixed point theorem*, Ph.D. thesis, Marathwada University, Aurangabad, India, 1984.
- [2] ———, *Generalised metric spaces and mappings with fixed point*, Bull. Calcutta Math. Soc. **84** (1992), no. 4, 329–336.

- [3] ———, *Generalized D-metric spaces and multi-valued contraction mappings*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.) **44** (1998), no. 1, 179–200.
- [4] ———, *A common fixed point principle in D-metric spaces*, Bull. Calcutta Math. Soc. **91** (1999), no. 6, 475–480.
- [5] ———, *On common fixed points of pairs of coincidentally commuting mappings in D-metric spaces*, Indian J. Pure Appl. Math. **30** (1999), no. 4, 395–406.
- [6] ———, *Applications of some fixed point theorems in approximation theory*, Math. Sci. Res. Hot-Line **4** (2000), no. 9, 63–69.
- [7] ———, *A common fixed point theorem for a pair of limit coincidentally commuting mappings in D-metric spaces*, Math. Sci. Res. Hot-Line **4** (2000), no. 2, 45–55.
- [8] ———, *Generalized metric spaces and topological structure. I*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.) **46** (2000), no. 1, 3–24.
- [9] ———, *On some fixed point theorems for contractive mapping in D-metric spaces*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.) **46** (2000), no. 1, 31–38.
- [10] ———, *A result in best approximation theory*, Indian J. Pure Appl. Math. **32** (2001), no. 8, 1209–1216.
- [11] G. Jungck and B. E. Rhoades, *Fixed points for set valued functions without continuity*, Indian J. Pure Appl. Math. **29** (1998), no. 3, 227–238.
- [12] B. E. Rhoades, *A fixed point theorem for generalized metric spaces*, Int. J. Math. Math. Sci. **19** (1996), no. 3, 457–460.

B. C. Dhage: Kasubai, Gurukul Colony, Ahmedpur 413515, Maharashtra, India
E-mail address: bcd20012001@yahoo.co.in

A. Jennifer Asha: Ramanujan Institute for Advanced Study in Mathematics, University of Madras, Chennai 600005, India

S. M. Kang: Department of Mathematics and Research Institute of Natural Science, Gyeongsang National University, Chinju 660-701, Korea
E-mail address: smkang@nongae.gsnu.ac.kr