THE CONJUGATION OPERATOR ON $A_q(G)$

SANJIV KUMAR GUPTA

Received 5 March 2002

Let $q > 2$. We prove that the conjugation operator H does not extend to a bounded operator on the space of integrable functions defined on any compact abelian group with the Fourier transform in l_q .

2000 Mathematics Subject Classification: 42A50, 43A17.

Let G be a compact abelian group with dual Γ . For $1 \leq q < \infty$, the space A_q is defined as

$$A_q(G) = \{f : f \in L^1(G), \hat{f} \in l_q(\Gamma)\} \quad (1)$$

with the norm $\|f\|_{A_q} = \|f\|_{L^1} + \|\hat{f}\|_{l_q}$. Then $A_q(G)$ is a commutative semisimple Banach algebra with maximal ideal space Γ , in which the set of trigonometric polynomials is dense [4]. The A_p -spaces have been studied in [1, 6].

If G is, in addition, a connected group, then its dual can be ordered; there exists a semigroup $P \subset \Gamma$ such that $P \cap -P = \{0\}$, $P \cup -P = \Gamma$ (see [5]), and we say that $\gamma \in \Gamma$ is positive if $\gamma \in P$. If $f = \sum_{\gamma \in F} \hat{f}(\gamma)\gamma$ is a trigonometric polynomial, the conjugation operator H is defined as

$$Hf = \sum_{\gamma \in F} \text{sgn}(\gamma) \hat{f}(\gamma)\gamma, \quad (2)$$

where $\text{sgn}(\gamma) = +1$ if $\gamma \in P$, -1 if $\gamma \in -P$, and 0 if $\gamma = 0$.

If $1 \leq q \leq 2$, then $A_q(G) \subset L^2(G)$, and it is easy to see that H extends to a bounded operator on $A_q(G)$. It was mentioned in [5] that the corresponding result for $q > 2$ is not known. Note that H extends to a bounded operator on $A_q(G)$ if and only if H extends to a bounded operator from $A_q(G)$ to $L^1(G)$. In [5], the following theorem was proved.

THEOREM 1. *Let G be a compact, connected abelian group and P any fixed order on Γ . If $q > 2$ and ϕ is a Young's function, then the conjugation operator H does not extend to a bounded operator from $A_q(G)$ to $L^\phi(G)$.*

We prove in [Theorem 2](#) that H does not extend to a bounded operator on $A_q(G)$, $q > 2$, thus answering the problem mentioned in [5]. Also, [Theorem 1](#) follows from our theorem ([Theorem 2](#)). [Theorem 2](#) was proved for the circle group in [2] but for the completeness sake, we give it below.

THEOREM 2. *Let G be a compact, connected abelian group and P any fixed order on Γ . If $q > 2$, then the conjugation operator H does not extend to a bounded operator on $A_q(G)$.*

PROOF. We will show that H does not extend to a bounded operator from $A_q(G)$ to $L^1(G)$. The proof is divided into three steps.

STEP 1. Let $G = \mathbf{T}$, the circle group. Suppose that H extends as a bounded operator from A_q to L^1 . Choose $\mu_0 \in M(\mathbf{T})$, $\hat{\mu}_0 \in l_q$ such that μ_0 is not absolutely continuous. Define $T : L^1 \rightarrow L^1$ by

$$Tf = H(f * \mu_0), \quad (3)$$

where T is well defined as $f * \mu_0 \in A_q$ and H maps A_q into L^1 by our assumption on H . Hence, T is a multiplier from L^1 to L^1 , and therefore is given by a measure $\mu \in M(\mathbf{T})$ (see [3]). Hence

$$\operatorname{sgn}(n)\hat{\mu}_0(n) = \hat{\mu}(n). \quad (4)$$

Observe that

$$\hat{\mu}_0 = \frac{\hat{\mu}_0 + \hat{\mu}}{2} + \frac{\hat{\mu}_0 - \hat{\mu}}{2}. \quad (5)$$

Now, $(\hat{\mu}_0 + \hat{\mu})/2$ and $(\hat{\mu}_0 - \hat{\mu})/2$ are absolutely continuous by F. and M. Riesz theorem. Hence, $\hat{\mu}_0$ is absolutely continuous, which contradicts the choice of μ_0 . Hence, H is unbounded on A_q , $q > 2$.

STEP 2. Let I be a closed subgroup of G such that H does not extend as a bounded operator on $A_q(G/I)$. Then H does not extend as a bounded operator on $A_q(G)$.

PROOF. Let (f_n) be a sequence of trigonometric polynomials on G/I such that

$$\|Hf_n\|_{L^1(G/I)} \rightarrow \infty, \quad \|f_n\|_{A_q(G/I)} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (6)$$

Let $g_n = f_n \circ q$, where $q : G \rightarrow G/I$ is the quotient map. Then it is easily seen that $Hg_n = (Hf_n) \circ q$, $\|Hg_n\|_{L^1(G)} = \|Hf_n\|_{L^1(G/I)}$, and $\|f_n \circ q\|_{A_q(G)} = \|f_n\|_{A_q(G/I)}$. Hence

$$\|Hg_n\|_{L^1(G)} \rightarrow \infty, \quad \|g_n\|_{A_q(G)} \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (7)$$

and Step 2 follows. $\square$

STEP 3. Since G is connected, Γ contains an element of infinite order, say y_0 (see [3]). Let S denote the subgroup generated by y_0 and let $I = S^\perp$. Then G/H is isomorphic to the circle group $\mathbf{T}$. Now, the proof of the theorem follows from Steps 1 and 2. $\square$

ACKNOWLEDGMENT. This research was supported by Sultan Qaboos University, Oman.

REFERENCES

- [1] L. M. Bloom and W. R. Bloom, *Multipliers on spaces of functions with p -summable Fourier transforms*, Harmonic Analysis (Luxembourg, 1987), Lecture Notes in Math., vol. 1359, Springer, Berlin, 1988, pp. 100–112.
- [2] J. T. Burnham, H. E. Krogstad, and R. Larsen, *Multipliers and the Hilbert distribution*, Nanta Math. **8** (1975), no. 2, 95–103.
- [3] E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis. Vol. I: Structure of Topological Groups. Integration Theory, Group Representations*, Die Grundlehren der Mathematischen Wissenschaften, vol. 115, Academic Press, New York, 1963.
- [4] ———, *Abstract Harmonic Analysis. Vol. II: Structure and Analysis for Compact Groups. Analysis on Locally Compact Abelian Groups*, Die Grundlehren der Mathematischen Wissenschaften, vol. 152, Springer-Verlag, New York, 1970.
- [5] W. Rudin, *Fourier Analysis on Groups*, Interscience Tracts in Pure and Applied Mathematics, no. 12, Interscience Publishers, New York, 1962.
- [6] U. B. Tewari and A. K. Gupta, *The algebra of functions with Fourier transforms in a given function space*, Bull. Austral. Math. Soc. **9** (1973), 73–82.

Sanjiv Kumar Gupta: Department of Mathematics and Statistics, College of Science, Sultan Qaboos University, P.O. Box 36, Al-Khaid 123, Sultanate of Oman
E-mail address: gupta@squ.edu.om