α -COMPACTNESS IN SMOOTH TOPOLOGICAL SPACES

CHUN-KEE PARK, WON KEUN MIN, and MYEONG HWAN KIM

Received 3 March 2003

We introduce the concepts of smooth α -closure and smooth α -interior of a fuzzy set which are generalizations of smooth closure and smooth interior of a fuzzy set defined by Demirci (1997) and obtain some of their structural properties.

2000 Mathematics Subject Classification: 54A40.

1. Introduction. Badard [1] introduced the concept of a smooth topological space which is a generalization of Chang's fuzzy topological space [2]. Many mathematical structures in smooth topological spaces were introduced and studied. In particular, Gayyar et al. [5] and Demirci [3, 4] introduced the concepts of smooth closure and smooth interior of a fuzzy set and several types of compactness in smooth topological spaces and obtained some properties of them.

In this paper, we define the smooth α -closure and smooth α -interior of a fuzzy set and investigate some of their properties. In fact, the smooth α -closure and smooth α -interior of a fuzzy set coincide with the smooth closure and smooth interior of a fuzzy set defined in [3] when $\alpha = 0$. We also introduce the concepts of several types of α -compactness using smooth α -closure and smooth α -interior of a fuzzy set and investigate some of their properties.

2. Preliminaries. In this section, we give some notations and definitions which are to be used in the sequel. Let X be a set and let $I = [0, 1]$ be the unit interval of the real line. Let I^X denote the set of all fuzzy sets of X . Let 0_X and 1_X denote the characteristic functions of ϕ and X , respectively.

A smooth topological space (s.t.s.) [6] is an ordered pair (X, τ) , where X is a nonempty set and $\tau : I^X \rightarrow I$ is a mapping satisfying the following conditions:

- (1) $\tau(0_X) = \tau(1_X) = 1$;
- (2) for all $A, B \in I^X$, $\tau(A \cap B) \geq \tau(A) \wedge \tau(B)$;
- (3) for every subfamily $\{A_i : i \in J\} \subseteq I^X$, $\tau(\cup_{i \in J} A_i) \geq \wedge_{i \in J} \tau(A_i)$.

Then the mapping $\tau : I^X \rightarrow I$ is called a smooth topology on X . The number $\tau(A)$ is called the degree of openness of A .

A mapping $\tau^* : I^X \rightarrow I$ is called a smooth cotopology [6] if and only if the following three conditions are satisfied:

- (1) $\tau^*(0_X) = \tau^*(1_X) = 1$;
- (2) for all $A, B \in I^X$, $\tau^*(A \cup B) \geq \tau^*(A) \wedge \tau^*(B)$;

(3) for every subfamily $\{A_i : i \in J\} \subseteq I^X$, $\tau^*(\cap_{i \in J} A_i) \geq \wedge_{i \in J} \tau^*(A_i)$.

If τ is a smooth topology on X , then the mapping $\tau^* : I^X \rightarrow I$, defined by $\tau^*(A) = \tau(A^c)$ where A^c denotes the complement of A , is a smooth cotopology on X . Conversely, if τ^* is a smooth cotopology on X , then the mapping $\tau : I^X \rightarrow I$, defined by $\tau(A) = \tau^*(A^c)$, is a smooth topology on X [6].

For the s.t.s. (X, τ) and $\alpha \in [0, 1]$, the family $\tau_\alpha = \{A \in I^X : \tau(A) \geq \alpha\}$ defines a Chang's fuzzy topology (CFT) on X [2]. The family of all closed fuzzy sets with respect to τ_α is denoted by τ_α^* and we have $\tau_\alpha^* = \{A \in I^X : \tau^*(A) \geq \alpha\}$. For $A \in I^X$ and $\alpha \in [0, 1]$, the τ_α -closure (resp., τ_α -interior) of A , denoted by $\text{cl}_\alpha(A)$ (resp., $\text{int}_\alpha(A)$), is defined by $\text{cl}_\alpha(A) = \cap \{K \in \tau_\alpha^* : A \subseteq K\}$ (resp., $\text{int}_\alpha(A) = \cup \{K \in \tau_\alpha : K \subseteq A\}$).

Demirci [3] introduced the concepts of smooth closure and smooth interior in smooth topological spaces as follows.

Let (X, τ) be an s.t.s. and $A \in I^X$. Then the τ -smooth closure (resp., τ -smooth interior) of A , denoted by $\bar{A}$ (resp., A°), is defined by $\bar{A} = \cap \{K \in I^X : \tau^*(K) > 0, A \subseteq K\}$ (resp., $A^\circ = \cup \{K \in I^X : \tau(K) > 0, K \subseteq A\}$).

Let (X, τ) and (Y, σ) be two smooth topological spaces. A function $f : X \rightarrow Y$ is called smooth continuous with respect to τ and σ [6] if and only if $\tau(f^{-1}(A)) \geq \sigma(A)$ for every $A \in I^Y$. A function $f : X \rightarrow Y$ is called weakly smooth continuous with respect to τ and σ [6] if and only if $\sigma(A) > 0 \Rightarrow \tau(f^{-1}(A)) > 0$ for every $A \in I^Y$.

A function $f : X \rightarrow Y$ is smooth continuous with respect to τ and σ if and only if $\tau^*(f^{-1}(A)) \geq \sigma^*(A)$ for every $A \in I^Y$. A function $f : X \rightarrow Y$ is weakly smooth continuous with respect to τ and σ if and only if $\sigma^*(A) > 0 \Rightarrow \tau^*(f^{-1}(A)) > 0$ for every $A \in I^Y$ [6].

A function $f : X \rightarrow Y$ is called smooth open (resp., smooth closed) with respect to τ and σ [6] if and only if $\tau(A) \leq \sigma(f(A))$ (resp., $\tau^*(A) \leq \sigma^*(f(A))$) for every $A \in I^X$.

A function $f : X \rightarrow Y$ is called smooth preserving (resp., strict smooth preserving) with respect to τ and σ [5] if and only if $\sigma(A) \geq \sigma(B) \Leftrightarrow \tau(f^{-1}(A)) \geq \tau(f^{-1}(B))$ (resp., $\sigma(A) > \sigma(B) \Leftrightarrow \tau(f^{-1}(A)) > \tau(f^{-1}(B))$) for every $A, B \in I^Y$.

If $f : X \rightarrow Y$ is a smooth preserving function (resp., a strict smooth preserving function) with respect to τ and σ , then $\sigma^*(A) \geq \sigma^*(B) \Leftrightarrow \tau^*(f^{-1}(A)) \geq \tau^*(f^{-1}(B))$ (resp., $\sigma^*(A) > \sigma^*(B) \Leftrightarrow \tau^*(f^{-1}(A)) > \tau^*(f^{-1}(B))$) for every $A, B \in I^Y$ [5].

A function $f : X \rightarrow Y$ is called smooth open preserving (resp., strict smooth open preserving) with respect to τ and σ [5] if and only if $\tau(A) \geq \tau(B) \Rightarrow \sigma(f(A)) \geq \sigma(f(B))$ (resp., $\tau(A) > \tau(B) \Rightarrow \sigma(f(A)) > \sigma(f(B))$) for every $A, B \in I^X$.

3. Smooth α -closure and smooth α -interior. In this section, we introduce the concepts of smooth α -closure and smooth α -interior of a fuzzy set in smooth topological spaces and investigate some properties of them.

DEFINITION 3.1. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A \in I^X$. The τ -smooth α -closure (resp., τ -smooth α -interior) of A , denoted by $\overline{A}_\alpha$ (resp., A_α^o), is defined by $\overline{A}_\alpha = \cap \{K \in I^X : \tau^*(K) > \alpha\tau^*(A), A \subseteq K\}$ (resp., $A_\alpha^o = \cup \{K \in I^X : \tau(K) > \alpha\tau(A), K \subseteq A\}$).

THEOREM 3.2. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A, B \in I^X$. Then

- (a) $\tau^*(\overline{A}_\alpha) \geq \alpha\tau^*(A)$,
- (b) $\tau(A_\alpha^o) \geq \alpha\tau(A)$,
- (c) $A \subseteq B$ and $\tau^*(A) \leq \tau^*(B) \Rightarrow \overline{A}_\alpha \subseteq \overline{B}_\alpha$,
- (d) $A \subseteq B$ and $\tau(B) \leq \tau(A) \Rightarrow A_\alpha^o \subseteq B_\alpha^o$.

PROOF. (a) and (b) follow directly from [Definition 3.1](#).

(c) If $A \subseteq B$ and $\tau^*(A) \leq \tau^*(B)$, then $K \in \{K \in I^X : \tau^*(K) > \alpha\tau^*(B), B \subseteq K\} \Rightarrow K \in \{K \in I^X : \tau^*(K) > \alpha\tau^*(A), A \subseteq K\}$. Hence $\overline{A}_\alpha \subseteq \overline{B}_\alpha$.

(d) The proof is similar to the proof of (c). □

THEOREM 3.3. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A \in I^X$. Then

- (a) $(\overline{A}_\alpha)^c = (A^c)_\alpha^o$,
- (b) $\overline{A}_\alpha = ((A^c)_\alpha^o)^c$,
- (c) $(A_\alpha^o)^c = (\overline{A^c})_\alpha$,
- (d) $A_\alpha^o = (\overline{(A^c)_\alpha})^c$.

PROOF. (a) From [Definition 3.1](#), we have

$$\begin{aligned}
 (\overline{A}_\alpha)^c &= (\cap \{K \in I^X : \tau^*(K) > \alpha\tau^*(A), A \subseteq K\})^c \\
 &= \cup \{K^c : K \in I^X, \tau(K^c) = \tau^*(K) > \alpha\tau^*(A) = \alpha\tau(A^c), K^c \subseteq A^c\} \\
 &= \cup \{U \in I^X : \tau(U) > \alpha\tau(A^c), U \subseteq A^c\} \\
 &= (A^c)_\alpha^o.
 \end{aligned} \tag{3.1}$$

(b), (c), and (d) are easily obtained from (a). □

THEOREM 3.4. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A, B \in I^X$. Then

- (a) $(0_X)_\alpha = 0_X$,
- (b) $A \subseteq \overline{A}_\alpha$,
- (c) $\overline{A}_\alpha \subseteq (\overline{A}_\alpha)_\alpha$,
- (d) $\overline{A}_\alpha \cap \overline{B}_\alpha \subseteq \overline{(A \cup B)}_\alpha$.

PROOF. (a) and (b) are easily obtained from [Definition 3.1](#). (c) follows directly from (b).

(d) For every $A, B \in I^X$, we have

$$\begin{aligned}
 \overline{(A \cup B)}_\alpha &= \cap \{K \in I^X : \tau^*(K) > \alpha\tau^*(A \cup B), A \cup B \subseteq K\} \\
 &\supseteq \cap \{K \in I^X : \tau^*(K) > \alpha\tau^*(A) \wedge \alpha\tau^*(B), A \cup B \subseteq K\}
 \end{aligned}$$

$$\begin{aligned}
&= \cap \{K \in I^X : \tau^*(K) > \alpha \tau^*(A) \\
&\quad \text{or } \tau^*(K) > \alpha \tau^*(B), A \subseteq K, B \subseteq K\} \\
&= \cap \{K \in I^X : (\tau^*(K) > \alpha \tau^*(A), A \subseteq K, B \subseteq K) \\
&\quad \text{or } (\tau^*(K) > \alpha \tau^*(B), A \subseteq K, B \subseteq K)\} \\
&\supseteq \cap [\{K \in I^X : \tau^*(K) > \alpha \tau^*(A), A \subseteq K\} \\
&\quad \cup \{K \in I^X : \tau^*(K) > \alpha \tau^*(B), B \subseteq K\}] \\
&= [\cap \{K \in I^X : \tau^*(K) > \alpha \tau^*(A), A \subseteq K\}] \\
&\quad \cap [\cap \{K \in I^X : \tau^*(K) > \alpha \tau^*(B), B \subseteq K\}] \\
&= \overline{A}_\alpha \cap \overline{B}_\alpha.
\end{aligned} \tag{3.2}$$

□

THEOREM 3.5. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A, B \in I^X$. Then

- (a) $(1_X)_\alpha^o = 1_X$,
- (b) $A_\alpha^o \subseteq A$,
- (c) $(A_\alpha^o)_\alpha^o \subseteq A_\alpha^o$,
- (d) $(A \cap B)_\alpha^o \subseteq A_\alpha^o \cup B_\alpha^o$.

PROOF. The proof is similar to the proof of [Theorem 3.4](#). □

THEOREM 3.6. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A \in I^X$. Then

- (a) $\tau^*(A) > 0 \Rightarrow \overline{A}_\alpha = A$,
- (b) $\tau(A) > 0 \Rightarrow A_\alpha^o = A$.

PROOF. (a) Let $\tau^*(A) > 0$. Then $A \in \{K \in I^X : \tau^*(K) > \alpha \tau^*(A), A \subseteq K\}$. By [Definition 3.1](#), $\overline{A}_\alpha \subseteq A$. By [Theorem 3.4](#), $A \subseteq \overline{A}_\alpha$. Hence $\overline{A}_\alpha = A$.

(b) Let $\tau(A) > 0$. Then $A \in \{K \in I^X : \tau(K) > \alpha \tau(A), K \subseteq A\}$. By [Definition 3.1](#), $A \subseteq A_\alpha^o$. By [Theorem 3.5](#), $A_\alpha^o \subseteq A$. Hence $A_\alpha^o = A$. □

REMARK 3.7. Let (X, τ) be an s.t.s., $\alpha_1, \alpha_2 \in [0, 1)$ with $\alpha_1 \leq \alpha_2$, and $A \in I^X$. Then $\overline{A}_{\alpha_1} \subseteq \overline{A}_{\alpha_2}$ and $A_{\alpha_2}^o \subseteq A_{\alpha_1}^o$.

THEOREM 3.8. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A \in I^X$. Then

- (a) $\overline{A}_\alpha = \cap_{\beta > \alpha \tau^*(A)} \text{cl}_\beta(A)$,
- (b) $A_\alpha^o = \cup_{\beta > \alpha \tau(A)} \text{int}_\beta(A)$.

PROOF. (a) For each $x \in X$, we have

$$\begin{aligned}
\overline{A}_\alpha(x) &= [\cap \{K \in I^X : \tau^*(K) > \alpha \tau^*(A), A \subseteq K\}](x) \\
&= \inf \{K(x) : K \in I^X, \tau^*(K) > \alpha \tau^*(A), A \subseteq K\} \\
&= \inf_{\beta > \alpha \tau^*(A)} \inf \{K(x) : K \in I^X, \tau^*(K) \geq \beta, A \subseteq K\}
\end{aligned}$$

$$\begin{aligned}
&= \inf_{\beta > \alpha\tau^*(A)} [\cap \{K \in I^X : \tau^*(K) \geq \beta, A \subseteq K\}](x) \\
&= \inf_{\beta > \alpha\tau^*(A)} \text{cl}_\beta(A)(x) \\
&= [\cap_{\beta > \alpha\tau^*(A)} \text{cl}_\beta(A)](x).
\end{aligned}
\tag{3.3}$$

Hence, $\overline{A}_\alpha = \cap_{\beta > \alpha\tau^*(A)} \text{cl}_\beta(A)$.

(b) The proof is similar to that of (a). $\square$

REMARK 3.9. Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and $A \in I^X$. From Theorems 3.4, 3.5, and 3.8, we easily obtain the following:

- (a) if there exists a $\beta \in (\alpha\tau^*(A), 1]$ such that $A = \text{cl}_\beta(A)$, then $A = \overline{A}_\alpha$;
- (b) if there exists a $\beta \in (\alpha\tau(A), 1]$ such that $A = \text{int}_\beta(A)$, then $A = A_\alpha^o$.

DEFINITION 3.10. Let (X, τ) and (Y, σ) be two smooth topological spaces and let $\alpha \in [0, 1)$. A function $f : X \rightarrow Y$ is called smooth α -preserving (resp., strict smooth α -preserving) with respect to τ and σ if and only if $\sigma(A) \geq \alpha\sigma(B) \Leftrightarrow \tau(f^{-1}(A)) \geq \alpha\tau(f^{-1}(B))$ (resp., $\sigma(A) > \alpha\sigma(B) \Leftrightarrow \tau(f^{-1}(A)) > \alpha\tau(f^{-1}(B))$) for every $A, B \in I^Y$.

A function $f : X \rightarrow Y$ is called smooth open α -preserving (resp., strict smooth open α -preserving) with respect to τ and σ if and only if $\tau(A) \geq \alpha\tau(B) \Rightarrow \sigma(f(A)) \geq \alpha\sigma(f(B))$ (resp., $\tau(A) > \alpha\tau(B) \Rightarrow \sigma(f(A)) > \alpha\sigma(f(B))$) for every $A, B \in I^X$.

THEOREM 3.11. Let (X, τ) and (Y, σ) be two smooth topological spaces and let $\alpha \in [0, 1)$. If $f : X \rightarrow Y$ is a smooth α -preserving function (resp., a strict smooth α -preserving function) with respect to τ and σ , then $\sigma^*(A) \geq \alpha\sigma^*(B) \Leftrightarrow \tau^*(f^{-1}(A)) \geq \alpha\tau^*(f^{-1}(B))$ (resp., $\sigma^*(A) > \alpha\sigma^*(B) \Leftrightarrow \tau^*(f^{-1}(A)) > \alpha\tau^*(f^{-1}(B))$) for every $A, B \in I^Y$.

PROOF. If $f : X \rightarrow Y$ is a smooth α -preserving function with respect to τ and σ , then

$$\begin{aligned}
\sigma^*(A) \geq \alpha\sigma^*(B) &\Leftrightarrow \sigma(A^c) \geq \alpha\sigma(B^c) \\
&\Leftrightarrow \tau(f^{-1}(A^c)) \geq \alpha\tau(f^{-1}(B^c)) \\
&\Leftrightarrow \tau((f^{-1}(A))^c) \geq \alpha\tau((f^{-1}(B))^c) \\
&\Leftrightarrow \tau^*(f^{-1}(A)) \geq \alpha\tau^*(f^{-1}(B))
\end{aligned}
\tag{3.4}$$

for every $A, B \in I^Y$.

The proof is similar when $f : X \rightarrow Y$ is a strict smooth α -preserving function with respect to τ and σ . $\square$

THEOREM 3.12. *Let (X, τ) and (Y, σ) be two smooth topological spaces and let $\alpha \in [0, 1)$. If a function $f : X \rightarrow Y$ is injective and strict smooth α -preserving with respect to τ and σ , then $f(\overline{A}_\alpha) \subseteq \overline{(f(A))}_\alpha$ for every $A \in I^X$.*

PROOF. For every $A \in I^X$, we have

$$\begin{aligned}
 f^{-1}(\overline{(f(A))}_\alpha) &= f^{-1}[\cap \{U \in I^Y : \sigma^*(U) > \alpha \sigma^*(f(A)), f(A) \subseteq U\}] \\
 &\supseteq f^{-1}[\cap \{U \in I^Y : \tau^*(f^{-1}(U)) > \alpha \tau^*(A), A \subseteq f^{-1}(U)\}] \\
 &= \cap \{f^{-1}(U) \in I^X : \tau^*(f^{-1}(U)) > \alpha \tau^*(A), A \subseteq f^{-1}(U)\} \quad (3.5) \\
 &\supseteq \cap \{K \in I^X : \tau^*(K) > \alpha \tau^*(A), A \subseteq K\} \\
 &= \overline{A}_\alpha.
 \end{aligned}$$

Hence, $f(\overline{A}_\alpha) \subseteq \overline{(f(A))}_\alpha$. □

THEOREM 3.13. *Let (X, τ) and (Y, σ) be two smooth topological spaces and let $\alpha \in [0, 1)$. If a function $f : X \rightarrow Y$ is strict smooth α -preserving with respect to τ and σ , then*

- (a) $\overline{(f^{-1}(A))}_\alpha \subseteq f^{-1}(\overline{A}_\alpha)$ for every $A \in I^Y$,
- (b) $f^{-1}(A_\alpha^o) \subseteq (f^{-1}(A))_\alpha^o$ for every $A \in I^Y$.

PROOF. (a) For every $A \in I^Y$, we have

$$\begin{aligned}
 f^{-1}(\overline{A}_\alpha) &= f^{-1}[\cap \{U \in I^Y : \sigma^*(U) > \alpha \sigma^*(A), A \subseteq U\}] \\
 &\supseteq f^{-1}[\cap \{U \in I^Y : \tau^*(f^{-1}(U)) > \alpha \tau^*(f^{-1}(A)), f^{-1}(A) \subseteq f^{-1}(U)\}] \\
 &= \cap \{f^{-1}(U) \in I^X : \tau^*(f^{-1}(U)) > \alpha \tau^*(f^{-1}(A)), f^{-1}(A) \subseteq f^{-1}(U)\} \\
 &\supseteq \cap \{K \in I^X : \tau^*(K) > \alpha \tau^*(f^{-1}(A)), f^{-1}(A) \subseteq K\} \\
 &= \overline{(f^{-1}(A))}_\alpha. \quad (3.6)
 \end{aligned}$$

(b) For every $A \in I^Y$, we have

$$\begin{aligned}
 f^{-1}(A_\alpha^o) &= f^{-1}[\cup \{U \in I^Y : \sigma(U) > \alpha \sigma(A), U \subseteq A\}] \\
 &\subseteq f^{-1}[\cup \{U \in I^Y : \tau(f^{-1}(U)) > \alpha \tau(f^{-1}(A)), f^{-1}(U) \subseteq f^{-1}(A)\}] \\
 &= \cup \{f^{-1}(U) \in I^X : \tau(f^{-1}(U)) > \alpha \tau(f^{-1}(A)), f^{-1}(U) \subseteq f^{-1}(A)\} \\
 &\subseteq \cup \{K \in I^X : \tau(K) > \alpha \tau(f^{-1}(A)), K \subseteq f^{-1}(A)\} \\
 &= (f^{-1}(A))_\alpha^o. \quad (3.7)
 \end{aligned}$$

□

THEOREM 3.14. *Let (X, τ) and (Y, σ) be two smooth topological spaces and let $\alpha \in [0, 1)$. If a function $f : X \rightarrow Y$ is strict smooth open α -preserving with respect to τ and σ , then $f(A_\alpha^o) \subseteq (f(A))_\alpha^o$ for every $A \in I^X$.*

PROOF. For every $A \in I^X$, we have

$$\begin{aligned} f(A_\alpha^o) &= f[\cup \{U \in I^X : \tau(U) > \alpha\tau(A), U \subseteq A\}] \\ &= f[\cup \{U \in I^X : \sigma(f(U)) > \alpha\sigma(f(A)), f(U) \subseteq f(A)\}] \\ &= \cup \{f(U) \in I^Y : \sigma(f(U)) > \alpha\sigma(f(A)), f(U) \subseteq f(A)\} \\ &= \cup \{K \in I^Y : \sigma(K) > \alpha\sigma(f(A)), K \subseteq f(A)\} \\ &= (f(A))_\alpha^o. \end{aligned} \quad (3.8) \quad \square$$

4. Types of smooth α -compactness. In this section, we introduce the concepts of several types of smooth α -compactness in smooth topological spaces and investigate some properties of them.

DEFINITION 4.1 [5]. An s.t.s. (X, τ) is called smooth compact if and only if for every family $\{A_i : i \in J\}$ in $\{A \in I^X : \tau(A) > 0\}$ covering X , there exists a finite subset J_0 of J such that $\cup_{i \in J_0} A_i = 1_X$.

THEOREM 4.2 [4]. *Let (X, τ) and (Y, σ) be two smooth topological spaces and $f : X \rightarrow Y$ a surjective weakly smooth continuous function with respect to τ and σ . If (X, τ) is smooth compact, then so is (Y, σ) .*

DEFINITION 4.3. Let $\alpha \in [0, 1)$. An s.t.s. (X, τ) is called smooth nearly α -compact if and only if for every family $\{A_i : i \in J\}$ in $\{A \in I^X : \tau(A) > 0\}$ covering X , there exists a finite subset J_0 of J such that $\cup_{i \in J_0} (\overline{A_i})_\alpha^o = 1_X$.

DEFINITION 4.4. Let $\alpha \in [0, 1)$. An s.t.s. (X, τ) is called smooth almost α -compact if and only if for every family $\{A_i : i \in J\}$ in $\{A \in I^X : \tau(A) > 0\}$ covering X , there exists a finite subset J_0 of J such that $\cup_{i \in J_0} \overline{A_i} = 1_X$.

DEFINITION 4.5. Let $\alpha \in [0, 1)$. An s.t.s. (X, τ) is called smooth α -regular if and only if each fuzzy set $A \in I^X$ satisfying $\tau(A) > 0$ can be written as $A = \cup \{K \in I^X : \tau(K) \geq \tau(A), \overline{K}_\alpha \subseteq A\}$.

DEFINITION 4.6. A smooth topology $\tau : I^X \rightarrow I$ on X is called monotonic increasing (resp., monotonic decreasing) if and only if $A \subseteq B \Rightarrow \tau(A) \leq \tau(B)$ (resp., $A \subseteq B \Rightarrow \tau(A) \geq \tau(B)$) for every $A, B \in I^X$.

THEOREM 4.7. *Let (X, τ) be an s.t.s., $\alpha \in [0, 1)$, and τ a monotonic decreasing smooth topology. If (X, τ) is smooth compact, then (X, τ) is smooth nearly α -compact.*

PROOF. Let (X, τ) be a smooth compact s.t.s. Then for every family $\{A_i : i \in J\}$ in $\{A \in I^X : \tau(A) > 0\}$ covering X , there exists a finite subset J_0 of J such

that $\cup_{i \in J_0} A_i = 1_X$. Since $\tau(A_i) > 0$ for each $i \in J$, we have $A_i = (A_i)_\alpha^0$ for each $i \in J$ by [Theorem 3.6](#). Since τ is monotonic decreasing and $A_i \subseteq \overline{(A_i)_\alpha}$ for each $i \in J$, we have $\tau(A_i) \geq \tau(\overline{(A_i)_\alpha})$ for each $i \in J$. Hence from [Theorem 3.2](#), we have $A_i = (A_i)_\alpha^0 \subseteq \overline{((A_i)_\alpha)_\alpha^0}$ for each $i \in J$. Thus $1_X = \cup_{i \in J_0} A_i \subseteq \cup_{i \in J_0} \overline{((A_i)_\alpha)_\alpha^0}$, that is, $\cup_{i \in J_0} \overline{((A_i)_\alpha)_\alpha^0} = 1_X$. Hence (X, τ) is smooth nearly α -compact. $\square$

THEOREM 4.8. *Let $\alpha \in [0, 1)$. Then a smooth nearly α -compact s.t.s. (X, τ) is smooth almost α -compact.*

PROOF. Let (X, τ) be a smooth nearly α -compact s.t.s. Then for every family $\{A_i : i \in J\}$ in $\{A \in I^X : \tau(A) > 0\}$ covering X , there exists a finite subset J_0 of J such that $\cup_{i \in J_0} \overline{((A_i)_\alpha)_\alpha^0} = 1_X$. Since $\overline{((A_i)_\alpha)_\alpha^0} \subseteq \overline{(A_i)_\alpha}$ for each $i \in J$ by [Theorem 3.5](#), $1_X = \cup_{i \in J_0} \overline{((A_i)_\alpha)_\alpha^0} \subseteq \cup_{i \in J_0} \overline{(A_i)_\alpha}$. So $\cup_{i \in J_0} \overline{(A_i)_\alpha} = 1_X$. Hence (X, τ) is smooth almost α -compact. $\square$

THEOREM 4.9. *Let (X, τ) and (Y, σ) be two smooth topological spaces, $\alpha \in [0, 1)$, and $f : X \rightarrow Y$ a surjective, weakly smooth continuous, and strict smooth α -preserving function with respect to τ and σ . If (X, τ) is smooth almost α -compact, then so is (Y, σ) .*

PROOF. Let $\{A_i : i \in J\}$ be a family in $\{A \in I^Y : \sigma(A) > 0\}$ covering Y , that is, $\cup_{i \in J} A_i = 1_Y$. Then $1_X = f^{-1}(1_Y) = \cup_{i \in J} f^{-1}(A_i)$. Since f is weakly smooth continuous, $\tau(f^{-1}(A_i)) > 0$ for each $i \in J$. Since (X, τ) is smooth almost α -compact, there exists a finite subset J_0 of J such that $\cup_{i \in J_0} \overline{(f^{-1}(A_i))_\alpha} = 1_X$. From the surjectivity of f we have $1_Y = f(1_X) = f(\cup_{i \in J_0} \overline{(f^{-1}(A_i))_\alpha}) = \cup_{i \in J_0} f(\overline{(f^{-1}(A_i))_\alpha})$. Since $f : X \rightarrow Y$ is strict smooth α -preserving with respect to τ and σ , from [Theorem 3.13](#) we have $\overline{(f^{-1}(A_i))_\alpha} \subseteq f^{-1}(\overline{(A_i)_\alpha})$ for each $i \in J$. Hence we have $1_Y = \cup_{i \in J_0} f(\overline{(f^{-1}(A_i))_\alpha}) \subseteq \cup_{i \in J_0} f(f^{-1}(\overline{(A_i)_\alpha})) = \cup_{i \in J_0} \overline{(A_i)_\alpha}$, that is, $\cup_{i \in J_0} \overline{(A_i)_\alpha} = 1_Y$. Thus (Y, σ) is smooth almost α -compact. $\square$

We obtain the following corollary from [Theorems 4.8](#) and [4.9](#).

COROLLARY 4.10. *Let (X, τ) and (Y, σ) be two smooth topological spaces, $\alpha \in [0, 1)$, and $f : X \rightarrow Y$ a surjective, weakly smooth continuous, and strict smooth α -preserving function with respect to τ and σ . If (X, τ) is smooth nearly α -compact, then (Y, σ) is smooth almost α -compact.*

THEOREM 4.11. *Let (X, τ) and (Y, σ) be two smooth topological spaces, $\alpha \in [0, 1)$, and $f : X \rightarrow Y$ a surjective, weakly smooth continuous, strict smooth α -preserving, and strict smooth open α -preserving function with respect to τ and σ . If (X, τ) is smooth nearly α -compact, then so is (Y, σ) .*

PROOF. Let $\{A_i : i \in J\}$ be a family in $\{A \in I^Y : \sigma(A) > 0\}$ covering Y , that is, $\cup_{i \in J} A_i = 1_Y$. Then $1_X = f^{-1}(1_Y) = \cup_{i \in J} f^{-1}(A_i)$. Since f is weakly smooth continuous, $\tau(f^{-1}(A_i)) > 0$ for each $i \in J$. Since (X, τ) is smooth nearly α -compact, there exists a finite subset J_0 of J such that $\cup_{i \in J_0} \overline{(f^{-1}(A_i))_\alpha} = 1_X$.

From the surjectivity of f we have $1_Y = f(1_X) = f(\cup_{i \in J_0} (\overline{(f^{-1}(A_i))_\alpha})_\alpha^o) = \cup_{i \in J_0} f((\overline{(f^{-1}(A_i))_\alpha})_\alpha^o)$. Since $f : X \rightarrow Y$ is strict smooth open α -preserving with respect to τ and σ , from Theorem 3.14 we have $f((\overline{(f^{-1}(A_i))_\alpha})_\alpha^o) \subseteq (f(\overline{(f^{-1}(A_i))_\alpha}))_\alpha^o$ for each $i \in J$. Since $f : X \rightarrow Y$ is strict smooth α -preserving with respect to τ and σ , from Theorem 3.13 we have $(f^{-1}(A_i))_\alpha \subseteq f^{-1}(\overline{(A_i)_\alpha})$ for each $i \in J$. Hence, we have

$$\begin{aligned} 1_Y &= \cup_{i \in J_0} f((\overline{(f^{-1}(A_i))_\alpha})_\alpha^o) \\ &\subseteq \cup_{i \in J_0} (f(\overline{(f^{-1}(A_i))_\alpha}))_\alpha^o \\ &\subseteq \cup_{i \in J_0} (f(f^{-1}(\overline{(A_i)_\alpha})))_\alpha^o \\ &= \cup_{i \in J_0} (\overline{(A_i)_\alpha})_\alpha^o. \end{aligned} \tag{4.1}$$

Hence, $\cup_{i \in J_0} (\overline{(A_i)_\alpha})_\alpha^o = 1_Y$. Thus (Y, σ) is smooth nearly α -compact. $\square$

THEOREM 4.12. *Let $\alpha \in [0, 1)$. Then a smooth almost α -compact smooth α -regular s.t.s. (X, τ) is smooth compact.*

PROOF. Let $\{A_i : i \in J\}$ be a family in $\{A \in I^X : \sigma(A) > 0\}$ covering X , that is, $\cup_{i \in J} A_i = 1_X$. Since (X, τ) is smooth α -regular, $A_i = \cup_{j_i \in J_i} \{K_{j_i} \in I^X : \tau(K_{j_i}) \geq \tau(A_i), \overline{(K_{j_i})_\alpha} \subseteq A_i\}$ for each $i \in J$. Since $\cup_{i \in J} A_i = \cup_{i \in J} [\cup_{j_i \in J_i} K_{j_i}] = 1_X$ and (X, τ) is smooth almost α -compact, there exists a finite subfamily $\{K_l \in I^X : \tau(K_l) > 0, l \in L\}$ such that $\cup_{l \in L} \overline{(K_l)_\alpha} = 1_X$. Since for each $l \in L$ there exists $i \in J$ such that $\overline{(K_l)_\alpha} \subseteq A_i$, we have $\cup_{i \in J_0} A_i = 1_X$, where J_0 is a finite subset of J . Hence (X, τ) is smooth compact. $\square$

We obtain the following corollary from Theorems 4.8 and 4.12.

COROLLARY 4.13. *Let $\alpha \in [0, 1)$. Then a smooth nearly α -compact smooth α -regular s.t.s. (X, τ) is smooth compact.*

REFERENCES

- [1] R. Badard, *Smooth axiomatics*, First IFSA Congress, Palma de Mallorca, 1986.
- [2] C. L. Chang, *Fuzzy topological spaces*, J. Math. Anal. Appl. **24** (1968), 182-190.
- [3] M. Demirci, *On several types of compactness in smooth topological spaces*, Fuzzy Sets and Systems **90** (1997), no. 1, 83-88.
- [4] ———, *Three topological structures of smooth topological spaces*, Fuzzy Sets and Systems **101** (1999), no. 1, 185-190.
- [5] M. K. El Gayyar, E. E. Kerre, and A. A. Ramadan, *Almost compactness and near compactness in smooth topological spaces*, Fuzzy Sets and Systems **62** (1994), no. 2, 193-202.
- [6] A. A. Ramadan, *Smooth topological spaces*, Fuzzy Sets and Systems **48** (1992), no. 3, 371-375.

Chun-Kee Park: Department of Mathematics, Kangwon National University, Chuncheon 200-701, Korea

E-mail address: ckpark@kangwon.ac.kr

Won Keun Min: Department of Mathematics, Kangwon National University, Chuncheon 200-701, Korea

E-mail address: wkmin@cc.kangwon.ac.kr

Myeong Hwan Kim: Department of Mathematics, Kangwon National University, Chuncheon 200-701, Korea

E-mail address: kimmw@kangwon.ac.kr