ON A CLASS OF HOLOMORPHIC FUNCTIONS DEFINED BY THE RUSCHEWEYH DERIVATIVE

GEORGIA IRINA OROS

Received 23 September 2002

By using the Ruscheweyh operator $D^m f(z)$, $z \in U$, we will introduce a class of holomorphic functions, denoted by $M_n^m(\alpha)$, and obtain some inclusion relations.

2000 Mathematics Subject Classification: 30C45.

1. Introduction and preliminaries. Denote by U the unit disc of the complex plane

$$U = \{z \in \mathbb{C}; |z| < 1\}. \quad (1.1)$$

Let $\mathcal{H}(U)$ be the space of holomorphic functions in U .

We let

$$A_n = \{f \in \mathcal{H}(U), f(z) = z + a_{n+1}z^{n+1} + \dots, z_1 \in U\} \quad (1.2)$$

with $A_1 = A$.

We let $\mathcal{H}[a, n]$ denote the class of analytic functions in U of the form

$$f(z) = a + a_n z^n + a_{n+1} z^{n+1} + \dots, \quad z \in U. \quad (1.3)$$

If f and g are analytic in U , we say that f is subordinate to g , written $f < g$ or $f(z) < g(z)$, if there is a function w analytic in U , with $w(0) = 0$, $|w(z)| < 1$, for any $z \in U$, such that $f(z) = g(w(z))$, for $z \in U$.

If g is univalent, then $f < g$ if and only if $f(0) = g(0)$ and $f(U) \subset g(U)$.

Let $K = \{f \in A : \operatorname{Re}(zf''(z)/f'(z)) + 1 > 0, z \in U\}$ denote the class of normalized convex functions in U . We use the following subordination results.

LEMMA 1.1 (Miller and Mocanu [2, page 71]). *Let h be a convex function with $h(0) = a$ and let $\gamma \in \mathbb{C}^*$ be a complex with $\operatorname{Re} \gamma \geq 0$. If $p \in \mathcal{H}[a, n]$ and*

$$p(z) + \frac{1}{\gamma} z p'(z) < h(z), \quad (1.4)$$

then $p(z) \prec g(z) \prec h(z)$, where

$$g(z) = \frac{\gamma}{nz^{\gamma/n}} \int_0^z h(t) \cdot t^{(\gamma/n)-1} dt. \quad (1.5)$$

The function g is convex and is the best (a, n) dominant.

LEMMA 1.2 (Miller and Mocanu [1]). *Let g be a convex function in U and let*

$$h(z) = g(z) + n\alpha z g'(z), \quad (1.6)$$

where $\alpha > 0$ and n is a positive integer. If $p(z) = g(0) + p_n z^n + \dots$ is holomorphic in U and

$$p(z) + \alpha z p'(z) \prec h(z), \quad (1.7)$$

then

$$p(z) \prec g(z) \quad (1.8)$$

and this result is sharp.

DEFINITION 1.3 [4]. For $f \in A$ and $m \geq 0$, the operator $D^m f$ is defined by

$$D^m f(z) = f(z) * \frac{z}{(1-z)^{m+1}} = \frac{z}{m!} [z^{m-1} f(z)]^{(m)}, \quad z \in U, \quad (1.9)$$

where $*$ stands for convolution.

REMARK 1.4. We have

$$\begin{aligned} D^0 f(z) &= f(z), \quad z \in U, \\ D^1 f(z) &= z f'(z), \quad z \in U, \\ 2D^2 f(z) &= z \cdot [D^1 f(z)]' + D^1 f(z), \\ (m+1)D^{m+1} f(z) &= z[D^m f(z)]' + mD^m f(z). \end{aligned} \quad (1.10)$$

2. Main results

DEFINITION 2.1. If $\alpha < 1$ and $m, n \in \mathbb{N}$, let $M_n^m(\alpha)$ denote the class of functions $f \in A_n$ which satisfy the inequality

$$\operatorname{Re} (D^m f)'(z) > \alpha. \quad (2.1)$$

THEOREM 2.2. *If $\alpha < 1$ and $m, n \in \mathbb{N}$, then*

$$M_n^{m+1}(\alpha) \subset M_n^m(\delta), \quad (2.2)$$

where

$$\delta = \delta(\alpha, n, m) = 2\alpha - 1 + 2 \cdot (1 - \alpha) \cdot \frac{m+1}{n} \beta\left(\frac{m+1}{n}\right), \quad (2.3)$$

$$\beta(x) = \int_0^1 \frac{t^{x-1}}{1+t} dt.$$

PROOF. Let $f \in M_n^{m+1}(\alpha)$. By using the properties of the operator $D^m f(z)$, we have

$$(m+1)D^{m+1}f(z) = z \cdot (D^m f)'(z) + mD^m f(z), \quad z \in U. \quad (2.4)$$

Differentiating (2.4), we obtain

$$\begin{aligned} (m+1)[D^{m+1}f(z)]' &= z \cdot (D^m f)''(z) + (D^m f)'(z) + m(D^m f)'(z) \\ &= z(D^m f)''(z) + (m+1)(D^m f)'(z). \end{aligned} \quad (2.5)$$

If we let $p(z) = (D^m f)'(z)$, then $p'(z) = (D^m f)''(z)$ and (2.4) becomes

$$[D^{m+1}f(z)]' = p(z) + \frac{1}{m+1} z \cdot p'(z). \quad (2.6)$$

Since $f \in M_n^{m+1}(\alpha)$, by using Definition 2.1, we have

$$\operatorname{Re} \left[p(z) + \frac{1}{m+1} z p'(z) \right] > \alpha \quad (2.7)$$

which is equivalent to

$$p(z) + \frac{1}{m+1} z p'(z) \prec \frac{1 + (2\alpha - 1)z}{1 + z} \equiv h(z). \quad (2.8)$$

By using Lemma 1.1, we have

$$p(z) \prec g(z) \prec h(z), \quad (2.9)$$

where

$$g(z) = \frac{m+1}{nz^{(m+1)/n}} \int_0^z \frac{1 + (2\alpha - 1)t}{1+t} \cdot t^{(m+1)/n-1} dt. \quad (2.10)$$

The function g is convex and is the best dominant.

From $p(z) \prec g(z)$, it results that

$$\operatorname{Re} p(z) > \delta = g(1) = \delta(\alpha, n, m), \quad (2.11)$$

where

$$\begin{aligned} g(1) &= \frac{m+1}{n} \int_0^1 t^{(m+1)/n-1} \cdot \frac{1+(2\alpha-1)t}{1+t} dt \\ &= 2\alpha - 1 + 2 \cdot \frac{m+1}{n} \cdot (1-\alpha) \beta\left(\frac{m+1}{n}\right), \end{aligned} \quad (2.12)$$

from which we deduce that $M_n^{m+1}(\alpha) \subset M_n^m(\delta)$. $\square$

For $n = 1$, this result was obtained in [3].

THEOREM 2.3. *Let g be a convex function, $g(0) = 1$, and let h be a function such that*

$$h(z) = g(z) + \frac{1}{m+1} z g'(z). \quad (2.13)$$

If $f \in A_n$ and verifies the differential subordination

$$(D^{m+1}f)'(z) \prec h(z), \quad (2.14)$$

then

$$(D^m f)'(z) \prec g(z). \quad (2.15)$$

PROOF. From

$$(m+1)D^{m+1}f(z) = z \cdot (D^m f)'(z) + mD^m f(z), \quad (2.16)$$

we obtain

$$\begin{aligned} (m+1)[D^{m+1}f(z)]' &= (D^m f)'(z) + z(D^m f)''(z) + m(D^m f)'(z) \\ &= z(D^m f)''(z) + (m+1)(D^m f)'(z). \end{aligned} \quad (2.17)$$

If we let $p(z) = (D^m f)'(z)$, then we obtain

$$[D^{m+1}f(z)]' = p(z) + \frac{1}{m+1} z p'(z) \quad (2.18)$$

and (2.14) becomes

$$p(z) + \frac{1}{m+1} z p'(z) \prec g(z) + \frac{1}{m+1} z g'(z) \equiv h(z). \quad (2.19)$$

By using Lemma 1.2, we have

$$p(z) \prec g(z), \quad \text{i.e., } (D^m f)'(z) \prec g(z). \quad (2.20)$$

$\square$

For $n = 1$, this result was obtained in [3].

THEOREM 2.4. Let $h \in \mathcal{H}[U]$, with $h(0) = 1$, $h'(0) \neq 0$, which verifies the inequality

$$\operatorname{Re} \left[1 + \frac{zh''(z)}{h'(z)} \right] > -\frac{1}{2(m+1)}, \quad m \geq 0. \quad (2.21)$$

If $f \in A_n$ and verifies the differential subordination

$$[D^{m+1}f(z)]' \prec h(z), \quad z \in U, \quad (2.22)$$

then

$$[D^m f(z)]' \prec g(z), \quad (2.23)$$

where

$$g(z) = \frac{m+1}{nz^{(m+1)/n}} \int_0^z h(t)t^{(m+1)/n-1} dt. \quad (2.24)$$

The function g is convex and is the best dominant.

PROOF. A simple application of the differential subordination technique [1, 2] shows that the function g is convex. From

$$(m+1)D^{m+1}f(z) = z[D^m f(z)]' + mD^m f(z), \quad (2.25)$$

we obtain

$$(m+1)[D^{m+1}f(z)]' = z[D^m f(z)]'' + (m+1)[D^m f(z)]'. \quad (2.26)$$

If we let $p(z) = [D^m f(z)]'$, then we obtain

$$[D^{m+1}f(z)]' = p(z) + \frac{1}{m+1}zp'(z) \quad (2.27)$$

and (2.22) becomes

$$p(z) + \frac{1}{m+1}zp'(z) \prec h(z). \quad (2.28)$$

By using Lemma 1.1, we have

$$p(z) \prec g(z) = \frac{m+1}{nz^{(m+1)/n}} \int_0^z h(t)t^{(m+1)/n-1} dt. \quad (2.29)$$

□

THEOREM 2.5. Let g be a convex function, $g(0) = 1$, and

$$h(z) = g(z) + n zg'(z). \quad (2.30)$$

If $f \in A_n$ and verifies the differential subordination

$$[D^m f(z)]' \prec h(z), \quad z \in U, \quad (2.31)$$

then

$$\frac{D^m f(z)}{z} \prec g(z). \quad (2.32)$$

PROOF. We let $p(z) = D^m f(z)/z$, $z \in U$, and we obtain

$$D^m f(z) = zp(z). \quad (2.33)$$

By differentiating, we obtain

$$[D^m f(z)]' = p(z) + zp'(z), \quad z \in U. \quad (2.34)$$

Then (2.31) becomes

$$p(z) + zp'(z) \prec h(z) = g(z) + zg'(z). \quad (2.35)$$

By using Lemma 1.2, we have (1.8). $\square$

REFERENCES

- [1] S. S. Miller and P. T. Mocanu, *On some classes of first-order differential subordinations*, Michigan Math. J. **32** (1985), no. 2, 185–195.
- [2] ———, *Differential Subordinations. Theory and Applications*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 225, Marcel Dekker, New York, 2000.
- [3] Gh. Oros and G. I. Oros, *A class of holomorphic functions*, Automat. Comput. Appl. Math. **11** (2002), no. 2, 77–80.
- [4] St. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc. **49** (1975), 109–115.

Georgia Irina Oros: Faculty of Mathematics and Computer Science, Babeş-Bolyai University, 400084 Cluj-Napoca, Romania

E-mail address: georgia_oros_ro@yahoo.com