A GENERAL LEMMA FOR FIXED-POINT THEOREMS INVOLVING MORE THAN TWO MAPS IN D -METRIC SPACES WITH APPLICATIONS

B. C. DHAGE, SMRATI ARYA, and JEONG SHEOK UME

Received 26 June 2002

A general procedural lemma for fixed-point theorems for three and four maps in a D -metric space is proved, and it is further applied for proving the common fixed-point theorems of three and four maps in a D -metric space satisfying certain contractive conditions.

2000 Mathematics Subject Classification: 47H10.

1. Introduction. The concept of a D -metric space is introduced by the first author in [2]. A nonempty set X , together with a function $\rho : X \times X \times X \rightarrow [0, \infty)$, is called a D -metric space with D -metric ρ if ρ satisfies the following properties:

- (i) $\rho(x, y, z) = 0 \Leftrightarrow x = y = z$ (coincidence),
- (ii) $\rho(x, y, z) = \rho(p\{x, z, y\})$ (symmetry), where p is a permutation function,
- (iii) $\rho(x, y, z) \leq \rho(x, y, a) + \rho(x, a, z) + \rho(a, y, z)$ for all $x, y, z, a \in X$ (tetrahedral inequality).

A few details along with some specific examples of a D -metric space appear in [3].

A sequence $\{x_n\} \subset X$ is said to be convergent to a point $x \in X$ if

$$\lim_{m, n \rightarrow \infty} \rho(x_m, x_n, x) = 0. \quad (1.1)$$

Similarly, a sequence $\{x_n\} \subset X$ is called D -Cauchy if

$$\lim_{m, n, p \rightarrow \infty} \rho(x_m, x_n, x_p) = 0. \quad (1.2)$$

A complete D -metric space is one in which every D -Cauchy sequence converges to a point. Further, a subset S of a D -metric space X is called bounded if there exists a constant $k > 0$ such that $\rho(x, y, z) \leq k$ for all $x, y, z \in S$, and the constant k is called a D -bound of S . The infimum of all such D -bounds k is called the diameter of S , and it is denoted by $\delta(S)$. Finally, it is known that a

mapping $f : X \rightarrow X$ is continuous if and only if, for any sequence $\{x_n\} \subset X$, $x_n \rightarrow x^*$ implies $fx_n \rightarrow fx^*$.

It has been shown in [5] that the D -metric ρ is continuous on X^3 in the topology of D -metric convergence which is Hausdorff. For details of a D -metric space, the reader is referred to Dhage [5].

The general existence principles for the fixed-point theorems for a single and a pair of maps in D -metric space have been established in Dhage and Rhoades [6] and Dhage [1], respectively. However, the extension of these existence principles to three or four maps is not possible.

In this paper, we just formulate the general procedure for common fixed-point theorems for more than two selfmaps of a D -metric space and discuss some of its applications.

We need the following auxiliary results in the sequence.

PROPOSITION 1.1. *Let $\{x_n\}$ be a sequence in a D -metric space X satisfying*

$$\rho(x_n, x_{n+1}, z) \leq \lambda \rho(x_{n-1}, x_n, z) \quad (1.3)$$

for all $n \in \mathbb{N}$ and $z \in \{x_n\}$, where $0 \leq \lambda < 1$. Then,

$$\rho(x_n, x_{n+1}, x_m) \leq \lambda^n k \quad (1.4)$$

for all $m > n$, where $k = (2/(1-\lambda)) \max\{\rho(x_0, x_0, x_1), \rho(x_0, x_1, x_1)\}$.

PROOF. From (1.3),

$$\rho(x_n, x_{n+1}, x_m) \leq \lambda \rho(x_{n-1}, x_n, x_m) \quad (1.5)$$

for each $m > n \in \mathbb{N}$. By induction,

$$\rho(x_n, x_{n+1}, x_m) \leq \lambda^n \rho(x_0, x_1, x_m). \quad (1.6)$$

Let $q = \max\{\rho(x_0, x_0, x_1), \rho(x_0, x_1, x_1)\}$. Using the tetrahedral inequality,

$$\begin{aligned} \rho(x_0, x_1, x_m) &\leq \rho(x_0, x_1, x_{m-1}) + \rho(x_{m-1}, x_0, x_m) + \rho(x_{m-1}, x_m, x_1) \\ &= \rho(x_0, x_1, x_{m-1}) + \rho(x_{m-1}, x_m, x_0) + \rho(x_{m-1}, x_m, x_1) \\ &\leq \rho(x_0, x_1, x_{m-1}) + \lambda^{m-1} \rho(x_0, x_1, x_0) + \lambda^{m-1} \rho(x_0, x_1, x_1) \\ &\leq \rho(x_0, x_1, x_{m-1}) + 2\lambda^{m-1} q \\ &\leq \rho(x_0, x_1, x_{m-2}) + 2\lambda^{m-1} q + 2\lambda^{m-2} q \\ &\vdots \\ &\leq \rho(x_0, x_1, x_1) + 2(\lambda + \lambda^2 + \cdots + \lambda^{m-1}) q \\ &\leq 2(1 + \lambda + \lambda^2 + \cdots + \lambda^{m-1}) q < \frac{2}{1-\lambda} q = k. \end{aligned} \quad (1.7)$$

Substituting (1.7) into (1.6) yields the desired inequality (1.4). $\square$

PROPOSITION 1.2. Every sequence $\{x_n\} \subset X$ satisfying (1.3) is bounded with a D -bound $k = (2/(1-\lambda)) \max\{\rho(x_0, x_0, x_1), \rho(x_0, x_1, x_1)\}$.

PROOF. Let $q = \max\{\rho(x_0, x_0, x_1), \rho(x_0, x_1, x_1)\}$. Then, for any integers $r \geq s \geq n$, there exists positive integer p and t such that

$$\begin{aligned}
 \rho(x_n, x_r, x_s) &= \rho(x_n, x_{n+p}, x_{n+t}) \\
 &\leq \rho(x_n, x_{n+1}, x_{n+t}) + \rho(x_n, x_{n+1}, x_{n+p}) + \rho(x_{n+1}, x_{n+p}, x_{n+t}) \\
 &\leq 2\lambda^n q + \rho(x_{n+1}, x_{n+p}, x_{n+q}) \\
 &\leq 2\lambda^n q + 2\lambda^{n+1} q + \rho(x_{n+2}, x_{n+p}, x_{n+t}) \\
 &\vdots \\
 &\leq 2 \left(\sum_{j=1}^{n+p-2} \lambda^j \right) q + \rho(x_{n+p-1}, x_{n+p}, x_{n+r}) \\
 &\leq 2 \left(\sum_{j=1}^{n+p-1} \lambda^j \right) q < 2 \left(\sum_{j=1}^{\infty} \lambda^j \right) q < \frac{2}{1-\lambda} q = k.
 \end{aligned} \tag{1.8}$$

Then, $\{x_n\}$ is bounded and the proof is complete. $\square$

2. Main results. Before going to the main results of this paper, we state a lemma proved in Dhage [4].

LEMMA 2.1 (D -Cauchy principle). Let $\{y_n\}$ be a bounded sequence in D -metric space with D -bound k satisfying

$$\rho(y_n, y_{n+1}, y_m) \leq \lambda^n k \tag{2.1}$$

for all $m > n \in \mathbb{N}$. Then, $\{y_n\}$ is D -Cauchy.

Let $A, B, S, T : X \rightarrow X$ be four maps such that

$$A(X) \subseteq T(X), \quad B(X) \subseteq S(X). \tag{2.2}$$

Condition (2.2) ensures that it is possible to define a sequence $\{y_n\}$ in X as follows. Let $x \in X$ be arbitrary. Then, in view of condition (2.2), there exists a sequence $\{x_n\}$ such that

$$x_0 = x, \quad Ax_{2n} = Tx_{2n+1}, \quad Bx_{2n+1} = Sx_{2n+2}, \quad n \geq 0. \tag{2.3}$$

Now, define $\{y_n\}$ in X by

$$y_0 = Sx_0, \quad y_{2n+1} = Tx_{2n+1}, \quad y_{2n+2} = Sx_{2n+2}, \quad n \geq 0. \tag{2.4}$$

A point $x \in X$ is called a coincidence point of two maps $A, B : X \rightarrow X$ if $Ax = Bx$, and in this case, the mappings A and B are called coincident on X .

Similarly, a coincidence point of three or four maps on a D -metric space is defined.

LEMMA 2.2. *Let $A, B, S, T : X \rightarrow X$ satisfy (2.2), and let $\{y_n\} \subset X$ be defined by (2.4). Further, assume that $\{y_n\}$ is complete. Suppose that there exists a $\lambda \in [0, 1)$ such that*

$$\rho(y_n, y_{n+1}, z) \leq \lambda \rho(y_{n-1}, y_n, z) \quad (2.5)$$

for all $n \in \mathbb{N}$ and $z \in \{y_n\}$. Then, either

- (a) A and S have a coincidence point,
- (b) B and T have a coincidence point,
- (c) A, S , and T have a coincidence point,
- (d) B, S , and T have a coincidence point, or
- (e) $\{y_n\}$ converges to a point $u \in X$ and, for all $m > n \in \mathbb{N}$,

$$\rho(y_n, y_m, u) \leq 2 \sum_{j=n}^m \lambda^j k, \quad \rho(y_n, u, u) \leq 2 \frac{\lambda^n}{1-\lambda} k, \quad (2.6)$$

where $k = \delta(\{y_n\})$.

PROOF. Suppose that $y_{2n} = y_{2n+1}$ for some n . Then, $Sx_{2n} = Tx_{2n+1} = Ax_{2n}$ and (a) holds. Also, if $x_{2n} = x_{2n+1}$, then $Tx_{2n} = Tx_{2n+1}$ and so (c) holds. Similarly, if $y_{2n+1} = y_{2n+2}$ for some n , then it is shown analogously that (b) and (d) hold.

Suppose now that $y_n \neq y_{n+1}$ for each n . Then, from Proposition 1.1, it follows that

$$\rho(y_n, y_{n+1}, y_m) \leq \lambda^n \rho(y_0, y_1, y_m) \leq \lambda^n k \quad (2.7)$$

for all $m > n \in \mathbb{N}$. Now, an application of Lemma 2.1 yields that $\{y_n\}$ is D -Cauchy. Since $\{y_n\}$ is complete, there exists a point $u \in X$ such that $\lim_n y_n = u$.

Now, for any positive integers m and n , $m > n$, by repeated application of the tetrahedral inequality,

$$\begin{aligned} \rho(y_n, y_m, u) &\leq \rho(y_n, y_{n+1}, y_m) + \rho(y_n, y_{n+1}, u) + \rho(y_{n+1}, y_m, u) \\ &\leq \lambda^n \rho(y_0, y_1, y_m) + \lambda^n \rho(y_0, y_1, u) + \rho(y_{n+1}, y_m, u) \\ &\leq 2\lambda^n k + \rho(y_{n+1}, y_m, u) \\ &\leq 2\lambda^n k + 2\lambda^{n+1} k + \rho(y_{n+2}, y_m, u) \\ &\vdots \\ &\leq 2(\lambda^n + \lambda^{n+1} + \dots + \lambda^m) k \\ &= 2 \sum_{j=n}^m \lambda^j k. \end{aligned} \quad (2.8)$$

The above inequality further gives that

$$\begin{aligned}\rho(y_n, y_m, u) &\leq 2 \sum_{j=n}^m \lambda^j k \\ &= 2\lambda^n (1 + \lambda + \cdots + \lambda^{m-n})k \\ &= 2\lambda^n \left(\frac{1 - \lambda^{m-n}}{1 - \lambda} \right) k.\end{aligned}\quad (2.9)$$

Taking the limit as $m \rightarrow \infty$ in the above inequality,

$$\rho(y_n, u, u) \leq 2 \frac{\lambda^n}{1 - \lambda} k. \quad (2.10)$$

The proof of [Lemma 2.2](#) is complete. $\square$

The three-maps version of [Lemma 2.2](#) is obtained in two ways: one by setting $S = T$ and the other by setting $A = B$. In the situation when $S = T$, condition [\(2.2\)](#) reduces to

$$A(X) \subseteq S(X), \quad B(X) \subseteq S(X). \quad (2.11)$$

Then, it is possible to choose a sequence $\{x_n\} \subset X$ such that

$$x_0 = x, \quad Ax_{2n} = Sx_{2n+1}, \quad Bx_{2n+1} = Sx_{2n+2}, \quad n \geq 0. \quad (2.12)$$

Now, define a sequence $\{y_n\}$ in X as follows:

$$y_0 = Sx_0, \quad y_{2n} = Sx_{2n}, \quad y_{2n+1} = Sx_{2n+1}, \quad n \in \mathbb{N}. \quad (2.13)$$

LEMMA 2.3. *Let $A, B, S : X \rightarrow X$ satisfy [\(2.11\)](#). Suppose that there exists an $x \in X$ such that the sequence $\{y_n\} \subset X$ defined by [\(2.13\)](#) is complete. Further, suppose that*

$$\rho(y_n, y_{n+1}, z) \leq \lambda \rho(y_{n-1}, y_n, z) \quad (2.14)$$

for all $n \in \mathbb{N}$ and $z \in \{y_n\}$, where $0 \leq \lambda < 1$. Then, either

- (a) A and S have a coincidence point,
- (b) B and S have a coincidence point, or
- (c) $\{y_n\}$ converges to a point $u \in X$ and, for all positive integers m and n , $m > n$,

$$\rho(y_n, y_m, u) \leq 2 \sum_{j=n}^m \lambda^j k, \quad \rho(y_n, u, u) \leq 2 \frac{\lambda^n}{1 - \lambda} k, \quad (2.15)$$

where $k = \delta(\{y_n\})$.

It is known that the fixed-point theorems for more than two maps require some sort of commutativity condition on the mappings under consideration.

Below, we will apply [Lemma 2.2](#) for proving the common fixed-point theorem for four maps on a D -metric space under a suitable commutativity condition.

A sequence $\{x_n\} \subset X$ is called a sequence of coincidence for the maps $A, B : X \rightarrow X$ if $\lim_n Ax_n = \lim_n Bx_n$. In this case, the mappings A and B are called limit coincident on X . Similarly, two maps $A, B : X \rightarrow X$ are called commuting or commutative if $(AB)(x) = (BA)(x)$ for all $x \in X$ and limit commuting if there exists a sequence $\{x_n\} \subset X$ such that

$$\lim_n (AB)(x_n) = \lim_n (BA)(x_n). \quad (2.16)$$

Finally, two maps $A, B : X \rightarrow X$ are called limit coincidentally commuting if their limit coincidence implies the limit commuting on X , that is, for any sequence $\{x_n\} \subset X$ if

$$\lim_n Ax_n = \lim_n Bx_n \Rightarrow \lim_n (AB)(x_n) = \lim_n (BA)(x_n). \quad (2.17)$$

It is known that the limit coincidentally commuting mappings commute at their coincidence points. See, for details, Dhage [4].

Now, we are ready to give some applications of [Lemma 2.2](#) for proving the existence of a common fixed point of four maps on a D -metric space X .

An orbit of four selfmaps A, B, S , and T of a D -metric space X at a point $x \in X$ is a set $O_{A,B}(S, T : x)$ in X defined by

$$O_{A,B}(S, T : x) = \{y_0 = Sx_0, y_{2n+1} = Ax_{2n} = Tx_{2n+1}, y_{2n+2} = Bx_{2n+1} = Sx_{2n+1} : n \geq 0\}. \quad (2.18)$$

Clearly, the orbit $O_{A,B}(S, T : x)$ is well defined if A, B, S , and T satisfy condition (2.2). By $\overline{O_{A,B}(S, T : x)}$, we denote the closure of the orbit $O_{A,B}(S, T : x)$ in X .

THEOREM 2.4. *Let $A, B, S, T : X \rightarrow X$ be four selfmaps of a D -metric space X satisfying (2.2) and*

$$\rho(Ax, By, z) \leq \lambda \max \{\rho(Sx, Ty, z), \rho(Sx, Ax, z), \rho(Ty, By, z)\} \quad (2.19)$$

for all $x, y, z \in X$, where $0 \leq \lambda < 1$. Assume further that

- (a) $\overline{O_{A,B}(S, T : x)}$ is complete for each $x \in X$,
- (b) $\{A, S\}$ and $\{B, T\}$ are limit coincidentally commuting,
- (c) any one A, B, S , or T continuous.

Then, A, B, S , and T have a unique common fixed point.

PROOF. Let $x \in X$ be arbitrary, and define a sequence $\{y_n\} \subset X$ by (2.4), which is possible in view of condition (2.4). Now, taking $x = x_{2n}$ and $y = x_{2n+1}$

in (2.19),

$$\begin{aligned} & \rho(y_{2n+1}, y_{2n+2}, z) \\ & \leq \lambda \max \{ \rho(y_{2n}, y_{2n+1}, z), \rho(y_{2n}, y_{2n+1}, z), \rho(y_{2n+1}, y_{2n+2}, z) \} \\ & = \lambda \rho(y_{2n}, y_{2n+1}, z) \end{aligned} \quad (2.20)$$

for all $n \geq 0$ and $z \in \{y_n\}$. Similarly, taking $x = x_{2n}$ and $y = x_{2n-1}$ in (2.19),

$$\begin{aligned} & \rho(y_{2n}, y_{2n+1}, z) \leq \lambda \max \{ \rho(y_{2n-1}, y_{2n}, z), \rho(y_{2n}, y_{2n+1}, z), \rho(y_{2n-1}, y_{2n}, z) \} \\ & = \lambda \rho(y_{2n-1}, y_{2n}, z) \end{aligned} \quad (2.21)$$

for all $n \in \mathbb{N}$ and $z \in \{y_n\}$. Hence, in general,

$$\rho(y_n, y_{n+1}, y_m) \leq \lambda \rho(y_{n-1}, y_n, y_m) \quad (2.22)$$

for all $m > n \in \mathbb{N}$ and $0 \leq \lambda < 1$.

We prove the conclusion of our theorem in two cases.

CASE 1. If $y_n = y_{n+1}$, then $y_n = y_{n+k}$ for all $k \geq 0$. If $y_{n+1} \neq y_{n+2}$, then, replacing n in (2.22) by $n+1$,

$$0 < \rho(y_{n+1}, y_{n+2}, y_{n+1}) \leq \lambda \rho(y_n, y_{n+1}, y_{n+1}) = 0, \quad (2.23)$$

which is a contradiction and $y_{n+1} = y_{n+2}$, and, by induction, $y_n = y_{n+k}$ for all $k \geq 0$. Therefore, by Lemma 2.2, there are points u and v in X such that $w_1 = Au = Su$ and $w_2 = Bv = Tv$.

We will show that $w_1 = w_2$. By (2.19),

$$\begin{aligned} & \rho(w_1, w_2, w_1) = \rho(Au, Bv, w_1) \\ & \leq \lambda \max \{ \rho(Su, Tv, w_1), \rho(Su, Au, w_1), \rho(Tv, Bv, w_1) \} \\ & = \lambda \max \{ \rho(w_1, w_2, w_1), \rho(w_1, w_1, w_1), \rho(w_2, w_2, w_1) \} \\ & = \lambda \rho(w_1, w_2, w_2). \end{aligned} \quad (2.24)$$

Again,

$$\begin{aligned} & \rho(w_1, w_2, w_2) = \rho(Au, Bv, w_2) \\ & \leq \lambda \max \{ \rho(Su, Tv, w_2), \rho(Su, Au, w_2), \rho(Tv, Bv, w_2) \} \\ & = \lambda \max \{ \rho(w_1, w_2, w_2), \rho(w_1, w_1, w_2), \rho(w_2, w_2, w_2) \} \\ & = \lambda \rho(w_1, w_2, w_1). \end{aligned} \quad (2.25)$$

Substituting (2.25) into (2.24),

$$\rho(w_1, w_2, w_1) \leq \lambda^2 \rho(w_1, w_2, w_1), \quad (2.26)$$

which is possible only when $w_1 = w_2$ since $\lambda < 1$. Hence, $Au = Bv = Su = Tv = w$. Next, we show that w is a coincidence of A, B, S , and T . Since $\{A, S\}$ and $\{B, T\}$ are limit coincidentally commuting, they commute at coincidence point. Therefore, $Sw = SAu = ASu = Aw$ and $Tw = TBv = BTv = Bw$. Now,

$$\begin{aligned}\rho(Aw, Bw, Aw) &\leq \lambda \max \{\rho(Sw, Tw, Aw), \rho(Sw, Aw, Aw), \rho(Tw, Bw, Aw)\} \\ &= \lambda \max \{\rho(Aw, Bw, Aw), \rho(Bw, Bw, Aw)\} \\ &= \lambda \rho(Aw, Bw, Bw).\end{aligned}\tag{2.27}$$

Similarly,

$$\rho(Aw, Bw, Bw) \leq \lambda \rho(Aw, Bw, Aw).\tag{2.28}$$

Substituting (2.28) into (2.27),

$$\rho(Aw, Bw, Aw) \leq \lambda^2 \rho(Aw, Bw, Aw),\tag{2.29}$$

which is possible only when $Aw = Bw$. Hence, $Aw = Sw = Tw = Bw$ is a coincidence point of the four maps A, B, S , and T . Finally, we prove that w is a common fixed point of A, B, S , and T . If $w \neq Aw$, then, by (2.19),

$$\begin{aligned}\rho(Aw, w, w) &= \rho(Aw, Bv, w) \\ &\leq \lambda \max \{\rho(Sw, Tv, w), \rho(Sw, Aw, w), \rho(Tv, Bv, w)\} \\ &= \lambda \rho(Aw, w, Aw).\end{aligned}\tag{2.30}$$

Similarly,

$$\rho(Aw, w, Aw) \leq \lambda \rho(Aw, w, w).\tag{2.31}$$

From (2.30) and (2.31),

$$\rho(Aw, w, w) \leq \lambda^2 \rho(Aw, w, w),\tag{2.32}$$

which is a contradiction to $Aw = w$ and hence, $w = Aw = Sw = Tw = Bw$.

CASE 2. Suppose that $y_n \neq y_{n+1}$ for each n . Then, by Lemma 2.2, there exists a point $w \in X$ such that $\lim_n y_n = w$. By definition of $\{y_n\}$,

$$\begin{aligned}\lim_n y_{2n} &= \lim_n Sx_{2n} = \lim_n Ax_{2n} = \lim_n Tx_{2n+1} = w, \\ \lim_n y_{2n+1} &= \lim_n Tx_{2n+1} = \lim_n Bx_{2n+1} = \lim_n y_{2n+2} = w.\end{aligned}\tag{2.33}$$

Since $\{A, S\}$ and $\{B, T\}$ are limit coincidentally commuting, then

$$\begin{aligned}\lim_n ASx_{2n} &= \lim_n SAX_{2n}, \\ \lim_n BTx_{2n+1} &= \lim_n TBx_{2n+1}.\end{aligned}\tag{2.34}$$

Suppose first that S is continuous on X . Then,

$$\lim_n SSx_{2n} = \lim_n SAx_{2n} = \lim_n ASx_{2n} = Sw. \quad (2.35)$$

First, we show that w is a fixed point of S . If $w \neq Sw$, then, by (2.19),

$$\begin{aligned} & \rho(Sw, w, w) \\ &= \lim_n \rho(ASx_{2n}, Bx_{2n+1}, w) \\ &\leq \lambda \lim_n \max \{ \rho(SSx_{2n}, Tx_{2n+1}, w), \rho(SBx_{2n}, ASx_{2n}, w), \rho(Tx_{2n+1}, Bx_{2n+1}, w) \} \\ &= \lambda \max \{ \rho(Sw, w, w), \rho(Sw, Sw, w) \} \\ &= \lambda \rho(Sw, Sw, w). \end{aligned} \quad (2.36)$$

Similarly,

$$\rho(Sw, w, Sw) \leq \lambda \rho(Sw, w, w). \quad (2.37)$$

Substituting (2.37) into (2.36),

$$\rho(Sw, w, w) \leq \lambda^2 \rho(Sw, w, w), \quad (2.38)$$

which is a contradiction and hence, $Sw = w$. Similarly,

$$\begin{aligned} & \rho(Aw, w, w) \\ &\leq \lim_n \rho(Aw, Bx_{2n+1}, w) \\ &\leq \lambda \lim_n \max \{ \rho(Sw, Tx_{2n+1}, w), \rho(Tx_{2n+1}, Bx_{2n+1}, w), \rho(Sw, Aw, w) \} \\ &= \lambda \max \{ 0, 0, \rho(Aw, w, w) \} \\ &= \lambda \rho(Aw, w, w), \end{aligned} \quad (2.39)$$

which implies that $Aw = w$, since $\lambda < 1$. From the condition $A(X) \subseteq T(X)$, it follows that there is a point $p \in X$ such that $w = Aw = Tp$. We show that $Bp = Tp$. If not, then

$$\begin{aligned} \rho(Tp, Bp, w) &= \rho(Aw, Bp, w) \\ &\leq \lambda \max \{ \rho(Sw, Tp, w), \rho(Sw, Aw, w), \rho(Tp, Bp, w) \} \\ &= \lambda \rho(Tp, Bp, w), \end{aligned} \quad (2.40)$$

which is a contradiction. Hence, $Bp = Tp$. Since $\{B, T\}$ are limit coincidentally commuting, we obtain $Bw = BTp = TBp = Bw$. Now,

$$\begin{aligned} \rho(Aw, Bw, w) &\leq \lambda \max \{ \rho(Sw, Tw, w), \rho(Sw, Aw, w), \rho(Tw, Bw, w) \} \\ &= \lambda \rho(Aw, Bw, Bw). \end{aligned} \quad (2.41)$$

Similarly,

$$\rho(Aw, Bw, Bw) \leq \lambda \rho(Aw, Bw, w). \quad (2.42)$$

Substituting (2.42) into (2.41),

$$\rho(Aw, Bw, w) \leq \lambda^2 \rho(Aw, Bw, w), \quad (2.43)$$

which is possible only when $Aw = Bw$. Thus, w is a common fixed point of A , B , S , and T .

Similarly, if T is continuous, then it is proved in an analogous way that A , B , S , and T have a common fixed point.

Next, suppose that A is continuous. Then, we have

$$\lim_n AAx_{2n} = \lim_n ASx_{2n} = \lim_n Sx_{2n} = Aw. \quad (2.44)$$

First, we show that $Aw = w$. If $Aw \neq w$, then

$$\begin{aligned} & \rho(Aw, w, w) \\ &= \lim_n \rho(AAx_{2n}, Bx_{2n+1}, w) \\ &\leq \lambda \lim_n \max \{ \rho(SAx_{2n}, Tx_{2n+1}, w), \rho(SAx_{2n}, AAx_{2n}, w), \rho(Tx_{2n+1}, Bx_{2n+1}, w) \} \\ &= \lambda \max \{ \rho(Aw, w, w), \rho(Aw, Aw, w) \} \\ &= \lambda \rho(Aw, Aw, w). \end{aligned} \quad (2.45)$$

Similarly,

$$\rho(Aw, w, Aw) \leq \lambda \rho(Aw, w, w). \quad (2.46)$$

Substituting (2.46) into (2.45),

$$\rho(Aw, w, w) \leq \lambda^2 \rho(Aw, w, w), \quad (2.47)$$

which is a contradiction. Hence, $Aw = w$. Using condition (2.2), there exists a point $p \in X$ such that $Tp = Aw = w$. We show that $Bp = Tp$. Now,

$$\begin{aligned} & \rho(Aw, Bp, w) \\ &= \lim_n \rho(AAx_{2n}, Bp, w) \\ &\leq \lambda \lim_n \max \{ \rho(SAx_{2n}, Tp, w), \rho(SAx_{2n}, AAx_{2n}, w), \rho(Tp, Bp, w) \} \\ &= \lambda \rho(w, Bp, w), \end{aligned} \quad (2.48)$$

which gives that $Bp = Tp$. Since $\{B, T\}$ are limit coincidentally commuting, they commute at coincidence point. Hence, $Tw = TBp = BTp = Bw$. Now,

$$\rho(Ax_{2n}, Bw, w) \leq \lambda \max \{ \rho(Sx_{2n}, Tw, w), \rho(Sx_{2n}, Ax_{2n}, w), \rho(Tw, Bw, w) \}. \quad (2.49)$$

Taking the limit as $n \rightarrow \infty$,

$$\rho(w, Bw, w) \leq \lambda \rho(w, Bw, Bw). \quad (2.50)$$

Similarly,

$$\rho(w, Bw, Bw) \leq \lambda \rho(w, Bw, w). \quad (2.51)$$

Substituting (2.51) into (2.50),

$$\rho(w, Bw, w) \leq \lambda^2 \rho(w, Bw, w), \quad (2.52)$$

which implies that $w = Bw = Tw = Aw$. Since $B(X) \subseteq S(X)$, there is a point $q \in X$ such that $w = Bw = Sq$. We show that $Aq = Sq$. Now,

$$\begin{aligned} \rho(Aq, Sq, w) &= \rho(Aq, Bw, w) \\ &\leq \lambda \max \{ \rho(Sq, Tw, w), \rho(Sq, Aq, w), \rho(Tw, Bw, w) \} \\ &= \lambda \rho(Sq, Aq, w), \end{aligned} \quad (2.53)$$

which implies that $Aq = Sq$. Since $\{A, S\}$ are limit coincidentally commuting $Sw = SAq = ASq = Aw = w$. Thus, $Aw = Sw = Tw = Bw$, that is, w is a common fixed point of A, B, S , and T . Similarly, if B is continuous, it is proved that A, B, S , and T have a common fixed point.

To prove the uniqueness, let $w^* (\neq w)$ be common fixed point of A, B, S , and T . Then,

$$\begin{aligned} \rho(w, w^*, w^*) &= \rho(Aw, Bw^*, w^*) \\ &\leq \lambda \max \{ \rho(Sw, Tw^*, w^*), \rho(Sw, Aw, w^*), \rho(Tw^*, Bw^*, w^*) \} \\ &= \lambda \rho(w, w, w^*). \end{aligned} \quad (2.54)$$

Similarly,

$$\rho(w, w, w^*) \leq \lambda \rho(w, w^*, w^*). \quad (2.55)$$

Substituting (2.54) into (2.55),

$$\rho(w, w, w^*) \leq \lambda^2 \rho(w, w, w^*), \quad (2.56)$$

which is a contradiction. Hence, $w = w^*$. This completes the proof. $\square$

Letting $S = T$ in [Theorem 2.4](#), we obtain the following corollary.

COROLLARY 2.5. *Let A , B , and S be three selfmappings of a D -metric space satisfying (2.11) and*

$$\rho(Ax, By, z) \leq \lambda \max \{ \rho(Sx, Sy, z), \rho(Sx, Ax, z), \rho(Sy, By, z) \} \quad (2.57)$$

for all $x, y, z \in X$, where $0 \leq \lambda < 1$.

Further assume that

- (a) $\overline{O_{A,B}(Sx)}$ is complete for each $x \in X$,
- (b) $\{A, S\}$ and $\{B, S\}$ are limit coincidentally commuting,
- (c) any one of A , B , and S is continuous.

Then A , B , and S have a unique common fixed point.

ACKNOWLEDGMENTS. We would like to express our thanks to the referee for his helpful suggestions. This work was supported by grant no. 2001-1-10100-005-2 from the Basic Research Program of the Korea Science and Engineering Foundation.

REFERENCES

- [1] B. C. Dhage, *On two general existence principles for proving the common fixed point theorems in D -metric spaces*, to appear in Indian J. Pure Appl. Math.
- [2] ———, *A study of some fixed point theorems*, Ph.D. thesis, Marathwada University, Aurangabad, India, 1984.
- [3] ———, *Generalised metric spaces and mappings with fixed point*, Bull. Calcutta Math. Soc. **84** (1992), no. 4, 329–336.
- [4] ———, *On common fixed points of pairs of coincidentally commuting mappings in D -metric spaces*, Indian J. Pure Appl. Math. **30** (1999), no. 4, 395–406.
- [5] ———, *Generalized metric spaces and topological structure. I*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi Sect. I a Mat. **46** (2000), no. 1, 3–24.
- [6] B. C. Dhage and B. E. Rhoades, *Proving fixed point theorems in D -metric spaces via general existence principle*, to appear in Indian J. Pure Appl. Math.

B. C. Dhage: Gurukul Colony, Ahmedpur-413 515, Latur District (Maharashtra), India
E-mail address: bcd20012001@yahoo.co.in

Smrati Arya: Chhatrapati Shivaji Ward, Kothari Gali, Harda-461331, India

Jeong Sheok Ume: Department of Applied Mathematics, Changwon National University, Changwon 641-773, Korea
E-mail address: jsume@changwon.ac.kr