DERIVATIONS ON BANACH ALGEBRAS

S. HEJAZIAN and S. TALEBI

Received 12 September 2002

Let D be a derivation on a Banach algebra; by using the operator D^2 , we give necessary and sufficient conditions for the separating ideal of D to be nilpotent. We also introduce an ideal $M(D)$ and apply it to find out more equivalent conditions for the continuity of D and for nilpotency of its separating ideal.

2000 Mathematics Subject Classification: 46H40, 47B47.

1. Introduction. Let A be a Banach algebra. By a derivation on A , we mean a linear mapping $D : A \rightarrow A$, which satisfies $D(ab) = aD(b) + D(a)b$ for all a and b in A . The separating space of D is the set

$$S(D) = \{a \in A : \exists \{a_n\} \subset A; a_n \rightarrow 0, D(a_n) \rightarrow a\}. \quad (1.1)$$

The set $S(D)$ is a closed ideal of A which, by the closed-graph theorem, is zero if and only if D is continuous.

DEFINITION 1.1. A closed ideal J of A is said to be a separating ideal if, for each sequence $\{a_n\}$ in A , there is a natural N such that

$$\overline{(Ja_n \cdots a_1)} = \overline{(Ja_N \cdots a_1)} \quad (n \geq N). \quad (1.2)$$

The separating space of a derivation on A is a separating ideal [2, Chapter 5]; it also satisfies the same property for the left products.

The following assertions are of the most famous conjectures about derivations on Banach algebras:

- (C1) every derivation on a Banach algebra has a nilpotent separating ideal;
- (C2) every derivation on a semiprime Banach algebra is continuous;
- (C3) every derivation on a prime Banach algebra is continuous;
- (C4) every derivation on a Banach algebra leaves each primitive ideal invariant.

Clearly, if (C1) is true, then the same for (C2) and (C3). Mathieu and Runde in [5] proved that (C1), (C2), and (C3) are equivalent. The conjecture (C4) is known as the noncommutative Singer-Wermer conjecture, and it has been proved in [1] that if each of the conjectures (C1), (C2), or (C3) hold, then (C4) is also true. The conjectures (C1), (C2), and (C3) are still open even if A is assumed

to be commutative, but (C4) is true in the commutative case, see [7]. These conjectures are also related to some other famous open problems; the reader is referred to [1, 3, 4, 5, 9] for more details.

In the next section, we deal with (C1), and although, for a derivation D on a Banach algebra, the operators D^n , $n = 2, 3, \dots$, are more complicated, by considering D^2 , we easily give some equivalent conditions for $S(D)$ to be nilpotent. As a consequence, we reprove some of the results in [8]. At the end of the next section, we introduce an ideal related to a derivation and apply it to obtain some equivalent conditions for continuity of D and for nilpotency of $S(D)$.

We recall that $S(D)$ is nilpotent if and only if $S(D) \cap R$ is nilpotent, see [1, Lemma 4.2].

2. The results. From now on, A is a Banach algebra, and R and L denote the Jacobson radical and the nil radical of A , respectively, (see [6, Chapter 4] for definitions). Note that D is a derivation on A , and $S(D)$ is the separating ideal of D . If B_i 's, $i = 1, 2, \dots, n$, are subsets of A , then $B_1 B_2 \cdots B_n$ denotes the linear span of the set $\{b_1 b_2 \cdots b_n : b_i \in B_i, \text{ for } i = 1, 2, \dots, n\}$, and if all of B_i 's coincide with each other, we denote this set by B^n .

THEOREM 2.1. *Let J be a closed left ideal of A . Then, $S(D) \cap J$ is nilpotent if and only if $D^2 \mid_{\bigcap_{n=1}^{\infty} (S(D) \cap J)^n}$ is continuous.*

PROOF. Suppose that D^2 is continuous on $\overline{\bigcap_{n=1}^{\infty} (S(D) \cap J)^n}$. Consider a in $S(D) \cap J$, then for each $n \in \mathbb{N}$, $a^n \in (S(D) \cap J)^n$, and since $S(D)$ is a separating ideal, there exists $N \in \mathbb{N}$ such that

$$\overline{S(D)a^n} = \overline{S(D)a^N} \quad (n \geq N). \quad (2.1)$$

Hence, by the Mittag-Leffler theorem [2, Theorem A.1.25] and the fact that $S(D)a^n \subseteq (S(D) \cap J)^n$, we have

$$\overline{S(D)a^N} = \bigcap_{n=1}^{\infty} \overline{S(D)a^n} = \bigcap_{n=1}^{\infty} \overline{S(D)a^n} \subseteq \bigcap_{n=1}^{\infty} \overline{(S(D) \cap J)^n}. \quad (2.2)$$

Now, let $\{x_n\} \subseteq A$, $x_n \rightarrow 0$, and $D(x_n) \rightarrow a^{N+1}$. Take $y_n = x_n a^{N+1}$, then $y_n \in S(D)a^N \subseteq \bigcap_{n=1}^{\infty} (S(D) \cap J)^n$, $y_n \rightarrow 0$, and $D(y_n) \rightarrow a^{2(N+1)}$, and by the hypothesis, $D^2(y_n) \rightarrow 0$ and $D^2(y_n^2) \rightarrow 0$. On the other hand,

$$D^2(y_n^2) = y_n D^2(y_n) + 2(Dy_n)^2 + D^2(y_n)y_n \rightarrow 2a^{4(N+1)}. \quad (2.3)$$

Therefore, $a^{4N+4} = 0$, that is, $S(D) \cap J$ is a nil and hence a nilpotent ideal by closedness [6, Theorem 4.4.11]. The converse is trivial. $\square$

REMARK 2.2. (i) Note that in [Theorem 2.1](#), we can replace J by a right ideal, see [\[2, Theorem 5.2.24\]](#).

(ii) The argument of [Theorem 2.1](#) shows that if J is not assumed to be closed and if D^2 is continuous on $\bigcap_{n=1}^{\infty} (S(D) \cap J)^n$, then $S(D) \cap J$ will be a nil ideal.

COROLLARY 2.3. *The set $S(D)$ is nilpotent if and only if $D^2|_{\overline{\bigcap_{n=1}^{\infty} (S(D) \cap R)^n}}$ is continuous.*

PROOF. If $S(D)$ is nilpotent, then the result is obvious. Conversely, by [Theorem 2.1](#), $S(D) \cap R$ is nilpotent, and by [\[1, Lemma 4.2\]](#), $S(D)$ is nilpotent. $\square$

COROLLARY 2.4. *If $\dim(\bigcap_{n=1}^{\infty} (S(D) \cap R)^n) < \infty$, then $S(D)$ is nilpotent.*

The assertions of the following theorem were proved by Villena in [\[8\]](#), see also [\[9, Theorem 4.4\]](#). Using [Theorem 2.1](#), we can reprove them in a different way.

THEOREM 2.5. *The derivation D is continuous if one of the following assertions hold:*

- (a) A is semiprime and $\dim(R \cap (\bigcap_{n=1}^{\infty} A^n)) < \infty$;
- (b) A is prime and $\dim(\bigcap_{n=1}^{\infty} (aA \cap R)^n) < \infty$ for some $a \in A$ with $a^2 \neq 0$;
- (c) A is an integral domain and $\dim(\bigcap_{n=1}^{\infty} (aA \cap R)^n) < \infty$ for some nonzero $a \in A$.

PROOF. (a) By [Corollary 2.4](#), $S(D)$ is nilpotent, and since A is semiprime, D is continuous.

(b) Without loss of generality, we may assume that A has an identity. By assumption, $\bigcap_{n=1}^{\infty} (aA \cap R \cap S(D))^n$ is finite dimensional; thus, D^2 is continuous on this space, and by [Remark 2.2\(ii\)](#), $aA \cap R \cap S(D)$ is a nil right ideal; therefore, $a(S(D) \cap R)$ is a nil right ideal, and by [\[6, Theorem 4.4.11\]](#), $a(S(D) \cap R) \subseteq L = \{0\}$. Thus, $AaA(S(D) \cap R) = \{0\}$, where AaA is the ideal generated by a . Since $a^2 \neq 0$ and A is prime, it follows that $S(D) \cap R = \{0\}$ and hence $S(D) \subseteq L = \{0\}$.

(c) The same argument as in (b) shows that $a(S(D) \cap R) = \{0\}$, and since A is an integral domain, $S(D) \cap R = \{0\}$ and D is continuous. $\square$

In the sequel, we give other equivalent conditions for $S(D)$ to be nilpotent, but first we introduce the set

$$M(D) = \{x \in S(D) \cap R : D(x) \in R\}. \quad (2.4)$$

Obviously, $M(D)$ is an ideal of A and $(S(D) \cap R)^2 \subseteq M(D)$. The following theorems show that this ideal can help us to study the continuity of a derivation or nilpotency of its separating ideal.

THEOREM 2.6. *The derivation D is continuous if and only if $M(D) = \{0\}$.*

PROOF. Clearly, if D is continuous, then $M(D) = \{0\}$. Conversely, let $M(D) = \{0\}$; then, $(S(D) \cap R)^2 = \{0\}$. Therefore, $(S(D) \cap R)$ and hence $S(D)$ is a nilpotent ideal. Therefore, $S(D) \subseteq L$; we also have $D(L) \subseteq L$ by [1, Lemma 4.1]; thus, $D(S(D)) \subseteq R$, that is, $S(D) \subseteq M(D) = \{0\}$ and D is continuous. $\square$

THEOREM 2.7. *The following assertions are equivalent:*

- (a) $S(D)$ is nilpotent;
- (b) $M(D)$ is a nil ideal;
- (c) $\bigcap_{n=1}^{\infty} M(D)^n = \{0\}$.

PROOF. Clearly, (a) implies (b). Suppose that (b) holds, then $(S(D) \cap R)^2$ is a nil ideal; therefore, $S(D)$ is a nilpotent ideal and (a) holds. Now, if $S(D)$ is nilpotent, then $\bigcap_{n=1}^{\infty} (S(D)^n) = \{0\}$ and this implies (c). Finally, if $\bigcap_{n=1}^{\infty} M(D)^n = \{0\}$, then by Theorem 2.1 and Remark 2.2 $M(D) = M(D) \cap S(D)$ is a nil ideal and (c) implies (b). $\square$

ACKNOWLEDGMENT. The authors would like to thank The Payame Noor University of Iran for the financial support.

REFERENCES

- [1] J. Cusack, *Automatic continuity and topologically simple radical Banach algebras*, J. London Math. Soc. (2) **16** (1977), no. 3, 493–500.
- [2] H. G. Dales, *Banach Algebras and Automatic Continuity*, London Mathematical Society Monographs. New Series, vol. 24, The Clarendon Press, New York, 2000.
- [3] M. Mathieu, *Where to find the image of a derivation*, Functional Analysis and Operator Theory (Warsaw, 1992), Banach Center Publ., vol. 30, Polish Academy of Sciences, Warsaw, 1994, pp. 237–249.
- [4] M. Mathieu and G. J. Murphy, *Derivations mapping into the radical*, Arch. Math. (Basel) **57** (1991), no. 5, 469–474.
- [5] M. Mathieu and V. Runde, *Derivations mapping into the radical. II*, Bull. London Math. Soc. **24** (1992), no. 5, 485–487.
- [6] T. W. Palmer, *Banach Algebras and the General Theory of *-Algebras. Vol. I*, Encyclopedia of Mathematics and Its Applications, vol. 49, Cambridge University Press, Cambridge, 1994.
- [7] M. P. Thomas, *The image of a derivation is contained in the radical*, Ann. of Math. (2) **128** (1988), no. 3, 435–460.
- [8] A. R. Villena, *Derivations with a hereditary domain. II*, Studia Math. **130** (1998), no. 3, 275–291.
- [9] ———, *Automatic continuity in associative and nonassociative context*, Irish Math. Soc. Bull. (2001), no. 46, 43–76.

S. Hejzian: Department of Mathematics, Ferdowsi University, Mashhad, Iran
E-mail address: hejazian@math.um.ac.ir

S. Talebi: Department of Mathematics, Payame Noor University, Mashhad, Iran
E-mail address: talebi@mshc.pnu.ac.ir