WAVELET ANALYSIS ON A BOEHMIAN SPACE

R. ROOPKUMAR

Received 24 May 2002

We extend the wavelet transform to the space of periodic Boehmians and discuss some of its properties.

2000 Mathematics Subject Classification: 44A15, 46F12, 42C40, 44A99.

1. Introduction. The concept of Boehmians was introduced by J. Mikusiński and P. Mikusiński [7], and the space of Boehmians with two notions of convergences was well established in [8]. Many integral transforms have been extended to the context of Bohmian spaces, for example, Fourier transform [9, 10, 11], Laplace transform [13, 17], Radon transform [14], and Hilbert transform [3, 5].

On the other hand, the theory of wavelet transform is recently developed, and it has various applications in signal processing, especially to analyze non-stationary signals by providing the time-frequency representation of the signal. For a fixed $g \in \mathcal{L}^2(\mathbb{R})$, called a mother wavelet, the wavelet transform $\Phi_g : \mathcal{L}^2(\mathbb{R}) \rightarrow \mathcal{L}^2(\mathbb{R} \times \mathbb{R}^+)$ is defined by

$$\Phi_g(f)(a, b) = \int_{-\infty}^{\infty} f(x) \overline{g_{a,b}(x)} dx \quad \text{for } a > 0, b \in \mathbb{R}, \quad (1.1)$$

where $g_{a,b}(x) = (1/\sqrt{a})g((x-b)/a)$, $x \in \mathbb{R}$, are called wavelets. For more details, we refer the reader to [6]. In [4], we extended the wavelet transform to a Bohmian space which properly contains $\mathcal{L}^2(\mathbb{R})$ and studied its properties.

Holschneider [2] introduced the wavelet transform on the space $C^\infty(\mathbb{T})$ of smooth functions on the unit circle $\mathbb{T}$ of the complex plane and gave an extension to the space of periodic distributions. In Section 2, we fix some notations and discuss the theory of wavelet transform on $C^\infty(\mathbb{T})$. In Section 3, we briefly recall the periodic Boehmians, construct a new Bohmian space $\mathcal{B}(\mathcal{S}(\mathbb{Y}), (C^\infty(\mathbb{T}), *, \odot, \Delta))$, and verify some auxiliary results. In Section 4, we define wavelet transform on the space of periodic Boehmians and prove that it is consistent with the wavelet transform on $C^\infty(\mathbb{T})$. Further, we establish that the extended wavelet transform is linear and continuous with respect to δ -convergence as well as Δ -convergence.

2. Preliminaries. The space $C^\infty(\mathbb{T})$ consists of infinitely differentiable, periodic functions on $\mathbb{R}$ of period 2π , with the Fréchet space topology induced by the increasing sequence of seminorms

$$\|\phi\|_{C^\infty(\mathbb{T});n} = \sum_{p=0}^n \sup_{t \in [0,2\pi]} |\partial^p \phi(t)|. \quad (2.1)$$

We know that

$$C^\infty(\mathbb{T}) = C_+^\infty(\mathbb{T}) \oplus C_-^\infty(\mathbb{T}) \oplus K(\mathbb{T}), \quad (2.2)$$

where $C_+^\infty(\mathbb{T})$ and $C_-^\infty(\mathbb{T})$ are the subspaces consisting of functions with positive and negative Fourier coefficients, respectively, and $K(\mathbb{T})$ is the space of constant functions.

Let $\mathcal{S}(\mathbb{R})$ denote the space of rapidly decreasing functions on $\mathbb{R}$. (See [1].) Given $f \in \mathcal{S}(\mathbb{R})$, $b \in [0, 2\pi]$, and $a > 0$, define $f_a, f_{b,a} \in C^\infty(\mathbb{T})$ by

$$\begin{aligned} f_a(x) &= \sum_{n \in \mathbb{Z}} \frac{1}{a} f\left(\frac{x + 2n\pi}{a}\right), \quad x \in [0, 2\pi], \\ f_{b,a}(x) &= f_a(x - b), \quad x \in [0, 2\pi]. \end{aligned} \quad (2.3)$$

Let $\mathcal{S}(\mathbb{Y})$ denote the Fréchet space of all smooth functions $\eta(b, a)$ of rapid descent on $\mathbb{R} \times \mathbb{R}^+$ which are periodic functions in the variable b of period 2π , with the following directed family of seminorms:

$$\|\eta\|_{\mathcal{S}(\mathbb{Y});n,\alpha,\beta} = \sum_{\substack{0 \leq p \leq n \\ 0 \leq l \leq \alpha \\ 0 \leq k \leq \beta}} \sup_{a > 0} \sup_{b \in [0, 2\pi]} |a^p \partial_a^l \partial_b^k \eta(b, a)|. \quad (2.4)$$

We choose a mother wavelet $g \in \mathcal{S}(\mathbb{R})$ with all moments $\int_{-\infty}^{\infty} x^n g(x) dx$ are equal to zero.

DEFINITION 2.1. The wavelet transform $T_g : C^\infty(\mathbb{T}) \rightarrow \mathcal{S}(\mathbb{Y})$ is defined by

$$T_g(\phi) = \int_0^{2\pi} \phi(x) \overline{g_{b,a}(x)} dx, \quad b \in \mathbb{R}, a > 0. \quad (2.5)$$

THEOREM 2.2. The wavelet transform $T_g : C^\infty(\mathbb{T}) \rightarrow \mathcal{S}(\mathbb{Y})$ is continuous and linear.

DEFINITION 2.3. The map $R_g : \mathcal{S}(\mathbb{Y}) \rightarrow C^\infty(\mathbb{T})$ is defined by

$$(R_g \eta)(x) = \int_0^{2\pi} \int_0^\infty g_{b,a}(x) \eta(b, a) \frac{da db}{a}. \quad (2.6)$$

THEOREM 2.4. *The map $R_g : \mathcal{S}(\mathbb{Y}) \rightarrow C^\infty(\mathbb{T})$ is continuous and linear.*

A partial inversion formula is given by the following theorem.

THEOREM 2.5. *If $\hat{g}$ is the Fourier transform of g and $C_g^+ = \int_0^\infty |\hat{g}(a)|^2 (da/a)$, $C_g^- = \int_0^\infty |\hat{g}(-a)|^2 (da/a)$, then*

$$\begin{aligned} R_g \circ T_g \phi &= C_g^+ \phi, \quad \forall \phi \in C_+^\infty(\mathbb{T}), \\ R_g \circ T_g \phi &= C_g^- \phi, \quad \forall \phi \in C_-^\infty(\mathbb{T}). \end{aligned} \quad (2.7)$$

3. Boehmian spaces. The triplet $(C^\infty(\mathbb{T}), *, \Delta)$, where $*$: $C^\infty(\mathbb{T}) \times C^\infty(\mathbb{T}) \rightarrow C^\infty(\mathbb{T})$ is defined by

$$(\phi * \psi)(x) = \int_0^{2\pi} \phi(x-t)\psi(t)dt, \quad x \in [0, 2\pi] \quad (3.1)$$

and Δ is the collection of all sequences (δ_k) from $C^\infty(\mathbb{T})$ satisfying

- (1) $\int_0^{2\pi} \delta_k(t)dt = 1$ for all $k \in \mathbb{N}$,
- (2) $\int_0^{2\pi} |\delta_k(t)|dt \leq M$ for all $k \in \mathbb{N}$, for some $M > 0$,
- (3) $s(\delta_k) \rightarrow 0$ as $n \rightarrow \infty$ where $s(\delta_k) = \sup\{t \in [0, 2\pi] : \delta_k(t) \neq 0\}$,

is the collection of all equivalence classes $[\phi_k/\delta_k]$ given by the equivalence relation $\sim$ defined by

$$((\phi_k), (\delta_k)) \sim ((\psi_k), (\epsilon_k)) \quad \text{if } \phi_k * \epsilon_j = \psi_j * \delta_k \quad \forall k, j \in \mathbb{N} \quad (3.2)$$

on the collection $\mathcal{A}$ of pair of sequences $((\phi_k), (\delta_k))$, $\phi_n \in C^\infty(\mathbb{T})$, $(\delta_k) \in \Delta$ satisfying

$$\phi_k * \delta_j = \phi_j * \delta_k, \quad \forall k, j \in \mathbb{N}. \quad (3.3)$$

This triplet with addition and scalar multiplication, defined by

$$\begin{aligned} \left[\frac{\phi_k}{\delta_k} \right] + \left[\frac{\psi_k}{\epsilon_k} \right] &= \left[\frac{\phi_k * \epsilon_k + \psi_k * \delta_k}{\delta_k * \epsilon_k} \right], \\ \alpha \left[\frac{\phi_k}{\delta_k} \right] &= \left[\frac{\alpha \phi_k}{\delta_k} \right], \end{aligned} \quad (3.4)$$

is called the periodic Boehmian space [15, 16], and we denote it by $\mathcal{B}_\mathbb{T}$.

DEFINITION 3.1 (δ -convergence). A sequence (x_n) δ -converges to x in $\mathcal{B}_\mathbb{T}$, denoted by $x_n \xrightarrow{\delta} x$ as $n \rightarrow \infty$ in $\mathcal{B}_\mathbb{T}$ if there exists $(\delta_k) \in \Delta$ such that

$x_n * \delta_k, x * \delta_k \in C^\infty(\mathbb{T})$, and for each $k \in \mathbb{N}$,

$$x_n * \delta_k \rightarrow x * \delta_k \quad \text{as } n \rightarrow \infty \text{ in } C^\infty(\mathbb{T}). \quad (3.5)$$

The following theorem is proved in [8].

THEOREM 3.2. *Let $x_n, x \in \mathcal{B}_{\mathbb{T}}$, $n \in \mathbb{N}$. $x_n \xrightarrow{\delta} x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{T}}$ if and only if there exist $\phi_{n,k}, \phi_k \in C^\infty(\mathbb{T})$ such that $x_n = [\phi_{n,k}/\delta_k], [\phi_k/\delta_k]$ and, for each $k \in \mathbb{N}$,*

$$\phi_{n,k} \rightarrow \phi_k \quad \text{as } n \rightarrow \infty \text{ in } C^\infty(\mathbb{T}). \quad (3.6)$$

DEFINITION 3.3 (Δ -convergence). A sequence (x_n) Δ -converges to x in $\mathcal{B}_{\mathbb{T}}$, denoted by $x_n \xrightarrow{\Delta} x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{T}}$ if there exists a delta-sequence (δ_n) such that $(x_n - x) * \delta_n \in C^\infty(\mathbb{T})$ for each $n \in \mathbb{N}$ and

$$(x_n - x) * \delta_n \rightarrow 0 \quad \text{as } n \rightarrow \infty \text{ in } C^\infty(\mathbb{T}). \quad (3.7)$$

Now, we construct a new Boehmian space as follows.

As in the context of Boehmian space defined in [12], we take the vector space Γ and the commutative semi-group as $\mathcal{S}(\mathbb{Y})$ and $(C^\infty(\mathbb{T}), *)$, respectively.

DEFINITION 3.4. Given $\eta \in \mathcal{S}(\mathbb{Y})$ and $\phi \in C^\infty(\mathbb{T})$, define

$$(\eta \odot \phi)(b, a) = \int_0^{2\pi} \eta(b - t, a) \phi(t) dt. \quad (3.8)$$

LEMMA 3.5. *If $\eta \in \mathcal{S}(\mathbb{Y})$ and $\phi \in C^\infty(\mathbb{T})$, then $\eta \odot \phi \in \mathcal{S}(\mathbb{Y})$.*

PROOF. To prove that $(\eta \odot \phi)(b, a)$ is infinitely differentiable, we show that

$$\begin{aligned} \partial_a(\eta \odot \phi)(b, a) &= (\partial_a \eta \odot \phi)(b, a), \\ \partial_b(\eta \odot \phi)(b, a) &= (\partial_b \eta \odot \phi)(b, a). \end{aligned} \quad (3.9)$$

Fix $a_0 > 0, b_0 \in \mathbb{R}$ arbitrarily.

Consider $((\eta \odot \phi)(b_0, a) - (\eta \odot \phi)(b_0, a_0)) / (a - a_0) = \int_0^{2\pi} (\eta(b_0 - t, a) - \eta(b_0 - t, a_0)) / (a - a_0) \phi(t) dt$. Using the mean-value theorem (in the variable a), we get that the integrand is dominated by $\|\eta\|_{\mathcal{S}(\mathbb{Y}); 0, 1, 0} \|\phi\|_{C^\infty(\mathbb{T}), 0}$. Therefore, we can apply Lebesgue dominated convergence theorem [18], and we get

$$\begin{aligned} \partial_a(\eta \odot \phi)(b_0, a_0) &= \lim_{a \rightarrow a_0} \int_0^{2\pi} \frac{\eta(b_0 - t, a) - \eta(b_0 - t, a_0)}{a - a_0} \phi(t) dt \\ &= \int_0^{2\pi} \lim_{a \rightarrow a_0} \frac{\eta(b_0 - t, a) - \eta(b_0 - t, a_0)}{a - a_0} \phi(t) dt \\ &= \int_0^{2\pi} \partial_a \eta(b_0 - t, a_0) \phi(t) dt \\ &= (\partial_a \eta \odot \phi)(b_0, a_0). \end{aligned} \quad (3.10)$$

By a similar argument, we can prove that $\partial_b(\eta \odot \phi)(b, a) = (\partial_b \eta \odot \phi)(b, a)$. Finally by a routine manipulation, we get

$$\|\eta \odot \phi\|_{\mathcal{G}(\mathbb{Y});n,\alpha,\beta} \leq \|\phi\|_{\mathcal{L}^1(\mathbb{T})} \|\eta\|_{\mathcal{G}(\mathbb{Y});n,\alpha,\beta}, \quad (3.11)$$

where $\|\phi\|_{\mathcal{L}^1(\mathbb{T})} = \int_0^{2\pi} |\phi(t)| dt$. Hence, $\eta \odot \phi \in \mathcal{G}(\mathbb{Y})$. $\square$

LEMMA 3.6. *If $\eta \in \mathcal{G}(\mathbb{Y})$ and $(\delta_n) \in \Delta$, then $\eta \odot \delta_n \rightarrow \phi$ as $n \rightarrow \infty$ in $\mathcal{G}(\mathbb{Y})$.*

PROOF. Let $p, k, l \in \mathbb{N}_0$ be arbitrary. Using the mean-value theorem and a property of δ -sequence, we get

$$\begin{aligned} |a^p \partial_a^l \partial_b^k (\eta \odot \delta_n - \eta)(b, a)| &= |a^p ((\partial_a^l \partial_b^k \eta) \odot \delta_n)(b, a) - a^p \partial_a^l \partial_b^k \eta(b, a)| \\ &\leq \int_0^{2\pi} |a^p (\partial_a^l \partial_b^k \eta(b - t, a) - \partial_a^l \partial_b^k \eta(b, a)) \delta_n(t)| dt \\ &\leq \|\eta\|_{\mathcal{G}(\mathbb{Y});p,l,k+1} \int_0^{2\pi} |t| |\delta_n(t)| dt \\ &\leq Ms(\delta_n) \|\eta\|_{\mathcal{G}(\mathbb{Y});p,l,k+1}, \end{aligned} \quad (3.12)$$

which tends to 0 as $n \rightarrow \infty$. This completes the proof of the lemma. $\square$

LEMMA 3.7. *If $\eta_n \rightarrow \eta$ as $n \rightarrow \infty$ in $\mathcal{G}(\mathbb{Y})$ and $\psi \in C^\infty(\mathbb{T})$, then $\eta_n \odot \psi \rightarrow \eta \odot \psi$ as $n \rightarrow \infty$.*

PROOF. Let $p, k, l \in \mathbb{N}_0$ be arbitrary. Now,

$$\begin{aligned} |a^p \partial_a^l \partial_b^k (\eta_n \odot \psi - \eta \odot \psi)(b, a)| &= \int_0^{2\pi} a^p |\partial_a^l \partial_b^k (\eta_n - \eta)(b, a)| |\psi(t)| dt \\ &\leq \|\psi\|_{\mathcal{L}^1(\mathbb{T})} \|\eta_n - \eta\|_{\mathcal{G}(\mathbb{Y});p,l,k} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (3.13)$$

Hence, the lemma follows. $\square$

LEMMA 3.8. *If $\eta_n \rightarrow \eta$ as $n \rightarrow \infty$ in $\mathcal{G}(\mathbb{Y})$ and $\delta_n \in \Delta$, then $\eta_n \odot \delta_n \rightarrow \eta$ as $n \rightarrow \infty$.*

PROOF. Since we have $\eta_n \odot \delta_n - \eta = \eta_n \odot \delta_n - \eta \odot \delta_n + \eta \odot \delta_n - \eta$ and [Lemma 3.6](#), we merely prove that $\eta_n \odot \delta_n - \eta \odot \delta_n \rightarrow 0$ as $n \rightarrow \infty$.

If $p, k, l \in \mathbb{N}_0$, then, using a property of delta-sequence, we get

$$\begin{aligned} |a^p \partial_a^l \partial_b^k (\eta_n - \eta) \odot \delta_n(b, a)| &= \int_0^{2\pi} |\delta_n(t)| |a^p \partial_a^l \partial_b^k (\eta_n - \eta)(b, a)| dt \\ &\leq \|\eta_n - \eta\|_{\mathcal{G}(\mathbb{Y});p,l,k} \int_0^{2\pi} |\delta_n(t)| dt \leq M \|\eta_n - \eta\|_{\mathcal{G}(\mathbb{Y});p,l,k}. \end{aligned} \quad (3.14)$$

The above inequalities prove the lemma. $\square$

Now using the above lemmas we can construct the Boehmian space $\mathcal{B}_{\mathbb{Y}} = (\mathcal{S}_{\mathbb{Y}}, (C^{\infty}, *), \odot, \Delta)$ in a canonical way.

4. Generalized wavelet transform

DEFINITION 4.1. Define $\mathcal{T}_g : \mathcal{B}_{\mathbb{T}} \rightarrow \mathcal{B}_{\mathbb{Y}}$ by

$$\mathcal{T}_g \left(\left[\frac{\phi_n}{\delta_n} \right] \right) = \left[\frac{T_g \phi_n}{\delta_n} \right]. \quad (4.1)$$

THEOREM 4.2. *The generalized wavelet transform $\mathcal{T}_g : \mathcal{B}_{\mathbb{T}} \rightarrow \mathcal{B}_{\mathbb{Y}}$ is well defined.*

First, we state and prove a lemma that will be useful.

LEMMA 4.3. *If $\phi, \psi \in C^{\infty}(\mathbb{T})$, then $T_g(\phi * \psi) = T_g \phi \odot \psi$.*

PROOF. Let $a \in \mathbb{R}^+$ and $b \in \mathbb{R}$ be arbitrary. Now

$$\begin{aligned} T_g(\phi * \psi)(b, a) &= \int_0^{2\pi} (\phi * \psi)(x) \overline{g_a(x-b)} dx \\ &= \int_0^{2\pi} \overline{g_a(x-b)} dx \int_0^{2\pi} \phi(x-t) \psi(t) dt. \end{aligned} \quad (4.2)$$

By an easy verification, we can apply Fubini's theorem and the last integral equals

$$\begin{aligned} &\int_0^{2\pi} \psi(t) dt \int_0^{2\pi} \phi(x-t) \overline{g_a(x-b)} dx \\ &= \int_0^{2\pi} \psi(t) dt \int_0^{2\pi} \phi(x) \overline{g_a(x-(b-t))} dx \\ &= (T_g \phi \odot \psi)(b, a). \end{aligned} \quad (4.3)$$

□

PROOF OF THEOREM 4.2. First, we show that $((T_g \phi_n), (\delta_n))$ is a quotient. Since $[\phi_n / \delta_n] \in \mathcal{B}_{\mathbb{T}}$, we have

$$\phi_k * \delta_j = \phi_j * \delta_k, \quad \forall j, k \in \mathbb{N}. \quad (4.4)$$

Applying the classical wavelet transform T_g on both sides, we get

$$T_g \phi_k \odot \delta_j = T_g \phi_j \odot \phi_k, \quad \forall j, k \in \mathbb{N} \text{ (by Lemma 4.3)}. \quad (4.5)$$

Next, we show that the definition of $\mathcal{T}_g$ is independent of the choice of the representative.

Let $[\phi_k/\epsilon_k] = [\psi_k/\delta_k]$ in $\mathcal{B}_{\mathbb{T}}$. Then, we have

$$\phi_k * \epsilon_j = \psi_j * \delta_k, \quad \forall j, k \in \mathbb{N}. \quad (4.6)$$

Again, applying the wavelet transform and using [Lemma 4.3](#), we get

$$\mathcal{T}_g \phi_k \odot \epsilon_j = \mathcal{T}_g \psi_j \odot \delta_k, \quad \forall j, k \in \mathbb{N}. \quad (4.7)$$

Hence, the theorem follows. $\square$

THEOREM 4.4 (consistency). *Let $\mathcal{I}_{\mathbb{T}} : C^\infty(\mathbb{T}) \rightarrow \mathcal{B}_{\mathbb{T}}$ and $\mathcal{I}_{\mathbb{V}} : \mathcal{F}(\mathbb{V}) \rightarrow \mathcal{B}_{\mathbb{V}}$ be the canonical identification defined, respectively, by*

$$\phi \mapsto \left[\frac{\phi * \delta_n}{\delta_n} \right], \quad \eta \mapsto \left[\frac{\eta \odot \delta_n}{\delta_n} \right], \quad (4.8)$$

where $(\delta_n) \in \Delta$, then $\mathcal{T}_g \circ \mathcal{I}_{\mathbb{T}} = \mathcal{I}_{\mathbb{V}} \circ T_g$.

PROOF. Let $\phi \in C^\infty(\mathbb{T})$, then

$$\begin{aligned} \mathcal{T}_g(\mathcal{I}_{\mathbb{T}}(\phi)) &= \mathcal{T}_g\left(\left[\frac{\phi * \delta_n}{\delta_n}\right]\right) = \left[\frac{T_g(\phi * \delta_n)}{\delta_n}\right] \\ &= \left[\frac{T_g \phi \odot \delta_n}{\delta_n}\right] \quad (\text{by } \text{Lemma 4.3}) \\ &= \mathcal{I}_{\mathbb{V}}(T_g(\phi)). \end{aligned} \quad (4.9)$$

$\square$

THEOREM 4.5. *The wavelet transform $\mathcal{T}_g : \mathcal{B}_{\mathbb{T}} \rightarrow \mathcal{B}_{\mathbb{V}}$ is a linear map.*

PROOF. If $[\phi_n/\delta_n], [\psi_n/\epsilon_n] \in \mathcal{B}_{\mathbb{T}}$, then

$$\begin{aligned} \mathcal{T}_g\left(\left[\frac{\phi_n}{\delta_n}\right] + \left[\frac{\psi_n}{\epsilon_n}\right]\right) &= \mathcal{T}_g\left(\left[\frac{\phi_n * \epsilon_n + \psi_n * \delta_n}{\delta_n * \epsilon_n}\right]\right) = \left[\frac{T_g(\phi_n * \epsilon_n + \psi_n * \delta_n)}{\delta_n * \epsilon_n}\right] \\ &= \left[\frac{T_g \phi_n \odot \epsilon_n + T_g \psi_n \odot \delta_n}{\delta_n * \epsilon_n}\right] = \left[\frac{T_g \phi_n}{\delta_n}\right] + \left[\frac{T_g \psi_n}{\epsilon_n}\right] \\ &= \mathcal{T}_g\left(\left[\frac{\phi_n}{\delta_n}\right]\right) + \mathcal{T}_g\left(\left[\frac{\psi_n}{\epsilon_n}\right]\right). \end{aligned} \quad (4.10)$$

If $\alpha \in \mathbb{C}$ and $[\phi_n/\delta_n] \in \mathcal{B}_{\mathbb{T}}$, then

$$\begin{aligned} \mathcal{T}_g\left(\alpha \left[\frac{\phi_n}{\delta_n}\right]\right) &= \mathcal{T}_g\left(\left[\frac{\alpha \phi_n}{\delta_n}\right]\right) = \left[\frac{T_g(\alpha \phi_n)}{\delta_n}\right] = \left[\frac{\alpha T_g \phi_n}{\delta_n}\right] \\ &= \alpha \left[\frac{T_g \phi_n}{\delta_n}\right] = \alpha \mathcal{T}_g\left(\left[\frac{\phi_n}{\delta_n}\right]\right). \end{aligned} \quad (4.11)$$

In the above proof, we have used the fact that T_g is linear wherever it is required. $\square$

From the following two theorems, we say that the generalized wavelet transform is continuous with respect to δ -convergence as well as Δ -convergence.

THEOREM 4.6. *If $x_n \xrightarrow{\delta} x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{T}}$, then $\mathcal{T}_g x_n \xrightarrow{\delta} \mathcal{T}_g x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{Y}}$.*

PROOF. If $x_n \xrightarrow{\delta} x$ as $n \rightarrow \infty$, then, by [Theorem 3.2](#), there exist $\phi_{n,k}, \phi_k \in C^\infty(\mathbb{T})$ and $(\delta_k) \in \Delta$ such that $x_n = [\phi_{n,k}/\delta_k]$ and $x = [\phi_k/\delta_k]$ and, for each $k \in \mathbb{N}$, $\phi_{n,k} \rightarrow \phi_k$ as $n \rightarrow \infty$ in $C^\infty(\mathbb{T})$.

By the continuity of the classical wavelet transform, we have, for each $k \in \mathbb{N}$,

$$T_g \phi_{n,k} \rightarrow T_g \phi_k \quad \text{as } n \rightarrow \infty \text{ in } \mathcal{S}_{\mathbb{Y}}. \quad (4.12)$$

Since $\mathcal{T}_g(x_n) = [T_g \phi_{n,k}/\delta_k]$ and $\mathcal{T}_g(x) = [T_g \phi_k/\delta_k]$, we get $\mathcal{T}_g(x_n) \xrightarrow{\delta} \mathcal{T}_g(x)$ as $n \rightarrow \infty$. Hence, the theorem follows. $\square$

THEOREM 4.7. *If $x_n \xrightarrow{\Delta} x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{T}}$, then $\mathcal{T}_g x_n \xrightarrow{\Delta} \mathcal{T}_g x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{Y}}$.*

PROOF. Let $x_n \xrightarrow{\Delta} x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{T}}$. Then, by definition, we can find $\phi_n \in C^\infty(\mathbb{T})$ and $(\delta_n) \in \Delta$ such that $(x_n - x) * \delta_n = [\phi_n * \delta_k/\delta_k]$ and

$$\phi_n \rightarrow 0 \quad \text{as } n \rightarrow 0 \text{ in } C^\infty(\mathbb{T}). \quad (4.13)$$

Applying the classical wavelet transform and using [Lemma 4.3](#), we get

$$T_g \phi_n \rightarrow 0 \quad \text{as } n \rightarrow 0 \text{ in } \mathcal{S}(\mathbb{Y}). \quad (4.14)$$

Hence, we get $\mathcal{T}_g x_n \xrightarrow{\Delta} \mathcal{T}_g x$ as $n \rightarrow \infty$ in $\mathcal{B}_{\mathbb{Y}}$. $\square$

LEMMA 4.8. *If $\eta \in \mathcal{S}(\mathbb{Y})$ and $\phi \in C^\infty(\mathbb{T})$, then $R_g(\eta \odot \phi) = R_g \eta * \phi$.*

PROOF. Using Fubini's theorem, we get

$$\begin{aligned} R_g(\eta \odot \phi)(x) &= \int_0^{2\pi} \int_0^\infty g_a(x-b)(\eta \odot \phi)(b,a) \frac{dad b}{a} \\ &= \int_0^{2\pi} \int_0^\infty g_a(x-b) \frac{dad b}{a} \int_0^{2\pi} \eta(b-t,a) \phi(t) dt \\ &= \int_0^{2\pi} \phi(t) dt \int_0^{2\pi} \int_0^\infty g_a(x-b) \eta(b-t,a) \frac{dad b}{a} \\ &= \int_0^{2\pi} \phi(t) dt \int_0^{2\pi} \int_0^\infty g_a((x-t)-c) \eta(c,a) \frac{dad c}{a} \quad (b-t=c) \\ &= \int_0^{2\pi} R_g \eta(x-t) \phi(t) dt \\ &= (R_g \eta * \phi)(x). \end{aligned} \quad (4.15)$$

$\square$

Therefore, we can give the following definition.

DEFINITION 4.9. Define $\mathcal{R}_g : \mathcal{B}_{\mathbb{V}} \rightarrow \mathcal{B}_{\mathbb{T}}$ by

$$\mathcal{R}_g \left(\left[\frac{\eta_n}{\delta_n} \right] \right) = \left[\frac{R_g \eta_n}{\delta_n} \right]. \quad (4.16)$$

THEOREM 4.10. *The map $\mathcal{R}_g : \mathcal{B}_{\mathbb{V}} \rightarrow \mathcal{B}_{\mathbb{T}}$ is linear.*

THEOREM 4.11. *The map $\mathcal{R}_g : \mathcal{B}_{\mathbb{V}} \rightarrow \mathcal{B}_{\mathbb{T}}$ is continuous with respect to δ -convergence as well as Δ -convergence.*

Using [Lemma 4.8](#) and [Theorem 2.4](#), we get a proof similar to that of [Theorems 4.6](#) and [4.7](#).

THEOREM 4.12 (an inversion formula). *If $x = [\phi_n / \delta_n] \in \mathcal{B}_{\mathbb{T}}$ such that $\phi_n \in C_{+(-)}^{\infty}(\mathbb{T})$ for all $n \in \mathbb{N}$, then*

$$\mathcal{R}_g \circ \mathcal{T}_g(x) = C_g^{+(-)} x. \quad (4.17)$$

PROOF. Now,

$$\begin{aligned} \mathcal{R}_g \circ \mathcal{T}_g(x) &= \mathcal{R}_g \left(\left[\frac{T_g \phi_n}{\delta_n} \right] \right) = \left[\frac{(R_g \circ T_g) \phi_n}{\delta_n} \right] \\ &= \left[\frac{C_g^{+(-)} \phi_n}{\delta_n} \right] = C_g^{+(-)} \left[\frac{\phi_n}{\delta_n} \right] = C_g^{+(-)} x. \end{aligned} \quad (4.18)$$

□

ACKNOWLEDGMENT. This work was supported by a Senior Research Fellowship from CSIR, India.

REFERENCES

- [1] I. M. Gel'fand and G. E. Shilov, *Generalized Functions. Vol. 2*, Academic Press, New York, 1968.
- [2] M. Holschneider, *Wavelet analysis on the circle*, J. Math. Phys. **31** (1990), no. 1, 39–44.
- [3] V. Karunakaran and N. V. Kalpakam, *Hilbert transform for Boehmians*, Integral Transform. Spec. Funct. **9** (2000), no. 1, 19–36.
- [4] V. Karunakaran and R. Roopkumar, *Wavelet transform for Boehmians*, Proceedings of the International Conference on Transform Techniques and Their Applications, St. Joseph's College, Irinjalakuda, India, 2001, to appear.
- [5] ———, *Boehmians and their Hilbert transforms*, Integral Transform. Spec. Funct. **13** (2002), no. 2, 131–141.
- [6] T. H. Koornwinder (ed.), *Wavelets: An Elementary Treatment of Theory and Applications*, Series in Approximations and Decompositions, vol. 1, World Scientific Publishing, New Jersey, 1993 (French).
- [7] J. Mikusiński and P. Mikusiński, *Quotients de suites et leurs applications dans l'analyse fonctionnelle [Quotients of sequences and their applications in functional analysis]*, C. R. Acad. Sci. Paris Sér. I Math. **293** (1981), no. 9, 463–464 (French).

- [8] P. Mikusiński, *Convergence of Boehmians*, Japan. J. Math. (N.S.) **9** (1983), no. 1, 159–179.
- [9] ———, *Fourier transform for integrable Boehmians*, Rocky Mountain J. Math. **17** (1987), no. 3, 577–582.
- [10] ———, *The Fourier transform of tempered Boehmians*, Fourier Analysis (Orono, Me, 1992), Lecture Notes in Pure and Applied Mathematics, vol. 157, Marcel Dekker, New York, 1994, pp. 303–309.
- [11] ———, *Tempered Boehmians and ultradistributions*, Proc. Amer. Math. Soc. **123** (1995), no. 3, 813–817.
- [12] ———, *On flexibility of Boehmians*, Integral Transform. Spec. Funct. **4** (1996), no. 1-2, 141–146.
- [13] P. Mikusiński, A. Morse, and D. Nemzer, *The two-sided Laplace transform for Boehmians*, Integral Transform. Spec. Funct. **2** (1994), no. 3, 219–230.
- [14] P. Mikusiński and A. Zayed, *The Radon transform of Boehmians*, Proc. Amer. Math. Soc. **118** (1993), no. 2, 561–570.
- [15] D. Nemzer, *Periodic Boehmians*, Int. J. Math. Math. Sci. **12** (1989), no. 4, 685–692.
- [16] ———, *Periodic Boehmians. II*, Bull. Austral. Math. Soc. **44** (1991), no. 2, 271–278.
- [17] ———, *The Laplace transform on a class of Boehmians*, Bull. Austral. Math. Soc. **46** (1992), no. 2, 347–352.
- [18] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, New York, 1987.

R. Roopkumar: School of Mathematics, Madurai Kamaraj University, Madurai 625021, India