

## GAUSSIAN QUADRATURE RULES AND $A$ -STABILITY OF GALERKIN SCHEMES FOR ODE

ALI BENSEBAH, FRANÇOIS DUBEAU, and JACQUES GÉLINAS

Received 20 November 2002

The  $A$ -stability properties of continuous and discontinuous Galerkin methods for solving ordinary differential equations (ODEs) are established using properties of Legendre polynomials and Gaussian quadrature rules. The influence on the  $A$ -stability of the numerical integration using Gaussian quadrature rules involving a parameter is analyzed.

2000 Mathematics Subject Classification: 65L05, 65L20, 65L60.

**1. Introduction.** In this paper, the  $A$ -stability of various (continuous and discontinuous) Galerkin schemes for the solution of an initial value problem in ordinary differential equation (ODE) is analyzed. Even if  $A$ -stability of a method for solving ODE is an old well-studied subject, the contribution of this paper is in the presentation of a new proof of  $A$ -stability in [Section 4.1](#) which points out the link between  $A$ -stability, Legendre polynomials, and Gaussian quadrature rules.

The stability results are obtained using a variety of Gaussian quadrature formulas of integrals defining the Galerkin finite element methods for problems of the form

$$\begin{aligned} \dot{y}(t) &= f(y(t), t), \quad 0 \leq t \leq T, \\ y(0) &= y_0. \end{aligned} \tag{1.1}$$

Continuous and discontinuous Galerkin methods have played an important role in the recently developed approach to global error estimation and control for numerical approximations of ODEs [\[8, 15, 16, 24\]](#). In particular, the stability analysis is an important issue and has motivated our work.

**2. Polynomial approximations and Galerkin methods.** Galerkin methods for [\(1.1\)](#) are based on a variational formulation and use a (continuous or discontinuous) piecewise polynomial approximation of the solution of the ODE. They can be briefly described in the following way [\[3, 5\]](#).

The interval  $[0, T]$  is partitioned into  $N$  intervals  $I_n = [t_{n-1}, t_n]$  ( $n = 1, \dots, N$ ) by specifying the sequence  $\{t_n\}_{n=0}^N$ ,  $0 = t_0 < t_1 < \dots < t_{N-1} < t_N = T$ , of real

numbers. If  $P^K(I_n)$  denotes the set of real polynomials of degree  $K$  on  $I_n$ , we consider the following variational problem.

**PROBLEM 2.1.** Let  $U_0 = y_0$ , and for  $n = 1, \dots, N$ , find  $u_n(\cdot) \in P^K(I_n)$  and  $U_n \in \mathbb{R}$  such that  $\langle Du_n - f(u_n), v_n \rangle_n + [u_n(t_{n-1}) - U_{n-1}]v_n(t_{n-1}) + [U_n - u_n(t_n)]v_n(t_n) = 0$  for all  $v_n(\cdot) \in P^{K+1-L}(I_n)$ , and  $u_n(\cdot)$  is subject to  $L$  additional collocation conditions.

In **Problem 2.1**,  $f(u)$  stands for  $f(u(t), t)$  and

$$\langle f, g \rangle_n = \int_{t_{n-1}}^{t_n} f(\tau)g(\tau)d\tau. \tag{2.1}$$

We say that we have an *approximate problem* if we use exact integration in **Problem 2.1** and we have a *discretized problem* if we use a quadrature rule to deal with the integrals.

Continuous Galerkin methods for ODEs have been introduced in [19, 20] and discontinuous Galerkin methods in [25]. Discontinuous Galerkin methods were first analyzed for linear nonstiff ODEs in [4], and later for nonlinear nonstiff systems in [2, 6]. A general framework and analysis of Galerkin methods for ODEs have been developed in [3, 5]. In particular, existence, uniqueness, and convergence results have been obtained for approximate and discretized problems under appropriate assumptions on  $f(\cdot, \cdot)$ .

Similarly, Galerkin methods for parabolic problems have been analyzed, for example, in [1, 14, 23, 26], and later used for finding adaptive finite element methods for parabolic problems in [9, 10, 11, 12, 13, 17].

**3. Gaussian quadrature rules.** Let  $y$  be a parameter in  $[-1, 1]$  and  $\pi_M(\cdot)$  be the Legendre polynomial of degree  $M$  on  $[-1, 1]$  such that  $\pi_M(1) = 1 = (-1)^M \pi_M(-1)$ . The quadrature rules we consider are based on the following result.

**LEMMA 3.1.** For  $y \in [-1, 1]$ , the  $M$  roots of the polynomial  $\pi_M(\cdot) - y\pi_{M-1}(\cdot)$ , denoted  $\tau_m(y)$  ( $m = 1, \dots, M$ ), are all real and distinct. Moreover,

- (i)  $-1 \leq \tau_1(y) < \dots < \tau_m(y) < \dots < \tau_M(y) \leq 1$ ;
- (ii)  $\tau_m(\cdot) \in C^\infty([-1, 1]; [-1, 1])$  for  $m = 1, \dots, M$ ;
- (iii)  $(d\tau_m/dy)(y) > 0$  for any  $y \in (-1, 1)$ ;
- (iv)  $\tau_1(-1) = -1$  and  $\tau_M(1) = 1$ .

**PROOF.** This result is obtained from the interlacing properties of the roots of Legendre polynomials and from the implicit function theorem. □

If we define the weights

$$\omega_m(y) = \int_{-1}^1 \prod_{\substack{k=1 \\ k \neq m}}^M \frac{\tau - \tau_k(y)}{\tau_m(y) - \tau_k(y)} d\tau, \tag{3.1}$$

for  $m = 1, \dots, M$ , the quadrature rule

$$\int_{-1}^1 \psi(\tau) d\tau \cong \sum_{m=1}^M \omega_m(y) \psi(\tau_m(y)), \tag{3.2}$$

which depends on a parameter  $y \in [-1, 1]$ , is exact for polynomials of degree  $M - 1$ . It follows that the formula is also exact for polynomials of degree  $2M - 2$  and for polynomials of degree  $2M - 1$  if  $y = 0$ . Gauss-Legendre quadrature rules correspond to  $y = 0$ , Gauss-Radau quadrature rules correspond to  $y = \pm 1$ , and for  $0 < |y| < 1$ , we obtain intermediate quadrature rules.

We will use the following notation for the interval  $[-1, 1]$ :

$$\begin{aligned} \langle f, g \rangle &= \int_{-1}^1 f(\tau) g(\tau) d\tau, \\ \langle f, g \rangle^d &= \sum_{m=1}^M \omega_m(y) f(\tau_m(y)) g(\tau_m(y)). \end{aligned} \tag{3.3}$$

We remark the following identities for Legendre polynomials:

$$\langle \pi_i, \pi_j \rangle^d = \langle \pi_i, \pi_j \rangle = \begin{cases} 0 & \text{if } i < j, \\ \frac{2}{2i+1} & \text{if } i = j, \end{cases} \tag{3.4}$$

for  $j = 0, \dots, M - 1$ ,

$$\langle \pi_i, \pi_M \rangle^d = \begin{cases} \langle \pi_i, \pi_M \rangle = 0 & \text{if } i < M - 1, \\ \frac{2y}{2M - 1} & \text{if } i = M - 1, \\ \frac{2y^2}{2M - 1} & \text{if } i = M, \end{cases} \tag{3.5}$$

since  $\pi_M(\tau_m(y)) = y\pi_{M-1}(\tau_m(y))$  and  $\langle \pi_i, \pi_M \rangle^d = y\langle \pi_i, \pi_{M-1} \rangle^d$ . Also, if  $i + j \leq 2M - 1$ , then

$$\begin{aligned} \langle D\pi_i, \pi_j \rangle^d &= \langle D\pi_i, \pi_j \rangle \\ &= \begin{cases} 0 & \text{if } i \leq j, \\ \pi_i \pi_j |_{-1}^1 - \langle \pi_i, D\pi_j \rangle = 1 - (-1)^{i+j} & \text{if } i > j. \end{cases} \end{aligned} \tag{3.6}$$

**4. A-stability analysis.** We will analyze the A-stability properties of the methods corresponding to approximate and discretized problems with respect to the parameter  $y$  for the following cases:

- (1)  $L = 0$ : the completely discontinuous method introduced in [3];
- (2)  $L = 2$  with  $u_n(t_{n-1}) = U_{n-1}$  and  $u_n(t_n) = U_n$ : the continuous method presented in [19, 20];

- (3)  $L = 1$  with  $u_n(t_n) = U_n$ : the discontinuous method described in [25], and also a special case of the discontinuous  $\alpha$ -method of [4] ( $\alpha_n = 1$  for all  $n$ );
- (4)  $L = 1$  with  $u_n(t_n) = U_{n-1}$ : a special case of the discontinuous  $\alpha$ -method of [4] ( $\alpha_n = 0$  for all  $n$ ).

For the  $A$ -stability analysis, we consider the following form of (1.1):

$$\begin{aligned} \dot{y}(t) &= \lambda y(t), \quad 0 \leq t \leq T, \\ y(0) &= y_0, \end{aligned} \tag{4.1}$$

and we solve Problem 2.1 to obtain

$$U_n = R\left(\frac{\lambda h_n}{2}\right)U_{n-1}, \tag{4.2}$$

where  $R(z) = P(z)/Q(z)$  is a rational function and  $h_n = t_n - t_{n-1}$ .

**DEFINITION 4.1.** The region of stability of a method is the set

$$S = \{z \in \mathbb{C} \mid |R(z)| \leq 1\}. \tag{4.3}$$

Let  $\text{Re}(z)$  be the real part of  $z$ ; a method is said to be

- (i) *A-stable* if  $|R(z)| < 1$  whenever  $\text{Re}(z) < 0$ ;
- (ii) *stiff A-stable* if it is  $A$ -stable and  $\lim_{\text{Re}(z) \rightarrow -\infty} |R(z)| = 0$ .

To obtain (4.2) from Problem 2.1, let

$$\pi_{ni}(t) = \pi_i\left(\frac{2t - (t_{n-1} + t_n)}{t_n - t_{n-1}}\right) \tag{4.4}$$

be the polynomial of degree  $i$  defined on  $I_n = [t_{n-1}, t_n]$  normalized such that  $\pi_{ni}(t_n) = 1 = (-1)^i \pi_{ni}(t_{n-1})$ . Then  $\{\pi_{ni}\}_{i=0}^K$  and  $\{\pi_{nj}\}_{j=0}^{K+1-L}$  form a basis for  $P^K(I_n)$  and  $P^{K+1-L}(I_n)$ . Hence, the polynomial  $u_n(\cdot)$  that we look for can be written as  $u_n(t) = \sum_{i=0}^K a_{ni} \pi_{ni}(t)$ . Then Problem 2.1 becomes the following problem.

**PROBLEM 4.2.** Let  $U_0 = y_0$ , and for  $n = 1, \dots, N$ , find  $a_{n0}, \dots, a_{nK}, U_n \in \mathbb{R}$ , such that  $\sum_{i=0}^K [(\lambda h_n/2) \langle \pi_i, \pi_j \rangle - \langle D\pi_i, \pi_j \rangle + 1 - (-1)^{i+j}] a_{ni} = U_n - (-1)^j U_{n-1}$  for  $j = 0, \dots, K + 1 - L$ , and  $u_n(\cdot)$  is subject to  $L$  additional conditions.

In the sequel of this paper, we will use the notation  $R_{L,K}(z; \gamma)$  for the amplifying factor  $R(\lambda h_n/2)$ , and  $\lambda h_n/2$  is replaced by  $z$ .

**4.1. The case  $L = 0$ .** For the interval  $I_n$ , we use the quadrature formula (3.2) with  $M = K + 1$ . Consequently,  $\langle D\pi_i, \pi_j \rangle^d$  is always exact, and  $\langle \pi_i, \pi_j \rangle^d$  is exact for  $i + j \leq 2K$  or for  $i + j \leq 2K + 1$  if  $\gamma = 0$ . Hence, for the stability analysis, the approximate problem is equivalent to the discretized problem for  $\gamma = 0$ .



**LEMMA 4.4.** For  $\ell \geq k$ ,  $A_{k\ell}$  is a polynomial in  $z^2$ . More precisely,

(i) if  $\ell - k = 2n + 1$ , then  $A_{k\ell} = z^{2n+2} + p_n(z^2)$ ,

(ii) if  $\ell - k = 2n$ , then  $A_{k\ell} = (n + 1)(2k + 2n + 1)z^{2n} + p_{n-1}(z^2)$ ,

for  $n = 0, 1, 2, \dots$ , and where  $p_j(z)$  is a polynomial of degree  $j$  in  $z$  ( $p_{-1}(z) = 0$ ).

As a direct consequence of Cramer's rule, we have

$$a_{n0} = U_{n-1} \frac{A_{1K} + yzA_{1K-1}}{(A_{0K} + yzA_{0K-1}) - z(A_{1K} + yzA_{1K-1})} \tag{4.9}$$

and from the properties of  $A_{k\ell}$ , we obtain

$$R_{0K}(z; y) = \frac{(A_{0K} + yzA_{0K-1}) + z(A_{1K} + yzA_{1K-1})}{(A_{0K} + yzA_{0K-1}) - z(A_{1K} + yzA_{1K-1})} \tag{4.10a}$$

or

$$R_{0K}(z; y) = \frac{(A_{0K} + zA_{1K}) + yz(A_{0K-1} + zA_{1K-1})}{(A_{0K} - zA_{1K}) + yz(A_{0K-1} - zA_{1K-1})}. \tag{4.10b}$$

Let

$$Q_K(z) = A_{0K-1} + zA_{1K-1} \tag{4.11}$$

for  $K = 0, 1, 2, \dots$ . From [Lemma 4.4](#),

$$Q_K(-z) = A_{0K-1} - zA_{1K-1} \tag{4.12}$$

and we have the following results.

**LEMMA 4.5.** The polynomials  $Q_k(z)$  can be generated recursively by  $Q_0(z) = 1$ ,  $Q_1(z) = 1 + z$ , and for  $K \geq 1$

$$Q_{K+1}(z) = (2K + 1)Q_K(z) + z^2Q_{K-1}(z). \tag{4.13}$$

**PROOF.** The proof is a direct consequence of [Lemma 4.3](#). □

**LEMMA 4.6.** For  $K \geq 0$ ,  $Q_K(z) = \sum_{i=0}^K ((2K - i)! / 2^{K-i}(K - i)!) z^i$ .

**PROOF.** Since  $Q_K(z)$  is a polynomial of degree  $K$ , let  $Q_K(z) = \sum_{i=0}^K a_{Ki} z^i$ . Then  $a_{00} = a_{10} = a_{11} = 1$ , and for  $K \geq 2$ , we have

$$a_{Ki} = (2K - 1)a_{K-1,i} + a_{K-2,i-2}, \tag{4.14}$$

for  $i = 0, \dots, K$ , considering  $a_{Kj} = 0$  if  $j < 0$  or  $j > K$ . Then the result follows by induction. □

**THEOREM 4.7.** Let  $P_{K+1}(z; \gamma) = Q_{K+1}(z) + \gamma z Q_K(z)$ . Then

$$P_{K+1}(z; \gamma) = \sum_{i=0}^{K+1} \frac{(2(K+1)-i)!}{2^{K+1-i}(K+1-i)!i!} \left[ 1 + \gamma \frac{i}{2(K+1)-i} \right] z^i, \tag{4.15}$$

$$R_{0K}(z; \gamma) = \frac{P_{K+1}(z; \gamma)}{P_{K+1}(-z; -\gamma)}.$$

Moreover, the following limits exist:

$$\lim_{|z| \rightarrow -\infty} |R_{0K}(z; \gamma)| = \frac{1+\gamma}{1-\gamma} \quad \text{for } \gamma \in [-1, 1), \tag{4.16}$$

$$\lim_{|z| \rightarrow +\infty} |R_{0K}(z; 1)| = +\infty \quad \text{for } \gamma = 1.$$

From the properties of  $A_{k\ell}$  of Lemma 4.3, we obtain the following expression.

**LEMMA 4.8.** For any complex number  $z$ ,

$$z(A_{1K} + \gamma z A_{1K-1}) \overline{(A_{0K} + \gamma z A_{0K-1})}$$

$$= \gamma |z|^{2K+2} + \sum_{i=0}^K (2i+1) z_i |z|^{2i} |A_{i+1,K} + \gamma z A_{i+1,K-1}|^2, \tag{4.17}$$

where  $z_i = z$  for  $i$  even and  $z_i = \bar{z}$  for  $i$  odd.

**LEMMA 4.9.** Let  $X$  and  $Y$  be two complex numbers, then

$$\frac{|X+Y|}{|X-Y|} < 1 \quad \text{iff } \operatorname{Re}(Y\bar{X}) < 0. \tag{4.18}$$

**THEOREM 4.10.** Let  $L = 0$  and  $K \geq 0$ .

(i) The method corresponding to the approximate problem (4.5) is A-stable but not stiff A-stable.

(ii) Let  $\gamma \in [-1, 1]$  and  $M = K + 1$  in (3.2). Then the method corresponding to the discretized problem (4.5) is A-stable for  $\gamma = [-1, 0]$  and stiff A-stable for  $\gamma = -1$ .

**PROOF.** We recall that the result for the approximate problem corresponds to the result for the discretized problem for  $\gamma = 0$ . For the A-stability, using Lemma 4.9, we observe that  $|R_{0K}(z; \gamma)| < 1$  is equivalent to

$$\operatorname{Re} \{ z(A_{1K} + \gamma z A_{1K-1}) \overline{(A_{0K} + \gamma z A_{0K-1})} \} < 0. \tag{4.19}$$

From (4.10) and the expression in Lemma 4.8, (4.19) is satisfied for  $\operatorname{Re}(z) < 0$  if  $\gamma \leq 0$ . From the limits of Theorem 4.7, it follows that for any  $\gamma > 0$ , there exists  $z$  such that  $\operatorname{Re}(z) < 0$  and  $|R_{0K}(z; \gamma)| > 1$ . Then the result on A-stability follows. The stiff A-stability for  $\gamma = -1$  follows also from Theorem 4.7.  $\square$

**EXAMPLE 4.11.** (i)  $K = 0$ ,

$$R_{00}(z; \gamma) = \frac{1 + (1 + \gamma)z}{1 - (1 - \gamma)z}; \tag{4.20}$$

(ii)  $K = 1$ ,

$$R_{01}(z; \gamma) = \frac{3 + (3 + \gamma)z + (1 + \gamma)z^2}{3 - (3 - \gamma)z + (1 - \gamma)z^2}; \tag{4.21}$$

(iii)  $K = 2$ ,

$$R_{02}(z; \gamma) = \frac{15 + (15 + 3\gamma)z + (6 + 3\gamma)z^2 + (1 + \gamma)z^3}{15 - (15 - 3\gamma)z + (6 - 3\gamma)z^2 - (1 - \gamma)z^3}; \tag{4.22}$$

(iv)  $K = 3$ ,

$$R_{03}(z; \gamma) = \frac{105 + (105 + 15\gamma)z + (45 + 15\gamma)z^2 + (10 + 6\gamma)z^3 + (1 + \gamma)z^4}{105 - (105 - 15\gamma)z + (45 - 15\gamma)z^2 - (10 - 6\gamma)z^3 + (1 - \gamma)z^4}. \tag{4.23}$$

**REMARK 4.12.** Since  $R_{0K}(z; \gamma) = 1/R_{0K}(-z; -\gamma)$ , the stability region for  $-1 \leq \gamma < 0$  is the exterior of the mirror image in the imaginary axis of the corresponding region for  $-\gamma$ . For  $\gamma = 0$ , the stability region is the left half-plane. Thus, it is sufficient to describe the bounded regions for  $0 < \gamma \leq 1$ .

**REMARK 4.13.** When  $\gamma = -1, 0$ , and  $1$ , the ratios of [Example 4.11](#) correspond to the subdiagonal, diagonal, and superdiagonal element of the Padé table for  $e^{2z}$ , respectively. Other values of  $\gamma$  give intermediate rational approximations of the exponential. These rational approximations of  $e^{2z}$  have already been analyzed; they appear in [\[7, 18, 27, 28\]](#).

**REMARK 4.14.** The proof of the  $A$ -stability presented here seems to be new in the sense that it uses only elementary properties of Legendre polynomials and Gaussian quadrature rules. However, we do not obtain the stability region in the case  $\gamma \in (0, 1]$ , we obtain only the fact that it is not  $A$ -stable. One way to obtain the stability region is to use the order stars approach [\[21, 22\]](#).

**4.2. The case  $L = 2$ :**  $u_n(t_{n-1}) = U_{n-1}$  and  $u_n(t_n) = U_n$ . In this case, we use the quadrature formula [\(3.2\)](#) with  $M = K$ . Hence,  $\langle D\pi_i, \pi_j \rangle^d$  is always exact, and  $\langle \pi_i, \pi_j \rangle^d$  is exact for  $i + j \leq 2K - 2$  or for  $i + j \leq 2K - 1$  if  $\gamma = 0$ .

Then the system is

$$\sum_{i=0}^j [1 - (-1)^{i+j}] a_{ni} + \frac{2z}{2j+1} a_{nj} = U_n - (-1)^j U_{n-1} \tag{4.24a}$$

for  $j = 0, \dots, K - 2$ ,

$$\sum_{i=0}^{K-1} [1 - (-1)^{i+K-1}] a_{ni} + \frac{2z}{2K-1} a_{nK-1} + \frac{2yz}{2K-1} a_{nK} = U_n - (-1)^{K-1} U_{n-1}, \tag{4.24b}$$

$$\sum_{i=0}^K (-1)^i a_{ni} = U_{n-1}, \tag{4.24c}$$

$$\sum_{i=0}^K a_{ni} = U_n, \tag{4.24d}$$

which is equivalent to

$$U_n = U_{n-1} + 2z a_{n0}, \tag{4.25a}$$

$$\sum_{i=0}^K (-1)^{i+1} a_{ni} = -U_{n-1}, \tag{4.25b}$$

$$-2z a_{nj} + (2j + 1) \sum_{i=j+1}^K [1 - (-1)^{i+j}] a_{ni} = 0, \tag{4.25c}$$

for  $j = 0, 2, \dots, K - 2$ , and

$$-z a_{nK-1} + [(2K - 1) - yz] a_{nK} = 0. \tag{4.25d}$$

Solving (4.25) for  $a_{n0}$  using Cramer's rule, we obtain

$$a_{n0} = U_{n-1} \frac{A_{1K-1} - yz A_{1K-2}}{(A_{0K-1} - yz A_{0K-2}) - z(A_{1K-1} - yz A_{1K-2})}, \tag{4.26}$$

$$R_{2K}(z; \gamma) = \frac{(A_{0K-1} - yz A_{0K-2}) + z(A_{1K-1} - yz A_{1K-2})}{(A_{0K-1} - yz A_{0K-2}) - z(A_{1K-1} - yz A_{1K-2})},$$

and the next result follows.

**THEOREM 4.15.** For any  $\gamma \in [-1, 1]$ ,

$$R_{2,K+1}(z; \gamma) = R_{0K}(z; -\gamma). \tag{4.27}$$

**THEOREM 4.16.** Let  $L = 2$ ,  $u_n(t_n) = U_n$ ,  $u_n(t_{n-1}) = U_{n-1}$ , and  $K \geq 1$ .

(1) The method corresponding to the approximate problem (4.24a) is A-stable but not stiff A-stable.

(2) Let  $\gamma \in [-1, 1]$  and  $M = K$  in (3.2). Then the method corresponding to the discretized problem (4.24a) is A-stable for  $\gamma \in [0, 1]$  and stiff A-stable for  $\gamma = 1$ .

**4.3. The case  $L = 1$ :**  $u_n(t_n) = U_n$ . In this case, using the quadrature rule (3.2) with  $M = K + 1$ , any term of the forms  $\langle \pi_i, \pi_j \rangle^d$  or  $\langle D\pi_i, \pi_j \rangle^d$  is integrated exactly for  $i, j \leq K$ . Hence, the approximate problem and the discretized problems have the same A-stability property.

The system is

$$\sum_{i=0}^j [1 - (-1)^{i+j}] a_{ni} + \frac{2z}{2j+1} a_{nj} = U_n - (-1)^j U_{n-1} \tag{4.28a}$$

for  $j = 0, \dots, K$ , and

$$\sum_{i=0}^K a_{ni} = U_n. \tag{4.28b}$$

But, this system is equivalent to

$$U_n = U_{n-1} + 2z a_{n0}, \tag{4.29a}$$

$$(1 - z) a_{n0} + \frac{z}{3} a_{n1} = U_{n-1}, \tag{4.29b}$$

$$-\frac{z}{2j-1} a_{nj-1} + a_{nj} + \frac{z}{2j+1} a_{nj+1} = 0, \tag{4.29c}$$

for  $j = 1, \dots, K-1$ , and

$$-\frac{z}{2K-1} a_{nK-1} + \left(1 - \frac{z}{2K+1}\right) a_{nK} = 0. \tag{4.29d}$$

Hence,

$$a_{n0} = U_{n-1} \frac{A_{1K} - zA_{1K-1}}{(A_{0K} - zA_{0K-1}) - z(A_{1k} - zA_{1K-1})}, \tag{4.30}$$

$$R_{1K}^r(z; \gamma) = \frac{(A_{0K} - zA_{0K-1}) + z(A_{1K} - zA_{1K-1})}{(A_{0K} - zA_{0K-1}) - z(A_{1K} - zA_{1K-1})},$$

and we have the next result.

**THEOREM 4.17.** For any  $\gamma \in [-1, 1]$ ,

$$R_{1K}^r(z; \gamma) = R_{0K}(z; -1), \tag{4.31}$$

and the methods corresponding to the approximate and the discretized problems (4.28) are stiff A-stable.

**4.4. The case  $L = 1$ :**  $u_n(t_{n-1}) = U_{n-1}$ . In this case, we have

$$\sum_{i=0}^j [1 - (-1)^{i+j}] a_{ni} + \frac{2z}{2j+1} a_{nj} = U_n - (-1)^j U_{n-1} \tag{4.32a}$$

for  $j = 0, \dots, K$ , and

$$\sum_{i=0}^K (-1)^i a_{ni} = U_{n-1}. \tag{4.32b}$$

This system is equivalent to

$$U_n = U_{n-1} + 2z a_{n0}, \tag{4.33a}$$

$$(1 - z)a_{n0} + \frac{z}{3}a_{n1} = U_{n-1}, \tag{4.33b}$$

$$-\frac{z}{2j-1}a_{nj-1} + a_{nj} + \frac{z}{2j+1}a_{nj+1} = 0, \tag{4.33c}$$

for  $j = 1, \dots, K-1$ , and

$$-\frac{z}{2K-1}a_{nK-1} + \left(1 + \frac{z}{2K+1}\right)a_{nK} = 0. \tag{4.33d}$$

Then

$$a_{n0} = U_{n-1} \frac{A_{1K} + zA_{1K-1}}{(A_{0K} + zA_{0K-1}) - z(A_{1K} + zA_{1K-1})}, \tag{4.34}$$

$$R_{1K}^\ell(z; \gamma) = \frac{(A_{0K} + zA_{0K-1}) + z(A_{1K} + zA_{1K-1})}{(A_{0K} + zA_{0K-1}) - z(A_{1K} + zA_{1K-1})},$$

and we obtain the last result.

**THEOREM 4.18.** For any  $\gamma \in [-1, 1]$ ,

$$R_{1K}^\ell(z; \gamma) = R_{0K}(z; 1), \tag{4.35}$$

and the methods corresponding to the approximate and the discretized problems (4.32) are not A-stable.

**ACKNOWLEDGMENT.** The research of the second author has been supported in part by the Natural Sciences and Engineering Research Council of Canada (NSERC).

**REFERENCES**

[1] G. A. Baker, J. H. Bramble, and V. Thomée, *Single step Galerkin approximations for parabolic problems*, Math. Comp. **31** (1977), no. 140, 818-847.  
 [2] A. Bensebah, F. Dubeau, and J. Gélinas, *Résultats d'existence et de superconvergence pour la méthode des  $\alpha$*  [Existence and superconvergence results for the  $\alpha$ -method], Ann. Sci. Math. Québec **17** (1993), no. 2, 115-138 (French).  
 [3] M. C. Delfour and F. Dubeau, *Discontinuous polynomial approximations in the theory of one-step, hybrid and multistep methods for nonlinear ordinary differential equations*, Math. Comp. **47** (1986), no. 175, 169-189, with a supplement S1-S8.  
 [4] M. C. Delfour, W. Hager, and F. Trochu, *Discontinuous Galerkin methods for ordinary differential equations*, Math. Comp. **36** (1981), no. 154, 455-473.  
 [5] F. Dubeau, *Approximations polynômiales par morceaux des équations différentielles*, Ph.D. thesis, University of Montréal, Montréal, Canada, 1981.  
 [6] ———, *Approximation discontinue des équations différentielles ordinaires: la méthode des  $\alpha$*  [Discontinuous approximation of ordinary differential equations: the method of  $\alpha$ 's], Ann. Sci. Math. Québec **10** (1986), no. 2, 153-179 (French).

- [7] B. L. Ehle, *On Padé approximation to the exponential function and A-stable methods for the numerical solution of initial value problems*, Ph.D. thesis, University of Waterloo, Waterloo, Canada, 1969.
- [8] K. Eriksson, D. Estep, P. Hansbo, and C. Johnson, *Introduction to adaptive methods for differential equations*, Acta Numerica, 1995, Acta Numer., Cambridge University Press, Cambridge, 1995, pp. 105–158.
- [9] K. Eriksson and C. Johnson, *Adaptive finite element methods for parabolic problems. I. A linear model problem*, SIAM J. Numer. Anal. **28** (1991), no. 1, 43–77.
- [10] ———, *Adaptive finite element methods for parabolic problems. II. Optimal error estimates in  $L_\infty L_2$  and  $L_\infty L_\infty$* , SIAM J. Numer. Anal. **32** (1995), no. 3, 706–740.
- [11] ———, *Adaptive finite element methods for parabolic problems. IV. Nonlinear problems*, SIAM J. Numer. Anal. **32** (1995), no. 6, 1729–1749.
- [12] ———, *Adaptive finite element methods for parabolic problems. V. Long-time integration*, SIAM J. Numer. Anal. **32** (1995), no. 6, 1750–1763.
- [13] K. Eriksson, C. Johnson, and S. Larsson, *Adaptive finite element methods for parabolic problems. VI. Analytic semigroups*, SIAM J. Numer. Anal. **35** (1998), no. 4, 1315–1325.
- [14] K. Eriksson, C. Johnson, and V. Thomée, *Time discretization of parabolic problems by the discontinuous Galerkin method*, RAIRO Modél. Math. Anal. Numér. **19** (1985), no. 4, 611–643.
- [15] D. Estep, *A posteriori error bounds and global error control for approximation of ordinary differential equations*, SIAM J. Numer. Anal. **32** (1995), no. 1, 1–48.
- [16] D. Estep and D. French, *Global error control for the continuous Galerkin finite element method for ordinary differential equations*, RAIRO Modél. Math. Anal. Numér. **28** (1994), no. 7, 815–852.
- [17] D. Estep, D. H. Hodges, and M. Warner, *Computational error estimation and adaptive error control for a finite element solution of launch vehicle trajectory problems*, SIAM J. Sci. Comput. **21** (1999/00), no. 4, 1609–1631.
- [18] E. Hairer, *Constructive characterization of A-stable approximations to  $\exp(z)$  and its connection with algebraically stable Runge-Kutta methods*, Numer. Math. **39** (1982), no. 2, 247–258.
- [19] B. L. Hulme, *Discrete Galerkin and related one-step methods for ordinary differential equations*, Math. Comp. **26** (1972), 881–891.
- [20] ———, *One-step piecewise polynomial Galerkin methods for initial value problems*, Math. Comp. **26** (1972), 415–426.
- [21] A. Iserles, *Order stars, approximations and finite differences. I. The general theory of order stars*, SIAM J. Math. Anal. **16** (1985), no. 3, 559–576.
- [22] ———, *Order stars, approximations and finite differences. II. Theorems in approximation theory*, SIAM J. Math. Anal. **16** (1985), no. 4, 785–802.
- [23] P. Jamet, *Galerkin-type approximations which are discontinuous in time for parabolic equations in a variable domain*, SIAM J. Numer. Anal. **15** (1978), no. 5, 912–928.
- [24] C. Johnson, *Error estimates and adaptive time-step control for a class of one-step methods for stiff ordinary differential equations*, SIAM J. Numer. Anal. **25** (1988), no. 4, 908–926.
- [25] P. Lasaint and P.-A. Raviart, *On a finite element method for solving the neutron transport equation*, Mathematical Aspects of Finite Elements in Partial Differential Equations (Proc. Sympos., Math. Res. Center, Univ. Wisconsin,

Madison, Wis., 1974) (C. DeBoor, ed.), Academic Press, New York, 1974, pp. 89-123.

- [26] M. Luskin and R. Rannacher, *On the smoothing property of the Galerkin method for parabolic equations*, SIAM J. Numer. Anal. **19** (1982), no. 1, 93-113.
- [27] S. P. Nørsett, *C-polynomials for rational approximation to the exponential function*, Numer. Math. **25** (1975/76), no. 1, 39-56.
- [28] S. P. Nørsett and G. Wanner, *Perturbed collocation and Runge-Kutta methods*, Numer. Math. **38** (1981/82), no. 2, 193-208.

Ali Bensebah: Commission Scolaire de Montréal (CSDM), 3737 rue Sherbrooke Est, Montréal, Quebec, Canada H1X 3B3

*E-mail address:* [bensebaha@csgm.qc.ca](mailto:bensebaha@csgm.qc.ca)

François Dubeau: Département de Mathématiques et d'Informatique, Faculté des Sciences, Université de Sherbrooke, 2500 boulevard Université, Sherbrooke, Quebec, Canada J1K 2R1

*E-mail address:* [francois.dubeau@dm1.usherb.ca](mailto:francois.dubeau@dm1.usherb.ca)

Jacques Gélinas: Defence Research and Development Canada-Ottawa, 3701 Carling Avenue, Ottawa, Ontario, Canada K1A 0Z4

*E-mail address:* [jacques.gelinas@drdc-rddc.gc.ca](mailto:jacques.gelinas@drdc-rddc.gc.ca)



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

