PROPERTIES OF CERTAIN p -VALENTLY CONVEX FUNCTIONS

DINGGONG YANG and SHIGEYOSHI OWA

Received 10 September 2002

A subclass $\mathcal{C}_p(\lambda, \mu)$ ($p \in \mathbb{N}$, $0 < \lambda < 1$, $-\lambda \leq \mu < 1$) of p -valently convex functions in the open unit disk $\mathbb{U}$ is introduced. The object of the present paper is to discuss some interesting properties of functions belonging to the class $\mathcal{C}_p(\lambda, \mu)$.

2000 Mathematics Subject Classification: 30C45.

1. Introduction. Let $\mathcal{A}_p$ denote the class of functions $f(z)$ of the form

$$f(z) = z^p + \sum_{n=1}^{\infty} a_{p+n} z^{p+n} \quad (p \in \mathbb{N} = \{1, 2, 3, \dots\}) \quad (1.1)$$

which are analytic in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$. A function $f(z)$ in $\mathcal{A}_p$ is said to be p -valently convex of order α if it satisfies

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > p\alpha \quad (z \in \mathbb{U}) \quad (1.2)$$

for some α ($0 \leq \alpha < 1$). We denote by $\mathcal{H}_p(\alpha)$ the subclass of $\mathcal{A}_p$ consisting of functions which are p -valently convex of order α in $\mathbb{U}$. In particular, we denote $\mathcal{H}_1(0) = \mathcal{K}$.

A function $f(z) \in \mathcal{A}_1$ is said to be uniformly convex in $\mathbb{U}$ if $f(z)$ is in the class $\mathcal{K}$ and has the property that the image arc $f(\gamma)$ is convex for every circular arc γ contained in $\mathbb{U}$ with center at $t \in \mathbb{U}$. We also denote by $\mathcal{UK}$ the subclass of $\mathcal{A}_1$ consisting of all uniformly convex functions in $\mathbb{U}$. Goodman [2] has introduced the class $\mathcal{UK}$ and given that $f(z) \in \mathcal{A}_1$ belongs to the class $\mathcal{UK}$ if and only if

$$\operatorname{Re} \left\{ 1 + (z-t) \frac{f''(z)}{f'(z)} \right\} \geq 0 \quad ((z, t) \in \mathbb{U} \times \mathbb{U}). \quad (1.3)$$

Ma and Minda [3] and Rønning [5] have showed a more applicable characterization for $\mathcal{UK}$. We state this as the following theorem.

THEOREM 1.1. *Let $f(z) \in \mathcal{A}_1$. Then $f(z) \in \mathcal{UK}$ if and only if*

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \left| \frac{zf''(z)}{f'(z)} \right| \quad (z \in \mathbb{U}). \quad (1.4)$$

In view of [Theorem 1.1](#), Owa [\[4\]](#) considered a subclass $\mathcal{UK}(\mu)$ ($-1 < \mu < 1$) of $\mathcal{A}_1$. A function $f(z) \in \mathcal{A}_1$ is said to be a member of the class $\mathcal{UK}(\mu)$ ($-1 < \mu < 1$) if and only if

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} - \mu > \left| \frac{zf''(z)}{f'(z)} \right| \quad (z \in \mathbb{U}). \quad (1.5)$$

In this paper, we investigate the following subclass of $\mathcal{A}_p$.

DEFINITION 1.2. A function $f(z) \in \mathcal{A}_p$ is said to be a member of the class $\mathcal{C}_p(\lambda, \mu)$ if

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} - p\mu > \lambda \left| 1 + \frac{zf''(z)}{f'(z)} - p \right| \quad (z \in \mathbb{U}) \quad (1.6)$$

for some λ ($0 < \lambda < 1$) and μ ($-\lambda \leq \mu < 1$).

Let $f(z)$ and $g(z)$ be analytic in $\mathbb{U}$. Then we say that $f(z)$ is subordinate to $g(z)$ in $\mathbb{U}$, written $f(z) < g(z)$, if there exists an analytic function $w(z)$ in $\mathbb{U}$ such that $|w(z)| \leq |z|$ and $f(z) = g(w(z))$. If $g(z)$ is univalent in $\mathbb{U}$, then the subordination $f(z) < g(z)$ is equivalent to $f(0) = g(0)$ and $f(\mathbb{U}) \subset g(\mathbb{U})$.

2. Subordination properties. Our first result for properties of functions $f(z) \in \mathcal{A}_p$ is contained in the following theorem.

THEOREM 2.1. A function $f(z) \in \mathcal{C}_p(\lambda, \mu)$ if and only if

$$1 + \frac{zf''(z)}{f'(z)} < h(z) \quad (2.1)$$

with

$$h(z) = p + \frac{p(1-\mu)}{2\sin^2\beta} \left\{ \left(\frac{1+\sqrt{z}}{1-\sqrt{z}} \right)^{2\beta/\pi} + \left(\frac{1-\sqrt{z}}{1+\sqrt{z}} \right)^{2\beta/\pi} - 2 \right\} \quad (\beta = \arccos\lambda). \quad (2.2)$$

PROOF. Let $1 + zf''(z)/f'(z) = w$ and $w = u + iv$. Then inequality (1.6) can be written as

$$u - p\mu > \lambda\sqrt{(u-p)^2 + v^2}. \quad (2.3)$$

By computing, we find that inequality (2.3) is equivalent to

$$\left(u + \frac{p(\lambda^2 - \mu)}{1 - \lambda^2} \right)^2 - \frac{\lambda^2}{1 - \lambda^2} v^2 > \left(\frac{p\lambda(1 - \mu)}{1 - \lambda^2} \right)^2, \quad (2.4)$$

$$u > \frac{p(\lambda + \mu)}{1 + \lambda}. \quad (2.5)$$

Thus the domain of the values of $1 + zf''(z)/f'(z)$ for $z \in \mathbb{U}$ is contained in

$$\mathbb{D} = \{w = u + iv : u \text{ and } v \text{ satisfy (2.4) and (2.5)}\}. \quad (2.6)$$

In order to prove our theorem, it suffices to show that the function $h(z)$ given by (2.2) maps $\mathbb{U}$ conformally onto the domain $\mathbb{D}$.

Consider the transformations

$$\begin{aligned} w_1 &= \frac{1-\lambda^2}{p(1-\mu)}w + \frac{\lambda^2-\mu}{1-\mu}, \\ t &= \frac{1}{2}\left(w_2^{\pi/\beta} + w_2^{-\pi/\beta}\right), \end{aligned} \quad (2.7)$$

where $\beta = \arccos \lambda$ and $w_2 = w_1 + \sqrt{w_1^2 - 1}$ is the inverse function of

$$w_1 = \frac{w_2 + 1/w_2}{2}. \quad (2.8)$$

It is easy to verify that composite function $t = t(w)$ maps $\mathbb{D}^+$ defined by

$$\mathbb{D}^+ = \{w = u + iv : u \text{ and } v \text{ satisfy (2.4), (2.5), and } v > 0\} \quad (2.9)$$

conformally onto the upper-half plane $\text{Im}(t) > 0$ so that $w = p$ corresponds to $t = 1$ and $w = p(\lambda + \mu)/(1 + \lambda)$ to $t = -1$. With the help of the symmetry principle, this function $t = t(w)$ maps $\mathbb{D}$ conformally onto the domain

$$\mathbb{G} = \{t : |\arg(t + 1)| < \pi\}. \quad (2.10)$$

Since

$$t = 2\left(\frac{1+z}{1-z}\right)^2 - 1 \quad (2.11)$$

maps $\mathbb{U}$ onto $\mathbb{G}$, we see that

$$\begin{aligned} w &= p + \frac{p(1-\mu)}{2(1-\lambda^2)} \left\{ \left(t + \sqrt{t^2 - 1}\right)^{\beta/\pi} + \left(t + \sqrt{t^2 - 1}\right)^{-\beta/\pi} - 2 \right\} \\ &= p + \frac{p(1-\mu)}{2\sin^2 \beta} \left\{ \left(\frac{1+\sqrt{z}}{1-\sqrt{z}}\right)^{2(\beta/\pi)} + \left(\frac{1-\sqrt{z}}{1+\sqrt{z}}\right)^{2(\beta/\pi)} - 2 \right\} \\ &= h(z) \end{aligned} \quad (2.12)$$

maps $\mathbb{U}$ onto $\mathbb{D}$ with $h(0) = p$. Hence the proof of the theorem is completed. $\square$

Theorem 2.1 gives the following corollaries.

COROLLARY 2.2. *If $f(z) \in \mathcal{C}_p(\lambda, \mu)$, then $f(z) \in \mathcal{H}_p((\lambda + \mu)/(1 + \lambda))$ and the order $(\lambda + \mu)/(1 + \lambda)$ is sharp with the extremal function*

$$f_0(z) = p \int_0^z \left(t_2^{p-1} \exp \int_0^{t_2} \frac{h(t_1) - p}{t_1} dt_1 \right) dt_2, \quad (2.13)$$

where $h(z)$ is given by (2.2).

PROOF. Using (2.5) in the proof of Theorem 2.1 and noting that

$$\operatorname{Re} \left(1 + \frac{zf_0''(z)}{f_0'(z)} \right) = \operatorname{Re} (h(z)) \rightarrow p \frac{\lambda + \mu}{1 + \lambda} \quad (2.14)$$

as $z = \operatorname{Re}(z) \rightarrow -1$, we have the corollary. $\square$

COROLLARY 2.3. If $f(z) \in \mathbb{C}_p(\lambda, \mu)$ and $-\lambda < \mu < \lambda < 1$, then

$$\left| \arg \left(1 + \frac{zf''(z)}{f'(z)} \right) \right| < \arctan \left(\frac{1 - \mu}{\sqrt{\lambda^2 - \mu^2}} \right) \quad (z \in \mathbb{U}). \quad (2.15)$$

The bound in (2.15) is sharp with the extremal function $f_0(z)$ given by (2.13).

PROOF. Let the function $h(z)$ be defined by (2.4). Then $h(\mathbb{U}) = \mathbb{D}$ and an easy calculation yields that

$$\min \{ \theta : |\arg(h(z))| < \theta (z \in \mathbb{U}) \} = \arctan \left(\frac{1 - \mu}{\sqrt{\lambda^2 - \mu^2}} \right) \quad (2.16)$$

for $-\lambda < \mu < \lambda < 1$. Therefore, the corollary follows immediately from Theorem 2.1. $\square$

Next we derive the following theorem.

THEOREM 2.4. Let $f(z) \in \mathbb{C}_p(\lambda, \mu)$ and let $h(z)$ be defined by (2.2). Then

$$\frac{f'(z)}{pz^{p-1}} < \exp \int_0^z \frac{h(t) - p}{t} dt, \quad (2.17)$$

$$\left| \frac{f'(z)}{pz^{p-1}} \right| \leq \exp \int_0^1 \frac{h(\rho) - p}{\rho} d\rho \quad (z \in \mathbb{U}). \quad (2.18)$$

The bound in (2.18) is sharp with the extremal function $f_0(z)$ given by (2.13).

PROOF. Since the function $h(z) - p$ is univalent and starlike (with respect to the origin), by Theorem 2.1 and the result due to Suffridge [6, Theorem 3], we have

$$\log \left(\frac{f'(z)}{pz^{p-1}} \right) = \int_0^z \left(\frac{f''(t)}{f'(t)} - \frac{p-1}{t} \right) dt < \int_0^z \frac{h(t) - p}{t} dt, \quad (2.19)$$

which implies the subordination (2.17).

Furthermore, noting that the univalent function $h(z)$ maps the disk $|z| < \rho$ ($0 < \rho \leq 1$) onto the domain which is convex and symmetric with respect to the real axis, we deduce that

$$\operatorname{Re} \int_0^z \frac{h(t) - p}{t} dt = \int_0^1 \frac{\operatorname{Re} \{h(\rho z) - p\}}{\rho} d\rho \leq \int_0^1 \frac{h(\rho) - p}{\rho} d\rho \quad (2.20)$$

for $z \in \mathbb{U}$. Thus inequality (2.18) follows from (2.19) and (2.20). $\square$

REMARK 2.5. If we let $\beta = \pi/4$ and $x = ((1 + \sqrt{\rho})/(1 - \sqrt{\rho}))^{1/2}$ ($0 \leq \rho < 1$), then

$$\begin{aligned} \int_0^1 \left\{ \left(\frac{1 + \sqrt{\rho}}{1 - \sqrt{\rho}} \right)^{2(\beta/\pi)} + \left(\frac{1 - \sqrt{\rho}}{1 + \sqrt{\rho}} \right)^{2(\beta/\pi)} - 2 \right\} \frac{d\rho}{\rho} \\ = 8 \int_1^{+\infty} \left(\frac{x}{x^2 + 1} - \frac{1}{x + 1} \right) dx = 4 \log 2. \end{aligned} \quad (2.21)$$

Thus, as the special case of [Theorem 2.4](#), we have that if $f(z) \in \mathcal{C}_p(1/\sqrt{2}, \mu)$ ($-1/\sqrt{2} \leq \mu < 1$), then

$$\left| \frac{f'(z)}{pz^{p-1}} \right| \leq 16^{p(1-\mu)} \quad (z \in \mathbb{U}) \quad (2.22)$$

and the result is sharp.

3. Coefficient inequalities

THEOREM 3.1. *If*

$$f(z) = z^p + \sum_{n=1}^{\infty} a_{p+n} z^{p+n} \quad (3.1)$$

belongs to $\mathcal{C}_p(\lambda, \mu)$, then

$$|a_{p+1}| \leq \frac{8p^2(1-\mu)}{p+1} \left(\frac{\beta}{\pi \sin \beta} \right)^2 \quad (\beta = \arccos \lambda). \quad (3.2)$$

PROOF. It can be easily verified that

$$\begin{aligned} 1 + \frac{zf''(z)}{f'(z)} &= p + \left(1 + \frac{1}{p} \right) a_{p+1} z + \dots, \\ h(z) &= p + \frac{p(1-\mu)}{2 \sin^2 \beta} \left(\frac{8\beta}{\pi} + \frac{8\beta}{\pi} \left(\frac{2\beta}{\pi} - 1 \right) \right) z + \dots \\ &= p + 8p(1-\mu) \left(\frac{\beta}{\pi \sin \beta} \right)^2 z + \dots, \end{aligned} \quad (3.3)$$

where $h(z)$ is given by [\(2.2\)](#). Since

$$f(z) = z^p + a_{p+1} z^{p+1} + \dots \in \mathcal{C}_p(\lambda, \mu), \quad (3.4)$$

it follows from [\(3.3\)](#) and [Theorem 2.1](#) that

$$\begin{aligned} \frac{\pi^2}{8p(1-\mu)} \left(\frac{\sin \beta}{\beta} \right)^2 \left(1 + \frac{zf''(z)}{f'(z)} - p \right) \\ = \frac{p+1}{8p^2(1-\mu)} \left(\frac{\pi \sin \beta}{\beta} \right)^2 a_{p+1} z + \dots \\ < \frac{\pi^2}{8p(1-\mu)} \left(\frac{\sin \beta}{\beta} \right)^2 (h(z) - p). \end{aligned} \quad (3.5)$$

It is well known that if

$$f(z) = \sum_{n=1}^{\infty} a_n z^n \prec g(z) \quad (3.6)$$

for $g(z) \in \mathcal{H}$, then (cf. Duren [1])

$$|a_n| \leq 1 \quad (n = 1, 2, 3, \dots). \quad (3.7)$$

Noting that

$$\frac{\pi^2}{8p(1-\mu)} \left(\frac{\sin \beta}{\beta} \right)^2 (h(z) - p) \in \mathcal{H}, \quad (3.8)$$

we get (3.2). Also the bound in (3.2) is sharp for the function $f_0(z)$ given by (2.13). $\square$

REFERENCES

- [1] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, vol. 259, Springer-Verlag, New York, 1983.
- [2] A. W. Goodman, *On uniformly convex functions*, Ann. Polon. Math. **56** (1991), no. 1, 87–92.
- [3] W. C. Ma and D. Minda, *Uniformly convex functions*, Ann. Polon. Math. **57** (1992), no. 2, 165–175.
- [4] S. Owa, *On uniformly convex functions*, Math. Japon. **48** (1998), no. 3, 377–384.
- [5] F. Rønning, *Uniformly convex functions and a corresponding class of starlike functions*, Proc. Amer. Math. Soc. **118** (1993), no. 1, 189–196.
- [6] T. J. Suffridge, *Some remarks on convex maps of the unit disk*, Duke Math. J. **37** (1970), 775–777.

Dinggong Yang: Department of Mathematics, Suzhou University, Suzhou, Jiangsu 215006, China

Shigeyoshi Owa: Department of Mathematics, Kinki University, Higashi-Osaka, Osaka 577-8502, Japan

E-mail address: owa@math.kindai.ac.jp