FUZZY SUPER IRRESOLUTE FUNCTIONS

S. E. ABBAS

Received 26 December 2002

The concept of fuzzy super irresolute function was considered and studied by Šostak's (1985). A comparison between this type and other existing ones is established. Several characterizations, properties, and their effect on some fuzzy topological spaces are studied. Also, a new class of fuzzy topological spaces under the terminology fuzzy S^* -closed spaces is introduced and investigated.

2000 Mathematics Subject Classification: 54A40.

1. Introduction and preliminaries. Šostak [10], introduced the fundamental concept of a fuzzy topological structure, as an extension of both crisp topology and Chang fuzzy topology [1], in the sense that not only the objects are fuzzified, but also the axiomatics. In [11, 12], Šostak gave some rules and showed how such an extension can be realized. Chattopadhyay et al. [2, 3] have redefined the same concept. In [8], Ramadan gave a similar definition, namely "smooth topological space." It has been developed in many directions [4, 5, 6, 7, 13].

In the present note, some counterexamples and characterizations of fuzzy super irresolute functions are examined. It is seen that fuzzy super irresolute function implies each of fuzzy irresolute [9] and fuzzy continuity [10], but not conversely. Also, properties preserved by fuzzy super irresolute functions are examined. Finally, we define a fuzzy S^* -closed space in fuzzy topological spaces in Šostak sense and characterize such a space from different angles. Our aim is to compare the introduced type of fuzzy covering property with the existing ones.

Throughout this note, let X be a nonempty set, $I = [0, 1]$, and $I_0 = (0, 1]$. For $\alpha \in I$, $\underline{\alpha}(x) = \alpha$ for all $x \in X$. The following definition and results which will be needed.

DEFINITION 1.1 [10]. A function $\tau : I^X \rightarrow I$ is called a *fuzzy topology* on X if it satisfies the following conditions:

- (1) $\tau(\underline{0}) = \tau(\underline{1}) = 1$,
- (2) $\tau(\mu_1 \wedge \mu_2) \geq \tau(\mu_1) \wedge \tau(\mu_2)$ for any $\mu_1, \mu_2 \in I^X$,
- (3) $\tau(\bigvee_{i \in \Gamma} \mu_i) \geq \bigwedge_{i \in \Gamma} \tau(\mu_i)$ for any $\{\mu\}_{i \in \Gamma} \subset I^X$.

The pair (X, τ) is called a *fuzzy topological space* (FTS).

REMARK 1.2. Let (X, τ) be an FTS. Then, for each $\alpha \in I$, $\tau_\alpha = \{\mu \in I^X : \tau(\mu) \geq r\}$ is a Chang's fuzzy topology on X .

THEOREM 1.3 [3]. Let (X, τ) be an FTS. Then, for each $r \in I_0$ and $\lambda \in I^X$, an operator $C_\tau : I^X \times I_0 \rightarrow I^X$ is defined as follows:

$$C_\tau(\lambda, r) = \bigwedge \{\mu \in I^X : \lambda \leq \mu, \tau(\underline{1} - \mu) \geq r\}. \quad (1.1)$$

For $\lambda, \mu \in I^X$ and $r, s \in I_0$, the operator C_τ satisfies the following conditions:

- (1) $C_\tau(\underline{0}, r) = \underline{0}$, $\lambda \leq C_\tau(\lambda, r)$,
- (2) $C_\tau(\lambda, r) \vee C_\tau(\mu, r) = C_\tau(\lambda \vee \mu, r)$,
- (3) $C_\tau(\lambda, r) \leq C_\tau(\lambda, s)$ if $r \leq s$,
- (4) $C_\tau(C_\tau(\lambda, r), r) = C_\tau(\lambda, r)$.

THEOREM 1.4 [9]. Let (X, τ) be an FTS. Then, for each $r \in I_0$ and $\lambda \in I^X$, an operator $I_\tau : I^X \times I_0 \rightarrow I^X$ is defined as follows:

$$I_\tau(\lambda, r) = \bigvee \{\mu \in I^X : \lambda \geq \mu, \tau(\mu) \geq r\}. \quad (1.2)$$

For $\lambda, \mu \in I^X$ and $r, s \in I_0$, the operator I_τ satisfies the following conditions:

- (1) $I_\tau(\underline{1} - \lambda, r) = \underline{1} - C_\tau(\lambda, r)$,
- (2) $I_\tau(\underline{1}, r) = \underline{1}$, $\lambda \geq I_\tau(\lambda, r)$,
- (3) $I_\tau(\lambda, r) \wedge I_\tau(\mu, r) = I_\tau(\lambda \wedge \mu, r)$,
- (4) $I_\tau(\lambda, r) \geq I_\tau(\lambda, s)$ if $r \leq s$,
- (5) $I_\tau(I_\tau(\lambda, r), r) = I_\tau(\lambda, r)$.

DEFINITION 1.5 [9]. Let (X, τ) be an FTS. Then, for each $r \in I_0$ and $\lambda \in I^X$, the following statements hold:

- (1) λ is called r -fuzzy semi-open (r -FSO) if there exists $v \in I^X$ with $\tau(v) \geq r$ such that $v \leq \lambda \leq C_\tau(v, r)$; equivalently, $\lambda \leq C_\tau(I_\tau(\lambda, r), r)$;
- (2) λ is called r -fuzzy semiclosed (r -FSC) if there exists $v \in I^X$ with $\tau(\underline{1} - v) \geq r$ such that $I_\tau(v, r) \leq \lambda \leq v$; equivalently, $I_\tau(C_\tau(\lambda, r), r) \leq \lambda$;
- (3) λ is called r -fuzzy semiclopen (r -FSCO) if λ is r -FSO and r -FSC;
- (4) λ is called r -fuzzy regular open (r -FRO) if $\lambda = I_\tau(C_\tau(\lambda, r), r)$;
- (5) the r -fuzzy semi-interior of λ , denoted $SI_\tau(\lambda, r)$, is defined by $SI_\tau(\lambda, r) = \bigvee \{v \in I^X : v \leq \lambda, v \text{ is } r\text{-FSO}\}$;
- (6) the r -fuzzy semiclosure of λ , denoted $SC_\tau(\lambda, r)$, is defined by $SC_\tau(\lambda, r) = \bigwedge \{v \in I^X : v \geq \lambda, v \text{ is } r\text{-FSC}\}$.

THEOREM 1.6 [9]. Let (X, τ) be an FTS. For $\lambda \in I^X$ and $r \in I_0$, the following statements are valid:

- (1) λ is r -FSO if and only if $\lambda = SI_\tau(\lambda, r)$, and λ is r -FSC if and only if $\lambda = SC_\tau(\lambda, r)$;
- (2) $I_\tau(\lambda, r) \leq SI_\tau(\lambda, r) \leq \lambda \leq SC_\tau(\lambda, r) \leq C_\tau(\lambda, r)$;

- (3) $SC_\tau(SC_\tau(\lambda, r), r) = SC_\tau(\lambda, r)$;
- (4) $C_\tau(SC_\tau(\lambda, r), r) = SC_\tau(C_\tau(\lambda, r), r) = C_\tau(\lambda, r)$;
- (5) $SI_\tau(\underline{1} - \lambda, r) = \underline{1} - SC_\tau(\lambda, r)$.

LEMMA 1.7. For any fuzzy set λ in an FTS (X, τ) and $r \in I_\circ$, if $\tau(\lambda) \geq r$, then $I_\tau(C_\tau(\lambda, r), r) = SC_\tau(\lambda, r)$.

PROOF. Since $SC_\tau(\lambda, r)$ is r -FSC, $I_\tau(C_\tau(SC_\tau(\lambda, r), r), r) \leq SC_\tau(\lambda, r)$ and hence, by Theorem 1.6(4), $I_\tau(C_\tau(\lambda, r), r) \leq SC_\tau(\lambda, r)$. To prove the opposite inclusion, since $\tau(\lambda) \geq r$, $r \in I_\circ$, we have $\lambda \leq I_\tau(C_\tau(\lambda, r), r)$ so that $\underline{1} - \lambda \geq \underline{1} - I_\tau(C_\tau(\lambda, r), r) = C_\tau(I_\tau(\underline{1} - \lambda, r), r)$. But $C_\tau(I_\tau(\underline{1} - \lambda, r), r)$ is r -FSO. Hence $C_\tau(I_\tau(\underline{1} - \lambda, r), r) \leq SI_\tau(\underline{1} - \lambda, r)$ and so $SC_\tau(\lambda, r) \leq I_\tau(C_\tau(\lambda, r), r)$. $\square$

DEFINITION 1.8. Let (X, τ) and (Y, η) be FTSs and let $f : X \rightarrow Y$ be a function which is called

- (1) fuzzy continuous (FC) if and only if $\eta(\mu) \leq \tau(f^{-1}(\mu))$ for each $\mu \in I^Y$ [10],
- (2) fuzzy open if and only if $\tau(\lambda) \leq \eta(f(\lambda))$ for each $\lambda \in I^X$ [10],
- (3) fuzzy semicontinuous (FSC) if and only if $f^{-1}(\mu)$ is r -FSO set of X for each $\eta(\mu) \geq r$, $r \in I_\circ$ [9],
- (4) fuzzy irresolute (FI) if and only if $f^{-1}(\mu)$ is r -FSO set of X for each μ is r -FSO set of Y , $r \in I_\circ$ [9].

2. Fuzzy super irresolute functions

DEFINITION 2.1. Let (X, τ) and (Y, η) be FTSs and let $f : X \rightarrow Y$ be a function which is called

- (1) fuzzy super irresolute (F-super I) if and only if $\tau(f^{-1}(\mu)) \geq r$ for each μ is r -FSO set of Y , $r \in I_\circ$,
- (2) fuzzy completely continuous (FCC) if and only if $f^{-1}(\mu)$ is r -FRO set of X for each $\mu \in I^Y$ and $\eta(\mu) \geq r$, $r \in I_\circ$,
- (3) fuzzy completely irresolute (FCI) if and only if $f^{-1}(\mu)$ is r -FRO set of X for each r -FSO set $\mu \in I^Y$ and $r \in I_\circ$.

REMARK 2.2. One can show the connection between these types and other existing ones by the following diagram:

$$\begin{array}{ccccc}
 \text{FCI} & \longrightarrow & \text{F-super I} & \longrightarrow & \text{FI} \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{FCC} & \longrightarrow & \text{FC} & \longrightarrow & \text{FSC}
 \end{array} \tag{2.1}$$

The converse of the previous implications need not be true in general as shown in the following counterexample.

COUNTEREXAMPLE 2.3. Let μ_1 , μ_2 , and μ_3 be fuzzy subsets of $X = \{a, b, c\}$ defined as follows:

$$\begin{aligned}\mu_1(a) &= 0.9, & \mu_1(b) &= 0.0, & \mu_1(c) &= 0.1, \\ \mu_2(a) &= 0.9, & \mu_2(b) &= 0.7, & \mu_2(c) &= 0.2, \\ \mu_3(a) &= 0.9, & \mu_3(b) &= 0.3, & \mu_3(c) &= 0.2.\end{aligned}\tag{2.2}$$

Then $\tau, \eta : I^X \rightarrow I$, defined as

$$\tau(\lambda) = \begin{cases} 1, & \text{if } \lambda = \underline{0}, \underline{1}, \\ \frac{1}{2}, & \text{if } \lambda = \mu_1, \\ \frac{1}{3}, & \text{if } \lambda = \mu_2, \\ 0, & \text{otherwise,} \end{cases} \quad \eta(\lambda) = \begin{cases} 1, & \text{if } \lambda = \underline{0}, \underline{1}, \\ \frac{1}{3}, & \text{if } \lambda = \mu_1, \mu_2, \\ \frac{1}{2}, & \text{if } \lambda = \mu_3, \\ 0, & \text{otherwise,} \end{cases}\tag{2.3}$$

are fuzzy topologies on X . Then,

- (1) the identity function $\text{id}_X : (X, \tau) \rightarrow (X, \eta)$ is FI but not F-super I because μ_3 is $1/3$ -FSO in (X, η) and $\tau(f^{-1}(\mu_3)) = \tau(\mu_3) = 0$;
- (2) the identity function $\text{id}_X : (X, \tau) \rightarrow (X, \tau)$ is FC but not F-super I function.

DEFINITION 2.4. An FTS (X, τ) is said to be fuzzy extremally disconnected if and only if $\tau(C_\tau(\lambda, r)) \geq r$ for every $\tau(\lambda) \geq r$ for each $\lambda \in I^X$ and $r \in I_\circ$.

THEOREM 2.5. For a function $f : X \rightarrow Y$, the following statements are true:

- (1) if X is fuzzy extremally disconnected and f is FI, then f is F-super I;
- (2) if Y is fuzzy extremally disconnected and f is FCI (resp., FC), then f is F-super I;
- (3) if both X and Y are fuzzy extremally disconnected, then the concepts F-super I, FCI, FI, FCC, FSC, and FC are equivalent.

PROOF. The proof is obvious. □

THEOREM 2.6. Let (X, τ_1) and (Y, τ_2) be FTSs. Let $f : X \rightarrow Y$ be a function. The following statements are equivalent:

- (1) a map f is F-super I;
- (2) for each r -FSC $\mu \in I^Y$, $\tau(\underline{1} - f^{-1}(\mu)) \geq r$, $r \in I_\circ$;
- (3) for each $\lambda \in I^X$ and $r \in I_\circ$, $f(C_{\tau_1}(\lambda, r)) \leq \text{SC}_{\tau_2}(f(\lambda), r)$;
- (4) for each $\mu \in I^Y$ and $r \in I_\circ$, $C_{\tau_1}(f^{-1}(\mu), r) \leq f^{-1}(\text{SC}_{\tau_2}(\mu, r))$;
- (5) for each $\mu \in I^Y$ and $r \in I_\circ$, $f^{-1}(\text{SI}_{\tau_2}(\mu, r)) \leq I_{\tau_1}(f^{-1}(\mu), r)$.

PROOF. (1) $\Leftrightarrow$ (2). It is easily proved from Theorem 1.4 and from $f^{-1}(\underline{1} - \mu) = \underline{1} - f^{-1}(\mu)$.

(2) $\Rightarrow$ (3). Suppose there exist $\lambda \in I^X$ and $r \in I_\circ$ such that

$$f(C_{\tau_1}(\lambda, r)) \not\leq \text{SC}_{\tau_2}(f(\lambda), r).\tag{2.4}$$

There exist $y \in Y$ and $t \in I_0$ such that

$$f(C_{\tau_1}(\lambda, r))(y) > t > SC_{\tau_2}(f(\lambda), r)(y). \quad (2.5)$$

If $f^{-1}(\{y\}) = \emptyset$, it is a contradiction because $f(C_{\tau_1}(\lambda, r))(y) = 0$.

If $f^{-1}(\{y\}) \neq \emptyset$, there exists $x \in f^{-1}(\{y\})$ such that

$$f(C_{\tau_1}(\lambda, r))(y) \geq C_{\tau_1}(\lambda, r)(x) > t > SC_{\tau_2}(f(\lambda), r)(f(x)). \quad (2.6)$$

Since $SC_{\tau_2}(f(\lambda), r)(f(x)) < t$, there exists r -FSC $\mu \in I^Y$ with $f(\lambda) \leq \mu$ such that

$$SC_{\tau_2}(f(\lambda), r)(f(x)) \leq \mu(f(x)) < t. \quad (2.7)$$

Moreover, $f(\lambda) \leq \mu$ implies $\lambda \leq f^{-1}(\mu)$. From (2), $\tau(\underline{1} - f^{-1}(\mu)) \geq r$. Thus, $C_{\tau_1}(\lambda, r)(x) \leq f^{-1}(\mu)(x) = \mu(f(x)) < t$, which is a contradiction to (2.6).

(3) $\Rightarrow$ (4). For all $\mu \in I^Y$, $r \in I_0$, put $\lambda = f^{-1}(\mu)$. From (3), we have

$$f(C_{\tau_1}(f^{-1}(\mu), r)) \leq SC_{\tau_2}(f(f^{-1}(\mu)), r) \leq SC_{\tau_2}(\mu, r), \quad (2.8)$$

which implies that

$$C_{\tau_1}(f^{-1}(\mu), r) \leq f^{-1}(f(C_{\tau_1}(f^{-1}(\mu), r))) \leq f^{-1}(SC_{\tau_2}(\mu, r)). \quad (2.9)$$

(4) $\Rightarrow$ (5). It is easily proved from [Theorem 1.4\(1\)](#).

(5) $\Rightarrow$ (1). Let μ be r -FSO set of Y . From [Theorem 1.6\(1\)](#), $\mu = SI_{\tau_2}(\mu, r)$. By (5),

$$f^{-1}(\mu) \leq I_{\tau_1}(f^{-1}(\mu), r). \quad (2.10)$$

On the other hand, by [Theorem 1.4\(2\)](#),

$$f^{-1}(\mu) \geq I_{\tau_1}(f^{-1}(\mu), r). \quad (2.11)$$

Thus, $f^{-1}(\mu) = I_{\tau_1}(f^{-1}(\mu), r)$, that is, $\tau(f^{-1}(\mu)) \geq r$. $\square$

3. Properties preserved by F-super I functions

DEFINITION 3.1. Let (X, τ) be an FTS and $r \in I_0$. Then

- (1) X is called r -fuzzy compact (resp., r -fuzzy almost compact and r -fuzzy nearly compact) if and only if for each family $\{\lambda_i \in I^X : \tau(\lambda_i) \geq r, i \in \Gamma\}$ such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, there exists a finite index set $\Gamma_0 \subset \Gamma$ such that $\bigvee_{i \in \Gamma_0} \lambda_i = \underline{1}$ (resp., $\bigvee_{i \in \Gamma_0} C_{\tau}(\lambda_i, r) = \underline{1}$ and $\bigvee_{i \in \Gamma_0} I_{\tau}(C_{\tau}(\lambda_i, r), r) = \underline{1}$);
- (2) X is called r -fuzzy semicompact (resp., r -fuzzy S -closed) if and only if for each family $\{\lambda_i \in I^X : \lambda_i \leq C_{\tau}(I_{\tau}(\lambda_i, r), r), i \in \Gamma\}$ such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, there exists a finite index set $\Gamma_0 \subset \Gamma$ such that $\bigvee_{i \in \Gamma_0} \lambda_i = \underline{1}$ (resp., $\bigvee_{i \in \Gamma_0} C_{\tau}(\lambda_i, r) = \underline{1}$).

THEOREM 3.2. *Every surjective F-super I image of r -fuzzy compact space is r -fuzzy semicompact, $r \in I_0$.*

PROOF. Let (X, τ) be r -fuzzy compact, $r \in I_0$, and let $f : (X, \tau) \rightarrow (Y, \eta)$ be F-super I surjective function. If $\{\lambda_i \in I^Y : \lambda_i \leq C_\eta(I_\eta(\lambda_i, r), r), i \in \Gamma\}$ with $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, then $\bigvee_{i \in \Gamma} f^{-1}(\lambda_i) = \underline{1}$. Since f is F-super I, $\tau(f^{-1}(\lambda_i)) \geq r$. Since X is r -fuzzy compact, there exists a finite subset $\Gamma_0 \subset \Gamma$ with $\bigvee_{i \in \Gamma_0} f^{-1}(\lambda_i) = \underline{1}$. From the surjectivity of f , we deduce

$$\underline{1} = f(\underline{1}) = f\left(\bigvee_{i \in \Gamma_0} f^{-1}(\lambda_i)\right) = \bigvee_{i \in \Gamma_0} f f^{-1}(\lambda_i) = \bigvee_{i \in \Gamma_0} \lambda_i. \quad (3.1)$$

So, Y is r -fuzzy semicompact. □

COROLLARY 3.3. *Every surjective F-super I image of r -fuzzy compact space is r -fuzzy S -closed, $r \in I_0$.*

THEOREM 3.4. *Every surjective F-super I image of r -fuzzy almost compact space is r -fuzzy S -closed, $r \in I_0$.*

PROOF. The proof is similar to that of [Theorem 3.2](#). □

COROLLARY 3.5. *r -fuzzy semicompactness and r -fuzzy S -closedness are preserved under an F-super I surjection function, $r \in I_0$.*

PROOF. The Corollary is a direct consequence of [Theorems 3.2](#) and [3.4](#). □

THEOREM 3.6. *Let $f : X \rightarrow Y$ be FSC and F-super I surjective function. If X is r -fuzzy nearly compact, then Y is r -fuzzy S -closed, $r \in I_0$.*

PROOF. Let (X, τ) be r -fuzzy nearly compact, and let $r \in I_0$, $f : (X, \tau) \rightarrow (Y, \eta)$ be FSC and F-super I surjective function. If $\{\lambda_i \in I^Y : \lambda_i \leq C_\eta(I_\eta(\lambda_i, r), r), i \in \Gamma\}$ with $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, then $\bigvee_{i \in \Gamma} f^{-1}(\lambda_i) = \underline{1}$. Since f is F-super I, $\tau(f^{-1}(\lambda_i)) \geq r$. Since X is r -fuzzy nearly compact, there exists a finite subset $\Gamma_0 \subset \Gamma$ with $\bigvee_{i \in \Gamma_0} I_\tau(C_\tau(f^{-1}(\lambda_i), r), r) = \underline{1}$. From the surjectivity of f , we deduce

$$\begin{aligned} \underline{1} = f(\underline{1}) &= f\left(\bigvee_{i \in \Gamma_0} I_\tau(C_\tau(f^{-1}(\lambda_i), r), r)\right) \\ &= \bigvee_{i \in \Gamma_0} f(I_\tau(C_\tau(f^{-1}(\lambda_i), r), r)) \\ &\leq \bigvee_{i \in \Gamma_0} f(f^{-1}(C_\eta(\lambda_i, r))) \quad (\text{since } f \text{ is FSC [9]}). \end{aligned} \quad (3.2)$$

Thus $\bigvee_{i \in \Gamma_0} C_\eta(\lambda_i, r) = \underline{1}$. Hence Y is r -fuzzy S -closed. □

4. Fuzzy S^* -closed spaces: characterizations and comparisons

DEFINITION 4.1. Let (X, τ) be an FTS and $r \in I_0$. Then X is called r -fuzzy S^* -closed if and only if for each family $\{\lambda_i \in I^X : \lambda_i \leq C_\tau(I_\tau(\lambda_i, r), r), i \in \Gamma\}$ such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, there exists a finite index set $\Gamma_0 \subset \Gamma$ such that

$$\bigvee_{i \in \Gamma_0} SC_\tau(\lambda_i, r) = \underline{1}. \quad (4.1)$$

THEOREM 4.2. For an FTS (X, τ) , $r \in I_0$, the following statements are equivalent:

- (1) X is r -fuzzy S^* -closed;
- (2) for every family $\{\lambda_i \in I^X : \lambda_i \text{ is } r\text{-FSCO}, i \in \Gamma\}$ such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, there exists a finite index set $\Gamma_0 \subset \Gamma$ such that $\bigvee_{i \in \Gamma_0} \lambda_i = \underline{1}$;
- (3) every family of r -FSCO sets having the finite intersection property has nonnull intersection;
- (4) for every family $\{\lambda_i \in I^X : \lambda_i \text{ is } r\text{-FSC}, i \in \Gamma\}$ such that $\bigwedge_{i \in \Gamma} \lambda_i = \underline{1}$, there exists a finite index set $\Gamma_0 \subset \Gamma$ such that $\bigwedge_{i \in \Gamma_0} SI_\tau(\lambda_i, r) = \underline{1}$.

PROOF. (1) $\Rightarrow$ (2). The proof is obvious.

(2) $\Rightarrow$ (3). Let $\{\lambda_i\}_{i \in \Gamma}$ be a family of r -FSCO sets having the finite intersection property. If possible, let $\bigwedge_{i \in \Gamma} \lambda_i = \underline{0}$. Then $\bigvee_{i \in \Gamma} (\underline{1} - \lambda_i) = \underline{1}$, where each $(\underline{1} - \lambda_i)$ is r -FSCO. By (2), there exists a finite subset Γ_0 of Γ such that $\bigvee_{i \in \Gamma_0} \underline{1} - \lambda_i = \underline{1}$, that is, $\bigwedge_{i \in \Gamma_0} \lambda_i = \underline{0}$, which is a contradiction.

(3) $\Rightarrow$ (1). Suppose that $\{\lambda_i : i \in \Gamma\}$ is a family of r -FSO sets of X with $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, and it has no finite subfamily $\{\lambda_{i_1}, \dots, \lambda_{i_n}\}$ such that $\bigvee_{j=1}^n SC_\tau(\lambda_{i_j}, r) = \underline{1}$. Then $\bigwedge_{i=1}^n (\underline{1} - SC_\tau(\lambda_{i_j}, r)) \neq \underline{0}$. Thus, $\{\underline{1} - SC_\tau(\lambda_i, r) : i \in \Gamma\}$ is a family of r -FSCO sets having the finite intersection property. By (3), $\bigwedge_{i \in \Gamma} (\underline{1} - SC_\tau(\lambda_i, r)) \neq \underline{0}$, and hence, $\bigvee_{i \in \Gamma} \lambda_i \neq \underline{1}$, which is a contradiction.

(1) $\Rightarrow$ (4). If $\{\lambda_i : i \in \Gamma\}$ is a family of nonnull r -FSC sets in X , $r \in I_0$ with $\bigwedge_{i \in \Gamma} \lambda_i = \underline{0}$, then $\{\underline{1} - \lambda_i : i \in \Gamma\}$ is r -FSO sets in X with $\bigvee_{i \in \Gamma} \underline{1} - \lambda_i = \underline{1}$. By (1), there is a finite subset $\Gamma_0 \subset \Gamma$ such that

$$\underline{1} = \bigvee_{i \in \Gamma_0} SC_\tau(\underline{1} - \lambda_i, r) = \underline{1} - \bigwedge_{i \in \Gamma_0} SI_\tau(\lambda_i, r), \quad (4.2)$$

that is, $\bigwedge_{i \in \Gamma_0} SI_\tau(\lambda_i, r) = \underline{0}$.

(4) $\Rightarrow$ (1). For any $\{\lambda_i \in I^X : \lambda_i \text{ is } r\text{-FSO}, i \in \Gamma\}$ such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, $\{\underline{1} - \lambda_i, i \in \Gamma\}$ is a family of r -FSC sets such that $\bigwedge_{i \in \Gamma} \underline{1} - \lambda_i = \underline{0}$. We can assume, without loss of generality, that each $\underline{1} - \lambda_i \neq \underline{0}$. By (4), there is a finite subset $\Gamma_0 \subset \Gamma$ such that $\bigwedge_{i \in \Gamma_0} SI_\tau(\underline{1} - \lambda_i, r) = \underline{0}$, that is, $\bigvee_{i \in \Gamma_0} SC_\tau(\lambda_i, r) = \underline{1}$, which proves the r -fuzzy S^* -closedness of X . $\square$

THEOREM 4.3. Let (X, τ) be an FTS and $r \in I_0$. If X is r -fuzzy semicompact, then X is r -fuzzy S^* -closed as well.

PROOF. Since for every $\lambda \in I^X$ and $r \in I_0$ we have $\lambda \leq SC_\tau(\lambda, r)$, this immediately follows from the definitions. $\square$

THEOREM 4.4. *Let (X, τ) be an FTS and $r \in I_0$. If X is r -fuzzy S^* -closed, then X is r -fuzzy S -closed as well.*

PROOF. Since for every $\lambda \in I^X$ and $r \in I_0$ we have $SC_\tau(\lambda, r) \leq C_\tau(\lambda, r)$, this immediately follows from the definitions. $\square$

That the converse is false is evident from the following counterexample.

COUNTEREXAMPLE 4.5. Let $\mathbb{N}$ denote the set of natural numbers with the fuzzy topology $\tau : I^{\mathbb{N}} \rightarrow I$ defined as

$$\tau(\lambda) = \begin{cases} 1, & \text{if } \lambda = \underline{0}, \underline{1}, \\ \frac{1}{3}, & \text{if } \lambda = \mu, \nu, \\ \frac{1}{2}, & \text{if } \lambda = \mu \vee \nu, \\ 0, & \text{otherwise,} \end{cases} \quad (4.3)$$

where $\mu(1) = 1$, $\mu(i) = 0$ (for $i = 2, 3, 4, \dots$), and $\nu(2) = 1$, $\nu(j) = 0$ (for $j = 1, 3, 4, \dots$). Let ρ_i^1 and ρ_i^2 (for $i = 3, 4, 5, \dots$) be the fuzzy sets in $I^{\mathbb{N}}$ given by

$$\begin{aligned} \rho_i^1(x) &= \begin{cases} 1, & \text{for } x = 1 \text{ and } i, \\ 0, & \text{otherwise,} \end{cases} \\ \rho_i^2(x) &= \begin{cases} 1, & \text{for } x = 2 \text{ and } i, \\ 0, & \text{otherwise.} \end{cases} \end{aligned} \quad (4.4)$$

Then $\mathcal{U} = \{\rho_i^1, \rho_i^2 : i = 3, 4, 5, \dots\}$ are $1/3$ -FSO sets with $\bigvee_{\rho \in \mathcal{U}} \rho = \underline{1}$ having no finite subcover. Hence $(\mathbb{N}, \tau)$ is not $1/3$ -fuzzy S^* -closed, but it is easily seen that $(\mathbb{N}, \tau)$ is $1/3$ -fuzzy S -closed.

THEOREM 4.6. *For any fuzzy extremally disconnected FTS (X, τ) and $r \in I_0$, X is r -fuzzy S^* -closed if and only if X is r -fuzzy S -closed.*

PROOF

NECESSITY. It follows from the proof of [Theorem 4.4](#).

SUFFICIENCY. We are going to prove that if (X, τ) is any fuzzy extremally disconnected FTS, then $C_\tau(\lambda, r) = SC_\tau(\lambda, r)$ for every r -FSO set λ in (X, τ) and $r \in I_0$. Then our result follows from [Definitions 3.1\(2\) and 4.1](#).

We always have $SC_\tau(\lambda, r) \leq C_\tau(\lambda, r)$ for every $\lambda \in I^X$ and $r \in I_0$. So, we have to prove that with our hypothesis we have $C_\tau(\lambda, r) \leq SC_\tau(\lambda, r)$ for every $\lambda \in I^X$ and $r \in I_0$.

If λ is r -FSO in (X, τ) , then there exists $\nu \in I^X$ with $\tau(\nu) \geq r$ such that $\nu \leq \lambda \leq C_\tau(\nu, r)$. So, $C_\tau(\lambda, r) = C_\tau(\nu, r)$, where $\tau(\nu) \geq r$. Because (X, τ) is

fuzzy extremally disconnected, we have that

$$C_\tau(\lambda, r) = C_\tau(v, r) = I_\tau(C_\tau(v, r), r) = I_\tau(C_\tau(\lambda, r), r). \quad (4.5)$$

By Lemma 1.7, we have $C_\tau(\lambda, r) = I_\tau(C_\tau(\lambda, r), r) \leq SC_\tau(\lambda, r)$. $\square$

REMARK 4.7. From Theorems 4.3 and 4.4, we have that r -fuzzy semicomactness implies r -fuzzy S -closedness, $r \in I_0$.

REMARK 4.8. Obviously, for $r \in I_0$, r -fuzzy S -closed space is r -fuzzy almost compact. Hence r -fuzzy compact space need not be r -fuzzy S^* -closed. That an r -fuzzy S^* -closed space is not necessarily r -fuzzy compact is shown by the following counterexample.

COUNTEREXAMPLE 4.9. Let X be any nonempty set and let $\tau : I^X \rightarrow I$ be defined as

$$\tau(\lambda) = \begin{cases} 1, & \text{if } \lambda = \underline{0}, \underline{1}, \\ \frac{1}{2}, & \text{if } \lambda = \underline{\alpha}, \text{ for } \frac{1}{2} < \alpha < 1, \\ 0, & \text{otherwise.} \end{cases} \quad (4.6)$$

Then (X, τ) is an FTS which is not $1/2$ -fuzzy compact. Now for any $\underline{\alpha} \in I^X$ with $\tau(\underline{\alpha}) \geq 1/2$, $C_\tau(\underline{\alpha}, 1/2) = \underline{1}$ and hence $I_\tau(C_\tau(\underline{\alpha}, 1/2), 1/2) = \underline{1}$, for all $\alpha \in (1/2, 1]$. Since, by Lemma 1.7, $SC_\tau(\underline{\alpha}, 1/2) = I_\tau(C_\tau(\underline{\alpha}, 1/2), 1/2) = \underline{1}$, we have for any r -FSO set λ , $SC_\tau(\lambda, 1/2) = \underline{1}$. Hence X is r -fuzzy S^* -closed.

However, we have the following theorem.

THEOREM 4.10. For $r \in I_0$, every r -fuzzy S^* -closed space is r -fuzzy nearly compact, $r \in I_0$.

PROOF. If X is not r -fuzzy nearly compact, then there exists $\{\lambda_i \in I^X, i \in \Gamma\}$ with $\tau(\lambda_i) \geq r$ and $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$ such that for any finite subset $\Gamma_0 \subset \Gamma$,

$$\bigvee_{i \in \Gamma_0} I_\tau(C_\tau(\lambda_i, r), r) \neq \underline{1}, \quad (4.7)$$

that is,

$$\bigvee_{i \in \Gamma_0} SC_\tau(\lambda_i, r) \neq \underline{1} \quad (4.8)$$

(by Lemma 1.7). Thus, X is not r -fuzzy S^* -closed. $\square$

In order to investigate for the condition under which r -fuzzy S^* -closed space is r -fuzzy compact, we set the following definition.

DEFINITION 4.11. An FTS (X, τ) is called r -fuzzy S -regular if and only if for each r -FSO set $\mu \in I^X$, $r \in I_\circ$,

$$\mu = \bigvee \{ \rho \in I^X \mid \rho \text{ is } r\text{-FSO, } SC_\tau(\rho, r) \leq \mu \}. \quad (4.9)$$

An FTS (X, τ) is called fuzzy S -regular if and only if it is r -fuzzy S -regular for each $r \in I_\circ$.

THEOREM 4.12. *If an FTS (X, τ) is r -fuzzy S -regular and r -fuzzy S^* -closed, $r \in I_\circ$, then it is r -fuzzy compact.*

PROOF. Let $\{\lambda_i \in I^X \mid \tau(\lambda_i) \geq r, i \in \Gamma\}$ be a family such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$. Since (X, τ) is r -fuzzy S -regular, for each $\tau(\lambda_i) \geq r$, λ_i is r -FSO,

$$\lambda_i = \bigvee_{i_k \in K_i} \{ \lambda_{i_k} \mid \lambda_{i_k} \text{ is } r\text{-FSO, } SC_\tau(\lambda_{i_k}, r) \leq \lambda_i \}. \quad (4.10)$$

Hence $\bigvee_{i \in \Gamma} (\bigvee_{i_k \in K_i} \lambda_{i_k}) = \underline{1}$. Since (X, τ) is r -fuzzy S^* -closed, there exists a finite index $J \times K_J$ such that

$$\underline{1} = \bigvee_{j \in J} \left(\bigvee_{j_k \in K_J} SC_\tau(\lambda_{j_k}, r) \right). \quad (4.11)$$

For each $j \in J$, since

$$\bigvee_{j_k \in K_J} SC_\tau(\lambda_{j_k}, r) \leq \lambda_j, \quad (4.12)$$

we have $\bigvee_{j \in J} \lambda_j = \underline{1}$. Hence (X, τ) is r -fuzzy compact. $\square$

It is evident that every FI function is FSC. That the converse is not always true is shown in [9]. Again, it is proved in [9] that $f : X \rightarrow Y$ is FI if and only if $f^{-1}(\mu)$ is r -FSC for every r -FSC set μ in Y and $r \in I_\circ$. Now we have the following theorem.

THEOREM 4.13. *The FI image of r -fuzzy S^* -closed space is r -fuzzy S^* -closed, $r \in I_\circ$.*

THEOREM 4.14. *If $f : (X, \tau) \rightarrow (Y, \eta)$ is FI surjective and X is r -fuzzy S^* -closed, then Y is r -fuzzy S -closed, $r \in I_\circ$.*

PROOF. If $\{\lambda_i \in I^Y : \lambda_i \text{ is } r\text{-FSO, } i \in \Gamma\}$ is a family such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$, then $\bigvee_{i \in \Gamma} f^{-1}(\lambda_i) = \underline{1}$. Since f is FI, then, for each $i \in \Gamma$, $f^{-1}(\lambda_i)$ is r -FSO set of X . By r -fuzzy S^* -closedness of X , there is a finite subset $\Gamma_\circ \subset \Gamma$ such that

$\bigvee_{i \in \Gamma_0} \text{SC}_\tau(f^{-1}(\lambda_i, r)) = \underline{1}$. Now,

$$\begin{aligned} \underline{1} &= f(\underline{1}) = f\left(\bigvee_{i \in \Gamma_0} \text{SC}_\tau(f^{-1}(\lambda_i, r))\right) \\ &\leq f\left(\bigvee_{i \in \Gamma_0} C_\tau(f^{-1}(\lambda_i, r))\right) \\ &\leq \bigvee_{i \in \Gamma_0} C_\eta(\lambda_i, r), \end{aligned} \quad (4.13)$$

which implies that Y is r -fuzzy S -closed. $\square$

THEOREM 4.15. *If $f : (X, \tau) \rightarrow (Y, \eta)$ is CI surjective and X is r -fuzzy nearly compact, then Y is r -fuzzy semicompact, $r \in I_0$.*

PROOF. The proof is similar to that of [Theorem 4.14](#). $\square$

DEFINITION 4.16. Let (X, τ) and (Y, η) be FTSS. A function $f : (X, \tau) \rightarrow (Y, \eta)$ is called semiweakly continuous if and only if

$$f^{-1}(\lambda) \leq \text{SI}_\tau(f^{-1}(\text{SC}_\eta(\lambda, r)), r), \quad (4.14)$$

for each r -FSO set λ in (Y, η) , $r \in I_0$.

THEOREM 4.17. *Let (X, τ) and (Y, η) be FTSS and let $f : (X, \tau) \rightarrow (Y, \eta)$ be a semiweakly continuous function. If X is r -fuzzy semicompact, then Y is r -fuzzy S^* -closed, $r \in I_0$.*

PROOF. If $\{\lambda_i \in I^Y : \lambda_i \text{ is } r\text{-FSO}, i \in \Gamma\}$ is a family such that $\bigvee_{i \in \Gamma} \lambda_i = \underline{1}$. From the semiweak continuity of f , we have $f^{-1}(\lambda_i) \leq \text{SI}_\tau(f^{-1}(\text{SC}_\eta(\lambda_i, r)), r)$. So, $\text{SI}_\tau(f^{-1}(\text{SC}_\eta(\lambda_i, r)), r)$ is a family of r -FSO sets in (X, τ) with

$$\bigvee_{i \in \Gamma} \text{SI}_\tau(f^{-1}(\text{SC}_\eta(\lambda_i, r)), r) = \underline{1}. \quad (4.15)$$

By the semicompactness of X , there exists a finite subset $\Gamma_0 \subset \Gamma$ such that $\bigvee_{i \in \Gamma_0} \text{SI}_\tau(f^{-1}(\text{SC}_\eta(\lambda_i, r)), r) = \underline{1}$. So,

$$\begin{aligned} \underline{1} &= f(\underline{1}) = f\left(\bigvee_{i \in \Gamma_0} \text{SI}_\tau(f^{-1}(\text{SC}_\eta(\lambda_i, r)), r)\right) \\ &\leq \bigvee_{i \in \Gamma_0} f f^{-1}(\text{SC}_\eta(\lambda_i, r)) \\ &\leq \bigvee_{i \in \Gamma_0} \text{SC}_\eta(\lambda_i, r). \end{aligned} \quad (4.16)$$

Hence, $\bigvee_{i \in \Gamma_0} \text{SC}_\eta(\lambda_i, r) = \underline{1}$ and Y is r -fuzzy S^* -closed. $\square$

ACKNOWLEDGMENT. The author is very grateful to the referees.

REFERENCES

- [1] C. L. Chang, *Fuzzy topological spaces*, J. Math. Anal. Appl. **24** (1968), 182–190.
- [2] K. C. Chattopadhyay, R. N. Hazra, and S. K. Samanta, *Gradation of openness: fuzzy topology*, Fuzzy Sets and Systems **49** (1992), no. 2, 237–242.
- [3] K. C. Chattopadhyay and S. K. Samanta, *Fuzzy topology: fuzzy closure operator, fuzzy compactness and fuzzy connectedness*, Fuzzy Sets and Systems **54** (1993), no. 2, 207–212.
- [4] U. Höhle, *Upper semicontinuous fuzzy sets and applications*, J. Math. Anal. Appl. **78** (1980), no. 2, 659–673.
- [5] U. Höhle and A. P. Šostak, *A general theory of fuzzy topological spaces*, Fuzzy Sets and Systems **73** (1995), no. 1, 131–149.
- [6] ———, *Axiomatic foundations of fixed-basis fuzzy topology*, Mathematics of Fuzzy Sets, Handb. Fuzzy Sets Ser., vol. 3, Kluwer Academic Publishers, Massachusetts, 1999, pp. 123–272.
- [7] T. Kubiak and A. P. Šostak, *Lower set-valued fuzzy topologies*, Quaestiones Math. **20** (1997), no. 3, 423–429.
- [8] A. A. Ramadan, *Smooth topological spaces*, Fuzzy Sets and Systems **48** (1992), no. 3, 371–375.
- [9] A. A. Ramadan, S. E. Abbas, and Y. C. Kim, *Fuzzy irresolute mappings in smooth fuzzy topological spaces*, J. Fuzzy Math. **9** (2001), no. 4, 865–877.
- [10] A. P. Šostak, *On a fuzzy topological structure*, Rend. Circ. Mat. Palermo (2) Suppl. (1985), no. 11, 89–103.
- [11] ———, *On the neighborhood structure of fuzzy topological spaces*, Zb. Rad. (1990), no. 4, 7–14.
- [12] ———, *Basic structures of fuzzy topology*, J. Math. Sci. **78** (1996), no. 6, 662–701.
- [13] D. Zhang, *On the relationship between several basic categories in fuzzy topology*, Quaestiones Math. **25** (2002), no. 3, 289–301.

S. E. Abbas: Department of Mathematics, Faculty of Science, South Valley University, Sohag 82524, Egypt

E-mail address: sabbas73@yahoo.com