q -RIEMANN ZETA FUNCTION

TAEKYUN KIM

Received 19 July 2003

We consider the modified q -analogue of Riemann zeta function which is defined by $\zeta_q(s) = \sum_{n=1}^{\infty} (q^{n(s-1)} / [n]^s)$, $0 < q < 1$, $s \in \mathbb{C}$. In this paper, we give q -Bernoulli numbers which can be viewed as interpolation of the above q -analogue of Riemann zeta function at negative integers in the same way that Riemann zeta function interpolates Bernoulli numbers at negative integers. Also, we will treat some identities of q -Bernoulli numbers using non-Archimedean q -integration.

2000 Mathematics Subject Classification: 11S80, 11B68.

1. Introduction. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$, and $\mathbb{C}_p$ will respectively denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field, and the completion of algebraic closure of $\mathbb{Q}_p$.

The p -adic absolute value in $\mathbb{C}_p$ is normalized so that $|p|_p = 1/p$. When one talks of q -extension, q is considered in many ways such as an indeterminate, a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, we normally assume $|q| < 1$. If $q \in \mathbb{C}_p$, then we normally assume $|q-1|_p < p^{-1/(p-1)}$ so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$. We use the notation

$$[x] = [x : q] = \frac{1-q^x}{1-q} = 1 + q + q^2 + \cdots + q^{x-1}. \quad (1.1)$$

Note that $\lim_{q \rightarrow 1} [x] = x$ for $x \in \mathbb{Z}_p$ in the p -adic case.

Let $UD(\mathbb{Z}_p)$ be denoted by the set of uniformly differentiable functions on $\mathbb{Z}_p$.

For $f \in UD(\mathbb{Z}_p)$, we start with the expression

$$\frac{1}{[p^N]} \sum_{0 \leq j < p^N} q^j f(j) = \sum_{0 \leq j < p^N} f(j) \mu_q(j + p^N \mathbb{Z}_p) \quad (1.2)$$

representing the analogue of Riemann's sums for f (cf. [4]).

The integral of f on $\mathbb{Z}_p$ will be defined as the limit ($N \rightarrow \infty$) of these sums, which exists. The p -adic q -integral of a function $f \in UD(\mathbb{Z}_p)$ is defined by (see [4])

$$\int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]} \sum_{0 \leq j < p^N} f(j) q^j. \quad (1.3)$$

For d that is a fixed positive integer with $(p, d) = 1$, let

$$\begin{aligned} X &= X_d = \varprojlim_N \frac{\mathbb{Z}}{dp^N \mathbb{Z}}, \quad X_1 = \mathbb{Z}_p, \\ X^* &= \bigcup_{\substack{0 < a < dp \\ (a, p) = 1}} a + dp\mathbb{Z}_p, \\ a + dp^N \mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\}, \end{aligned} \quad (1.4)$$

where $a \in \mathbb{Z}$ lies in $0 \leq a < dp^N$.

Let $\mathbb{N}$ be the set of positive integers. For $m, k \in \mathbb{N}$, the q -Bernoulli polynomials, $\beta_m^{(-m, k)}(x, q)$, of higher order for the variable x in $\mathbb{C}_p$ are defined using p -adic q -integral by (cf. [4])

$$\begin{aligned} \beta_m^{(-m, k)}(x, q) &= \underbrace{\int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k \text{ times}} [x + x_1 + x_2 + \cdots + x_k]^m \\ &\quad \cdot q^{-x_1(m+1) - x_2(m+2) - \cdots - x_k(m+k)} d\mu_q(x_1) d\mu_q(x_2) \cdots d\mu_q(x_k). \end{aligned} \quad (1.5)$$

Now, we define the q -Bernoulli numbers of higher order as follows (cf. [2, 4, 7]):

$$\beta_m^{(-m, k)} (= \beta_m^{(-m, k)}(q)) = \beta_m^{(-m, k)}(0, q). \quad (1.6)$$

By (1.5), it is known that (cf. [4])

$$\begin{aligned} \beta_m^{(-m, k)} &= \lim_{N \rightarrow \infty} \frac{1}{[p^N]^k} \sum_{x_1=0}^{p^N-1} \cdots \sum_{x_k=0}^{p^N-1} [x_1 + \cdots + x_k]^m q^{-x_1 m - x_2(m+1) - \cdots - x_k(m+k-1)} \\ &= \frac{1}{(1-q)^m} \sum_{i=0}^m \binom{m}{i} (-1)^i \frac{(i-m)(i-m-1) \cdots (i-m-k+1)}{[i-m][i-m-1] \cdots [i-m-k+1]}, \end{aligned} \quad (1.7)$$

where $\binom{m}{i}$ are the binomial coefficients.

Note that $\lim_{q \rightarrow 1} \beta_m^{(-m, k)} = B_m^{(k)}$, where $B_m^{(k)}$ are ordinary Bernoulli numbers of order k (cf. [2, 3, 5, 7, 9]). By (1.5) and (1.7), it is easy to see that

$$\begin{aligned} \beta_m^{(-m, 1)}(x, q) &= \sum_{i=0}^m \binom{m}{i} q^{xi} \beta_i^{(-m, 1)} [x]^{m-i} \\ &= \frac{1}{(1-q)^m} \sum_{j=0}^m q^{jx} \binom{m}{j} (-1)^j \frac{j-m}{[j-m]}. \end{aligned} \quad (1.8)$$

We modify the q -analogue of Riemann zeta function which is defined in [1] as follows: for $q \in \mathbb{C}$ with $0 < q < 1$, $s \in \mathbb{C}$, define

$$\zeta_q(s) = \sum_{n=1}^{\infty} \frac{q^{(s-1)n}}{[n]^s}. \quad (1.9)$$

The numerator ensures the analytic continuation for $\Re(s) > 1$. In (1.9), we can consider the following problem.

“Are there q -Bernoulli numbers which can be viewed as interpolation of $\zeta_q(s)$ at negative integers in the same way that Riemann zeta function interpolates Bernoulli numbers at negative integers?”

In this paper, we give the value $\zeta_q(-m)$ for $m \in \mathbb{N}$, which is the answer of the above problem, and construct a new complex q -analogue of Hurwitz’s zeta function and q - L -series. Also, we will treat some interesting identities of q -Bernoulli numbers.

2. Some identities of q -Bernoulli numbers $\beta_m^{(-m,1)}$. In this section, we assume $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-1/(p-1)}$. By (1.5), we have

$$\begin{aligned}\beta_n^{(-n,1)}(x, q) &= \int_X q^{-(n+1)t} [x+t]^n d\mu_q(t) \\ &= [d]^{n-1} \sum_{i=0}^{d-1} q^{-ni} \int_{\mathbb{Z}_p} q^{-(n+1)dx} \left[\frac{x+i}{d} : q^d \right]^n d\mu_{q^d}(x).\end{aligned}\quad (2.1)$$

Thus, we have

$$\beta_n^{(-n,1)}(x, q) = [d]^{n-1} \sum_{i=0}^{d-1} q^{-ni} \beta_n^{(-n,1)}\left(\frac{x+i}{d}, q^d\right), \quad (2.2)$$

where d, n are positive integers.

If we take $x = 0$, then we have

$$[n]\beta_m^{(-m,1)} - n[n]^m \beta_m^{(-m,1)}(q^n) = \sum_{k=0}^{m-1} \binom{m}{k} [n]^k \beta_k^{(-m,1)}(q^n) \sum_{j=1}^{n-1} q^{-(m-j)k} [j]^{m-k}. \quad (2.3)$$

It is easy to see that $\lim_{q \rightarrow 1} \beta_m^{(-m,1)} = B_m$, where B_m are ordinary Bernoulli numbers (cf. [7]).

REMARK 2.1. By (2.3), note that

$$n(1 - n^m)B_m = \sum_{k=0}^{m-1} \binom{m}{k} n^k B_k \sum_{j=1}^{n-1} j^{m-k}. \quad (2.4)$$

Let $F_q(t)$ be the generating function of $\beta_n^{(-n,1)}$ as follows:

$$F_q(t) = \sum_{k=0}^{\infty} \beta_k^{(-k,1)} \frac{t^k}{k!}. \quad (2.5)$$

By (1.7) and (2.5), we easily see that

$$F_q(t) = - \sum_{m=0}^{\infty} \left(m \sum_{n=0}^{\infty} q^{-mn} [n]^{m-1} \right) \frac{t^m}{m!}. \quad (2.6)$$

Through differentiating both sides with respect to t in (2.5) and (2.6), and comparing coefficients, we obtain the following proposition.

PROPOSITION 2.2. *For $m > 0$, there exists*

$$-\frac{\beta_m^{(-m,1)}}{m} = \sum_{n=1}^{\infty} q^{-nm} [n]^{m-1}. \quad (2.7)$$

Moreover, $\beta_0^{(0,1)} = (q-1)/\log q$.

REMARK 2.3. Note that Proposition 2.2 is a q -analogue of $\zeta(1-2m)$ for any positive integer m .

Let χ be a primitive Dirichlet character with conductor $f \in \mathbb{N}$.

For $m \in \mathbb{N}$, we define

$$\beta_{m,\chi}^{(-m,1)} = \int_{\mathbb{X}} q^{-(m+1)x} \chi(x) [x]^m d\mu_q(x), \quad \text{for } m \geq 0. \quad (2.8)$$

Note that

$$\beta_{m,\chi}^{(-m,1)} = [d]^{m-1} \sum_{i=0}^{d-1} \chi(i) q^{-mi} \beta_m^{(-m,1)} \left(\frac{i}{d}, q^d \right). \quad (2.9)$$

3. q -analogs of zeta functions. In this section, we assume $q \in \mathbb{C}$ with $|q| < 1$. In [1], the q -analogue of Riemann zeta function was defined by (cf. [1])

$$\zeta_q^*(s) = \sum_{n=1}^{\infty} \frac{q^{ns}}{[n]^s}, \quad \Re(s) > 0. \quad (3.1)$$

Now, we modify the above q -analogue of Riemann zeta function as follows: for $q \in \mathbb{C}$ with $0 < |q| < 1$, $s \in \mathbb{C}$, define

$$\zeta_q(s) = \sum_{n=1}^{\infty} \frac{q^{(s-1)n}}{[n]^s}. \quad (3.2)$$

By (2.5), (2.6), and (2.7), we obtain the following proposition.

PROPOSITION 3.1. *For $m \in \mathbb{N}$, there exists*

- (i) $\zeta_q(1-m) = -\beta_m^{(-m,1)}/m$, for $m \geq 1$;
- (ii) $\zeta_q(s)$ having simple pole at $s = 1$ with residue $(q-1)/\log q$.

By (1.7) and (1.8), we see that

$$\beta_n^{(-n,1)}(x, q) = -n \sum_{k=0}^{\infty} ([k]q^x + [x])^{n-1} q^{-n(k+x)}, \quad \text{where } 0 \leq x < 1. \quad (3.3)$$

Hence, we can define q -analogue of Hurwitz ζ -function as follows: for $s \in \mathbb{C}$, define

$$\zeta_q(s, x) = \sum_{n=0}^{\infty} \frac{q^{(s-1)(n+x)}}{([n]q^x + [x])^s}. \quad (3.4)$$

Note that $\zeta_q(s, x)$ has an analytic continuation in $\mathbb{C}$ with only one simple pole at $s = 1$.

By (3.3) and (3.4), we have the following theorem.

THEOREM 3.2. *For any positive integer k , there exists*

$$\zeta_q(1-k, x) = -\frac{\beta_k^{(-k,1)}(x, q)}{k}. \quad (3.5)$$

Let χ be Dirichlet character with conductor $d \in \mathbb{N}$. By (2.9), the generalized q -Bernoulli numbers with χ can be defined by

$$\beta_{m, \chi}^{(-m,1)} = [d]^{m-1} \sum_{i=0}^{d-1} \chi(i) q^{-mi} \beta_m^{(-m,1)} \left(\frac{i}{d}, q^d \right). \quad (3.6)$$

For $s \in \mathbb{C}$, we define

$$L_q(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n) q^{(s-1)n}}{[n]^s}. \quad (3.7)$$

It is easy to see that

$$L_q(\chi, s) = [d]^{-s} \sum_{a=1}^d \chi(a) q^{(s-1)a} \zeta_{q^d} \left(s, \frac{a}{d} \right). \quad (3.8)$$

By (3.6), (3.7), and (3.8), we obtain the following theorem.

THEOREM 3.3. *Let k be a positive integer. Then there exists*

$$L_q(1-k, \chi) = -\frac{\beta_{k, \chi}^{(-k,1)}}{k}. \quad (3.9)$$

Let a and F be integers with $0 < a < F$. For $s \in \mathbb{C}$, we consider the functions $H_q(s, a, F)$ as follows:

$$H_q(s, a, F) = \sum_{m \equiv a(F), m > 0} \frac{q^{m(s-1)}}{[m]^s} = [F]^{-s} \zeta_{q^F} \left(s, \frac{a}{F} \right). \quad (3.10)$$

Then we have

$$H_q(1-n, a, F) = -\frac{[F]^{n-1}}{n} \beta_n^{(-n,1)} \left(\frac{a}{F}, q^F \right), \quad (3.11)$$

where n is any positive integer.

Therefore, we obtain the following theorem.

THEOREM 3.4. *Let a and F be integers with $0 < a < F$. For $s \in \mathbb{C}$, there exists*

- (i) $H_q(1-n, a, F) = -([F]^{n-1}/n)\beta_n^{(-n,1)}(a/F, q^F)$;
- (ii) $H_q(s, a, F)$ having a simple pole at $s = 1$ with residue $(1/[F]F)((q^F - 1)/\log q)$.

In a recent paper, the q -analogue of Riemann zeta function was studied by Cherednik (see [1]). In [1], we can consider the q -Bernoulli numbers which can be viewed as an interpolation of the q -analogue of Riemann zeta function at negative integers. In this paper, we have shown that the q -analogue of zeta function interpolates q -Bernoulli numbers at negative integers in the same way that Riemann zeta function interpolates Bernoulli numbers at negative integers (cf. [2, 5, 7]).

REMARK 3.5. Let $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-1/(p-1)}$. Then the p -adic q -gamma function was defined as (see [8])

$$\Gamma_{p,q}(n) = (-1)^n \prod_{1 \leq j < n, (j,p)=1} [j]. \quad (3.12)$$

For all $x \in \mathbb{Z}_p$, we have

$$\Gamma_{p,q}(x+1) = \epsilon_{p,q}(x) \Gamma_{p,q}(x), \quad (3.13)$$

where $\epsilon_{p,q}(x) = -[x]$ for $|x|_p = 1$, and $\epsilon_{p,q}(x) = -1$ for $|x|_p < 1$, (see [8]). By (3.13), we easily see that (cf. [6])

$$\log \Gamma_{p,q}(x+1) = \log \epsilon_{p,q}(x) + \log \Gamma_{p,q}(x). \quad (3.14)$$

By the differentiation of both sides in (3.14), we have (cf. [6])

$$\frac{\Gamma'_{p,q}(x+1)}{\Gamma_{p,q}(x+1)} = \frac{\Gamma'_{p,q}(x)}{\Gamma_{p,q}(x)} + \frac{\epsilon'_{p,q}(x)}{\epsilon_{p,q}(x)}. \quad (3.15)$$

By (3.15), we easily see that (cf. [6])

$$\frac{\Gamma'_{p,q}(x)}{\Gamma_{p,q}(x)} = \left(\sum_{j=1}^{x-1} \frac{q^j}{[j]} \right) \frac{\log q}{q-1} + \frac{\Gamma'_{p,q}(1)}{\Gamma_{p,q}(1)}. \quad (3.16)$$

Define

$$L_{p,q}(x) = \sum_{j=0}^{x-1} \frac{\epsilon'_{p,q}(j)}{\epsilon_{p,q}(j)}. \quad (3.17)$$

It is easy to check that $L_{p,q}(1) = 0$. By (3.15), we also see that

$$\frac{\Gamma'_{p,q}(x)}{\Gamma_{p,q}(x)} = L_{p,q}(x) + \frac{\Gamma'_{p,q}(1)}{\Gamma_{p,q}(1)}, \quad \text{for } x \in \mathbb{Z}_p, \quad (3.18)$$

where $L_{p,q}(x)$ denotes the indefinite sum of $\epsilon'_{p,q}(x)/\epsilon_{p,q}(x)$. By using (3.18) after substituting $x = 1$, we obtain $L_{p,q}(1) = 0$. The classical Euler constant was known as $\gamma = -\Gamma'(1)/\Gamma(1)$. In [8], Koblitz defined the p -adic q -Euler constant $\gamma_{p,q} = -\Gamma'_{p,q}(1)/\Gamma_{p,q}(1)$ (cf. [6, 8]). By using (3.16) and the congruence of Andrews (cf. [3]), we obtain the following congruence:

$$\frac{q-1}{\log q} \left(\frac{\Gamma'_{p,q}(p)}{\Gamma_{p,q}(p)} - \gamma_{p,q} \right) = \sum_{j=1}^{p-1} \frac{q^j}{[j]} \equiv \frac{p-1}{2} (q-1) \pmod{[p]}. \quad (3.19)$$

ACKNOWLEDGMENTS. The author expresses his gratitude to the referees for their valuable suggestions and comments. This work was supported by the Korea Research Foundation Grant (KRF-2002-050-C00001).

REFERENCES

- [1] I. Cherednik, *On q -analogues of Riemann's zeta function*, Selecta Math. (N.S.) **7** (2001), no. 4, 447–491.
- [2] T. Kim, *On explicit formulas of p -adic q -L-functions*, Kyushu J. Math. **48** (1994), no. 1, 73–86.
- [3] ———, *On p -adic q -L-functions and sums of powers*, Discrete Math. **252** (2002), no. 1–3, 179–187.
- [4] ———, *q -Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), no. 3, 288–299.
- [5] ———, *Non-archimedean q -integrals associated with multiple Changhee q -Bernoulli polynomials*, Russ. J. Math. Phys. **10** (2003), 91–98.
- [6] T. Kim, L. C. Jang, K-H. Koh, and I-S. Pyung, *A note on analogue of Γ -functions*, Proceedings of the Conference on 5th Transcendental Number Theory, vol. 5, no. 1, Gakushuin University, Tokyo, 1997, pp. 37–44.
- [7] T. Kim and S-H. Rim, *Generalized Carlitz's q -Bernoulli numbers in the p -adic number field*, Adv. Stud. Contemp. Math. (Pusan) **2** (2000), 9–19.
- [8] N. Koblitz, *q -extension of the p -adic gamma function*, Trans. Amer. Math. Soc. **260** (1980), no. 2, 449–457.
- [9] Q.-M. Luo, Z.-L. Wei, and F. Qi, *Lower and upper bounds of $\zeta(3)$* , Adv. Stud. Contemp. Math. (Kyungshang) **6** (2003), no. 1, 47–51.

Taekyun Kim: Institute of Science Education, Kongju National University, Kongju 314-701, Korea

E-mail address: tkim@kongju.ac.kr