A GENERALIZATION OF A NECESSARY AND SUFFICIENT CONDITION FOR PRIMALITY DUE TO VANTIEGHEM

L. J. P. KILFORD

Received 10 March 2004

We present a family of congruences which hold if and only if a natural number n is prime.

2000 Mathematics Subject Classification: 11A51, 11A07.

The subject of primality testing has been in the mathematical and general news recently, with the announcement [1] that there exists a polynomial-time algorithm to determine whether an integer p is prime or not.

There are older deterministic primality tests which are less efficient; the classical example is Wilson's theorem, that

$$(n-1)! \equiv -1 \pmod{n} \quad (1)$$

if and only if n is prime. Although this is a deterministic algorithm, it does not provide a workable primality test because it requires much more calculation than trial division.

This note provides another family of congruences satisfied by primes and only by primes; it is a generalization of previous work. They could be used as examples of primality tests for students studying elementary number theory.

In Guy [3, Problem A17], the following result due to Vantieghem [4] is quoted as follows.

THEOREM 1 (Vantieghem [4]). *Let n be a natural number greater than 1. Then n is prime if and only if*

$$\prod_{d=1}^{n-1} (1-2^d) \equiv n \pmod{(2^n-1)}. \quad (2)$$

In this note, we will generalize this result to obtain the following theorem.

THEOREM 2. *Let m and n be natural numbers greater than 1. Then n is prime if and only if*

$$\prod_{d=1}^{n-1} (1-m^d) \equiv n \pmod{\frac{m^n-1}{m-1}}. \quad (3)$$

We note that these congruences are also much less efficient than trial division.

PROOF. We follow the method of Vantieghem, using a congruence satisfied by cyclotomic polynomials.

LEMMA 3 (Vantieghem). *Let m be a natural number greater than 1 and let $\Phi_m(X)$ be the m th cyclotomic polynomial. Then*

$$\prod_{\substack{d=1 \\ (d,m)=1}}^m (X - Y^d) \equiv \Phi_m(X) \pmod{\Phi_m(Y)} \quad \text{in } \mathbb{Z}[X, Y]. \quad (4)$$

PROOF OF LEMMA 3. We can write

$$\prod_{\substack{d=1 \\ (d,m)=1}}^m (X - Y^d) - \Phi_m(X) = f_0(Y) + f_1(Y)X + f_2(Y)X^2 + \cdots. \quad (5)$$

(Here the f_i are polynomials over $\mathbb{Z}$.)

Let ζ be a primitive m th root of unity. Now, if $Y = \zeta$, then we see that the left-hand side of this expression is identically 0 in X .

This implies that the f_i are zero at every ζ and every i . Therefore, we have $f_i(Y) \equiv 0 \pmod{\Phi_m(Y)}$, which is enough to prove the lemma. $\square$

Suppose that the natural number n in [Theorem 2](#) is prime. Let $p := n$. We have that $\Phi_p(X) = X^{p-1} + X^{p-2} + \cdots + X + 1$. Therefore, if we set $m = p$ in [Lemma 3](#), we find that

$$\prod_{d=1}^{p-1} (X - Y^d) \equiv X^{p-1} + X^{p-2} + \cdots + X + 1 \pmod{(Y^{p-1} + \cdots + 1)}. \quad (6)$$

We now set $X = 1$ and $Y = m$, to get

$$\prod_{d=1}^{p-1} (1 - m^d) \equiv p \pmod{\frac{m^p - 1}{m - 1}}. \quad (7)$$

This proves that if p is prime, then the congruence holds.

We now prove the converse, by supposing that the congruence [\(3\)](#) holds, and that p is not prime. Therefore p is composite, and hence has a smallest prime factor q . We write $p = q \cdot a$; now $q \leq a$, and also $p \leq a^2$.

Now we have that $m^a - 1$ divides $m^p - 1$ and $m^a - 1$ divides the product $\prod_{d=1}^{p-1} (m^d - 1)$. By combining this with the congruence [\(3\)](#) in [Theorem 2](#), this implies that $(m^a - 1)/(m - 1)$ divides p . Therefore we have

$$2^a - 1 \leq \frac{m^a - 1}{m - 1} \leq p \leq a^2. \quad (8)$$

The inequality $2^a - 1 \leq a^2$ forces a to be either 2 or 3; this means that $p \in \{4, 6, 9\}$ and $m \in \{2, 3\}$; one can check by hand that the congruence does not hold in this case, so we have proved [Theorem 2](#). $\square$

Guy also asks if there is a relationship between the congruence given by Vantieghem and Wilson's theorem. The following theorem gives an elementary congruence similar to that of Vantieghem between a product over integers and a cyclotomic polynomial. It is in fact equivalent to Wilson's theorem.

THEOREM 4. *Let m be a natural number greater than 2. Define the product $F(X)$ by*

$$F(X) := \prod_{\substack{i=1 \\ (i,m)=1}}^{m-1} (X - i - 1) + 1. \quad (9)$$

Then m is prime if and only if

$$\Phi_m(X) \equiv F(X) \pmod{m}. \quad (10)$$

PROOF OF THEOREM 4. Firstly, we prove that if m is not prime, the congruence (10) in Theorem 4 does not hold.

Recall that $\phi(m)$ is defined to be Euler's totient function; the number of integers in the set $\{1, \dots, m\}$ which are coprime to m .

The coefficient of $X^{\phi(m)-1}$ in $F(X)$ is given by the sum

$$-\sum_{\substack{i=1 \\ (i,m)=1}}^{m-1} (i+1) = -\phi(m) - \sum_{\substack{i=1 \\ (i,m)=1}}^{m-1} i. \quad (11)$$

We find that the following congruence holds:

$$-\phi(m) - \sum_{\substack{i=1 \\ (i,m)=1}}^{m-1} i \equiv -\phi(m) \pmod{m}. \quad (12)$$

This follows from the following identity:

$$\sum_{\substack{i=1 \\ (i,m)=1}}^{m-1} i = \frac{m\phi(m)}{2}. \quad (13)$$

Because $m > 2$, $\phi(m)$ is divisible by 2, the sum on the left-hand side of (12) is a multiple of m . We now use some theorems to be found in a paper by Gallot [2, Theorems 1.1 and 1.4].

THEOREM 5. *Let p be a prime and m a natural number.*

(1) *The following relations between cyclotomic polynomials hold:*

$$\Phi_{pm}(x) = \begin{cases} \Phi_m(x^p) & \text{if } p \mid m, \\ \frac{\Phi_m(x^p)}{\Phi_m(x)} & \text{if } p \nmid m. \end{cases} \quad (14)$$

(2) *If $m > 1$, then*

$$\Phi_n(1) = \begin{cases} p & \text{if } n \text{ is a power of a prime } p, \\ 1 & \text{otherwise.} \end{cases} \quad (15)$$

From these results, we see that if m is not a prime power, we then have $\Phi_n(1) \equiv 1 \pmod{m}$, and $F(1)$ is given by

$$1 + \prod_{\substack{i=1 \\ (i,m)=1}}^{m-1} (-i). \quad (16)$$

We see that this is not congruent to $1 \pmod{m}$ because the product is over those i which are coprime to m , so the product does not vanish modulo m .

If m is a prime power p^n , then we see from [Theorem 5](#) that $\Phi_{p^n}(x) = \Phi_p(x^{p^{n-1}})$; in particular, we see that the coefficient of $x^{\phi(p^n)-1}$ is 0, which differs from the coefficient of $x^{\phi(p^n)-1}$ in $F(X)$.

Therefore, if m is not prime, then the congruence does not hold. We now show that if m is prime, the congruence holds.

If m is prime, then $\Phi_m(x) = x^{m-1} + x^{m-2} + \dots + x + 1$. We consider the polynomials $\Phi_m(X+1)$ and $F(X+1)$. Now, modulo m we have

$$\Phi_m(X+1) = X^{m-1}, \quad F(X+1) = \prod_{\substack{i=1 \\ (i,m)=1}}^{m-1} (X-i) + 1. \quad (17)$$

Now if $x \not\equiv 0 \pmod{m}$, then we see that $\Phi_m(x+1) \equiv 1$ and that $F(x+1) \equiv 1$, because the product vanishes.

And if we have $x = 0$, then $\Phi_m(x) = 0$ and, by Wilson's theorem, $F(0) \equiv (m-1)! + 1 \equiv 0 \pmod{m}$.

Therefore we have proved [Theorem 4](#). □

REFERENCES

- [1] M. Agrawal, N. Kayal, and N. Saxena, *PRIMES is in P*, preprint, 2002, <http://www.cse.iitk.ac.in/news/primality.html>.
- [2] Y. Gallot, *Cyclotomic polynomials and prime numbers*, preprint, 2001, <http://perso.wanadoo.fr/yves.gallot/papers>.
- [3] R. K. Guy, *Unsolved Problems in Number Theory*, 2nd ed., Problem A17. Problem Books in Mathematics, I, Springer-Verlag, New York, 1994.
- [4] E. Vantieghem, *On a congruence only holding for primes*, Indag. Math. (N.S.) **2** (1991), no. 2, 253–255.

L. J. P. Kilford: The University of Oxford, Mathematical Institute, 24–29 Street Giles', Oxford OX1 3LB, UK