ON THE CLASS OF QS -ALGEBRAS

MICHIRO KONDO

Received 19 September 2003

We consider some fundamental properties of QS -algebras and show that (1) the theory of QS -algebras is logically equivalent to the theory of *Abelian groups*, that is, each theorem of QS -algebras is provable in the theory of Abelian groups, and conversely, each theorem of Abelian groups is provable in the theory of QS -algebras; and (2) a G -part $G(X)$ of a QS -algebra X is a normal subgroup generated by the class of all elements of order 2 of X when it is considered as a group.

2000 Mathematics Subject Classification: 06F35, 03G25, 03C07.

1. Introduction. In [3], the notion of Q -algebras is introduced and some fundamental properties are established. The algebras are extensions of the BCK/BCI-algebras which were proposed by Y. Imai and K. Iséki in 1966. It is usually important to generalize the algebraic structures. Neggers and Kim [4] introduced a class of algebras which is related to several classes of algebras such as BCK/BCI/BCH-algebras. They call them *B-algebras*, and they proved that every group $(X; \circ, 0)$ determines a B -algebra $(X; *, 0)$, which is called the *group-derived B-algebra*. Conversely, in [2], we prove that every B -algebra is group-derived and hence that the class of B -algebras and the class of all groups are the same. Ahn and Kim [1] proposed the notion of QS -algebras which is also a generalization of BCK/BCI-algebras and obtained several results. Here, we consider some fundamental properties of QS -algebras and show that

- (1) the theory of QS -algebras is logically equivalent to the theory of *Abelian groups*, that is, each theorem of QS -algebras is provable in the theory of Abelian groups and conversely each theorem of Abelian groups is provable in the theory of QS -algebras;
- (2) a subset $G(X)$ called G -part of a QS -algebra X is a normal subgroup which is generated by the class of all elements of order 2.

2. Preliminaries. A QS -algebra is a nonempty set X with a constant 0 and a binary operation “ $*$ ” satisfying the following axioms:

- (QS1) $x * x = 0$,
- (QS2) $x * 0 = x$,
- (QS3) $(x * y) * z = (x * z) * y$,
- (QS4) $(x * y) * (x * z) = y * z$,

for all x, y, z in X .

EXAMPLE 2.1. (1) Let $X = \{0, 1, 2\}$ be a set with an operation $*$ defined as follows:

$$\begin{array}{c|ccc}
 * & 0 & 1 & 2 \\
 \hline
 0 & 0 & 2 & 1 \\
 1 & 1 & 0 & 2 \\
 2 & 2 & 1 & 0
 \end{array} \tag{2.1}$$

Then $(X; *, 0)$ is a QS -algebra.

(2) Let X be the set of all integers. Define a binary operation $*$ on X by

$$x * y := x - y. \tag{2.2}$$

Then $(X; *, 0)$ is a QS -algebra.

We note that these examples are both Abelian groups and the operation $*$ corresponds to the minus operation “ $-$ ”. In the case of (1), X can be considered as the set $\mathbb{Z}_3$ of integers of modulo 3 and the operation $*$ as a minus “ $-$ ” modulo operation. It seems that any Abelian group gives an example of a QS -algebra. In fact, we can prove the fact.

THEOREM 2.2. *Let $(X; \cdot, -1, e)$ be an Abelian group. If $x * y = x \cdot y^{-1}$ is defined and $0 = e$, then $(X; *, 0)$ is a QS -algebra.*

PROOF. We only show that the conditions (QS3) and (QS4) of QS -algebras are satisfied. For the case of (QS3), since X is an Abelian group,

$$\begin{aligned}
 (x * y) * z &= (x \cdot y^{-1}) \cdot z^{-1} \\
 &= x \cdot (y^{-1} \cdot z^{-1}) \\
 &= x \cdot (z^{-1} \cdot y^{-1}) \\
 &= (x \cdot z^{-1}) \cdot y^{-1} \\
 &= (x * z) * y.
 \end{aligned} \tag{2.3}$$

For the case of (QS4), we also have

$$\begin{aligned}
 (x * y) * (x * z) &= (x \cdot y^{-1}) \cdot (x \cdot z^{-1})^{-1} \\
 &= (x \cdot y^{-1}) \cdot (z \cdot x^{-1}) \\
 &= x \cdot x^{-1} \cdot z \cdot y^{-1} \\
 &= z \cdot y^{-1} \\
 &= z * y.
 \end{aligned} \tag{2.4}$$

□

The theorem means that every Abelian group $(X; \cdot, -1, e)$ determines a QS -algebra $(X; *, 0)$; in other words, any Abelian group can be considered as a QS -algebra. Conversely, we will show in the next section that any QS -algebra determines an Abelian group, that is, every QS -algebra can be considered as an Abelian group. Hence, we are able to conclude that in this sense, the class of QS -algebras coincides with the class of Abelian groups.

3. Abelian groups can be derived from QS-algebras. We show that every QS-algebra determines an Abelian group. In order to do so, it is sufficient to construct an Abelian group from any QS-algebra. We need some lemmas to prove that.

Let $(X; *, 0)$ be a QS-algebra.

LEMMA 3.1. *For all $x, y, z \in X$, if $x * y = z$, then $x * z = y$.*

PROOF. Suppose that $x * y = z$. Then, since X is a QS-algebra, we have $x * z = (x * 0) * (x * y) = y * 0 = y$. $\square$

It follows from the above that the condition (QS4)' is established in any QS-algebra: (QS4)' $(x * z) * (y * z) = x * y$.

COROLLARY 3.2. *If $x * y = 0$, then $x = y$.*

Let $B(X) = \{x \in X \mid 0 * x = 0\}$. A QS-algebra X is called *p-semisimple* if $B(X) = \{0\}$ (cf. [1]). We can show that every QS-algebra is *p-semisimple*.

COROLLARY 3.3. *Every QS-algebra is p-semisimple.*

PROOF. Suppose that X is a QS-algebra. For all elements $x \in X$, since

$$\begin{aligned} x \in B(X) &\iff 0 * x = 0 \\ &\iff x = 0 \quad (\text{by Corollary 3.2}), \end{aligned} \tag{3.1}$$

we can conclude that X is *p-semisimple*. $\square$

REMARK 3.4. It is proved in [1] that every *associative* QS-algebra is *p-semisimple*. The corollary above means that the assumption of associativity is superfluous.

LEMMA 3.5. $0 * (x * y) = y * x$.

PROOF. $0 * (x * y) = (x * x) * (x * y) = y * x$. $\square$

COROLLARY 3.6. $0 * (0 * x) = x$.

LEMMA 3.7. $x * (0 * y) = y * (0 * x)$.

PROOF. Since

$$\begin{aligned} 0 * (x * (y * (0 * x))) &= (y * (0 * x)) * x \quad (\text{by Lemma 3.5}) \\ &= (y * x) * (0 * x) \quad (\text{by (QS3)}) \\ &= y * 0 \quad (\text{by (QS4)'}) \\ &= y, \end{aligned} \tag{3.2}$$

we have $0 * (0 * (x * (y * (0 * x)))) = 0 * y$. It follows from Corollary 3.6 that $x * (y * (0 * x)) = 0 * y$ and hence $x * (0 * y) = y * (0 * x)$ by Lemma 3.1. $\square$

These lemmas provide a proof that we can construct an Abelian group $(X; \cdot, e)$ from a QS-algebra $(X; *, 0)$.

THEOREM 3.8. *Let $(X; *, 0)$ be a QS-algebra. If $x \cdot y = x * (0 * y)$ is defined, $x^{-1} = 0 * x$, and $e = 0$, then the structure $(X; \cdot, -1, e)$ is an Abelian group.*

PROOF. We only show that the structure $(X; \cdot, -1, e)$ satisfies the conditions of associativity and of commutativity with respect to the operation “ $\cdot$ ”.

For associativity, we have

$$\begin{aligned}
 (x \cdot y) \cdot z &= (x * (0 * y)) * (0 * z) \\
 &= (y * (0 * x)) * (0 * z) \quad (\text{by Lemma 3.7}) \\
 &= (y * (0 * z)) * (0 * x) \\
 &= x * (0 * (y * (0 * z))) \quad (\text{by Lemma 3.7}) \\
 &= x \cdot (y \cdot z).
 \end{aligned} \tag{3.3}$$

For commutativity, we also have $x \cdot y = x * (0 * y) = y * (0 * x) = y \cdot x$. $\square$

Combining Theorems 2.2 and 3.8, we can conclude that the class of QS -algebras coincides with the class of Abelian groups.

In the following, we will describe our results in greater detail. We can show that each theorem of QS -algebras is translated to a formula of $\mathcal{AG}$ which is provable in the theory of Abelian groups and conversely each theorem of Abelian groups is proved in the theory of QS -algebras. To present our theorem precisely, we will develop the formal theories of QS -algebras and Abelian groups. Let $\mathcal{QS}$ and $\mathcal{AG}$ be the theories of QS -algebras and Abelian groups, respectively. Theories consist of languages and axioms. At first, we define languages of these theories which are needed to present statements formally in their theories. By $\mathcal{L}(\mathcal{QS})$ (or $\mathcal{L}(\mathcal{AG})$), we mean a language of the theory $\mathcal{QS}$ of QS -algebras (or the theory $\mathcal{AG}$ of groups). We define them as follows.

The language of the theory of $\mathcal{QS}$ of QS -algebras consists of

- (lq1) countable variables $x, y, z, \dots$,
- (lq2) binary operation symbol $*$,
- (lq3) constant symbol 0 ;

and the language of the theory of $\mathcal{AG}$ of QS -algebras consists of

- (lg1) countable variables $x, y, z, \dots$,
- (lg2) binary operation symbol $\circ$,
- (lg3) unary operation symbol -1 ,
- (lg4) constant symbol e .

Next we define *terms* which represent objects in the theory. By $\mathcal{T}(\mathcal{QS})$ (or $\mathcal{T}(\mathcal{AG})$) we mean the set of terms of $\mathcal{QS}$ (or $\mathcal{AG}$). Terms are defined as follows.

For terms of $\mathcal{QS}$,

- (tb1) each variable is a term,
- (tb2) the constant 0 is a term,
- (tb3) if u and v are terms, then $u * v$ is a term.

For terms of $\mathcal{AG}$,

- (tg1) each variable is a term,
- (tg2) the constant e is a term,
- (tg3) if u and v are terms, then so are $u \circ v$ and u^{-1} .

We also define *formulas* which represent statements in each theory. Formulas of $\mathcal{QS}$ (or $\mathcal{AG}$) are defined as the forms of $s = t$, where $s, t \in \mathcal{T}(\mathcal{QS})$ (or $s, t \in \mathcal{T}(\mathcal{AG})$). By $\mathcal{F}(\mathcal{QS})$

(or $\mathcal{F}(\mathcal{A}\mathcal{G})$) we mean the set of formulas of $\mathcal{Q}\mathcal{S}$ (or $\mathcal{A}\mathcal{G}$). We denote formulas simply by $A, B, C, \dots$.

As to the axioms of QS -algebras, we list the following:

$$(QS1) \quad x * x = 0,$$

$$(QS2) \quad x * 0 = x,$$

$$(QS3) \quad (x * y) * z = (x * z) * y,$$

$$(QS4) \quad (x * y) * (x * z) = y * z.$$

For the axioms of Abelian groups, we use the following:

$$(G1) \quad x \circ (y \circ z) = (x \circ y) \circ z,$$

$$(G2) \quad x \circ e = e \circ x = x,$$

$$(G3) \quad x \circ x^{-1} = x^{-1} \circ x = e,$$

$$(G4) \quad x \circ y = y \circ x.$$

Two formal theories $\mathcal{Q}\mathcal{S}$ and $\mathcal{A}\mathcal{G}$ have the same rules of inference concerning “equality,” for they have no predicate symbols.

RULES OF INFERENCE. For all terms $s, t, w, s_1, s_2, \dots \in \mathcal{T}(\mathcal{Q}\mathcal{S})$ (or $\mathcal{T}(\mathcal{A}\mathcal{G})$),

$$s = s, \quad \frac{s = t}{t = s}, \quad \frac{s = t, t = w}{s = w}, \quad \frac{s_1 = t_1, \dots, s_n = t_n}{\phi(s_1, \dots, s_n) = \phi(t_1, \dots, t_n)}, \quad (3.4)$$

where $\phi(x_1, \dots, x_n)$ is a term of $\mathcal{T}(\mathcal{Q}\mathcal{S})$ (or $\mathcal{T}(\mathcal{A}\mathcal{G})$) whose variables are contained in $\{x_1, \dots, x_n\}$.

We are now ready to present a formal theory of Abelian groups and QS -algebras. Let Γ be a subset of formulas of $\mathcal{F}(\mathcal{Q}\mathcal{S})$ (or $\mathcal{F}(\mathcal{A}\mathcal{G})$). By

$$\Gamma \vdash_{\mathcal{Q}\mathcal{S}} A (\Gamma \vdash_{\mathcal{A}\mathcal{G}} A), \quad (3.5)$$

we mean that there is a finite sequence of formulas $A_1, A_2, \dots, A_n$ of $\mathcal{F}(\mathcal{Q}\mathcal{S})$ ($\mathcal{F}(\mathcal{A}\mathcal{G})$) such that for each i ,

- (1) A_i is an axiom of $\mathcal{Q}\mathcal{S}(\mathcal{A}\mathcal{G})$,
- (2) $A_i \in \Gamma$,
- (3) there exists $j_1, \dots, j_k$ ($j_1, \dots, j_k < i$) such that

$$\frac{A_{j_1}, \dots, A_{j_k}}{A}. \quad (3.6)$$

We say that A is provable from Γ in $\mathcal{Q}\mathcal{S}(\mathcal{A}\mathcal{G})$ when $\Gamma \vdash_{\mathcal{A}\mathcal{G}} A (\Gamma \vdash_{\mathcal{Q}\mathcal{S}} A)$. In particular, in case of $\Gamma = \emptyset$, we say that A is a theorem of $\mathcal{Q}\mathcal{S}(\mathcal{A}\mathcal{G})$ and simply denote it by $\vdash_{\mathcal{Q}\mathcal{S}} A (\vdash_{\mathcal{A}\mathcal{G}} A)$.

As an example, we present the following which is called a cancelation rule in the theory of groups:

$$x \circ y = z \circ x \vdash_{\mathcal{A}\mathcal{G}} y = z. \quad (3.7)$$

Indeed, we have the following finite sequence of formulas:

$$\begin{aligned}
 x \circ y &= z \circ x, & z \circ x &= x \circ z, & x \circ y &= x \circ z, & x^{-1} \circ (x \circ y) &= x^{-1} \circ (x \circ z), \\
 x^{-1} \circ (x \circ y) &= (x^{-1} \circ x) \circ y, & x^{-1} \circ (x \circ z) &= (x^{-1} \circ x) \circ z, \\
 (x^{-1} \circ x) \circ y &= (x^{-1} \circ x) \circ z, & x^{-1} \circ x &= e, & e \circ y &= e \circ z, \\
 e \circ y &= y, & e \circ z &= z, & y &= z.
 \end{aligned} \tag{3.8}$$

Next, we define two maps ξ from the theory $\mathcal{QS}$ of QS -algebras to the theory $\mathcal{AG}$ of Abelian groups and η from $\mathcal{AG}$ to $\mathcal{QS}$ as follows. For $\mathcal{T}(\mathcal{QS})$,

$$\begin{aligned}
 \xi(x) &\equiv x \quad \text{for each variable } x, \\
 \xi(0) &\equiv e, \\
 \xi(s * t) &\equiv \xi(s) \circ \xi(t^{-1}),
 \end{aligned} \tag{3.9}$$

and for $\mathcal{F}(\mathcal{QS})$,

$$\begin{aligned}
 \xi(s = t) &\equiv \xi(s) = \xi(t), \\
 \xi(s = t \Rightarrow s' = t') &\equiv \xi(s = t) \Rightarrow \xi(s' = t'),
 \end{aligned} \tag{3.10}$$

where $s, s', t, t' \in \mathcal{T}(\mathcal{QS})$.

Conversely, we define a map $\eta: \mathcal{AG} \rightarrow \mathcal{QS}$ as follows. For $\mathcal{T}(\mathcal{AG})$,

$$\begin{aligned}
 \eta(x) &\equiv x \quad \text{for each variable } x, \\
 \eta(e) &\equiv 0, \\
 \eta(s^{-1}) &\equiv 0 * (\eta(s)), \\
 \eta(s \circ t) &\equiv \eta(s) * (0 * \eta(t)),
 \end{aligned} \tag{3.11}$$

and for $\mathcal{F}(\mathcal{AG})$,

$$\begin{aligned}
 \eta(s = t) &\equiv \eta(s) = \eta(t), \\
 \eta(s = t \Rightarrow s' = t') &\equiv \eta(s = t) \Rightarrow \eta(s' = t'),
 \end{aligned} \tag{3.12}$$

where $s, s', t, t' \in \mathcal{T}(\mathcal{AG})$.

We are now ready to state our theorem. It is as follows.

MAIN THEOREM 3.9. (1) For every $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{QS})$, if $\Gamma \vdash_{\mathcal{QS}} A$, then $\xi(\Gamma) \vdash_{\mathcal{AG}} \xi(A)$; conversely,

- (2) for every $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{AG})$, if $\Gamma \vdash_{\mathcal{AG}} A$, then $\eta(\Gamma) \vdash_{\mathcal{QS}} \eta(A)$; moreover,
- (3) $\Gamma \vdash_{\mathcal{QS}} \eta\xi(A)$ if and only if $\Gamma \vdash_{\mathcal{QS}} A$ for every $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{QS})$;
- (4) $\Gamma \vdash_{\mathcal{AG}} \xi\eta(A)$ if and only if $\Gamma \vdash_{\mathcal{AG}} A$ for every $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{AG})$.

This means that each theorem of QS -algebras can be translated immediately to that of groups and conversely, every theorem of groups is applied to that of QS -algebras.

At first we will establish the former part, that is, if $\Gamma \vdash_{\mathcal{QS}} A$, then $\xi(\Gamma) \vdash_{\mathcal{AG}} \xi(A)$.

THEOREM 3.10. For $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{Q}\mathcal{S})$, if $\Gamma \vdash_{\mathcal{Q}\mathcal{S}} A$, then $\xi(\Gamma) \vdash_{\mathcal{A}\mathcal{G}} \xi(A)$.

PROOF. It is sufficient to show that for every axiom A of QS-algebras, $\xi(A)$ is provable in the theory $\mathcal{A}\mathcal{G}$ of Abelian groups. For the sake of simplicity, we treat only the case of axiom (QS3): $(x * y) * z = x * (z * (0 * y))$. Other cases can be proved similarly.

Since

$$\begin{aligned} \xi((x * y) * z) &= (x \circ y^{-1}) \circ z^{-1}, \\ \xi(x * (z * (0 * y))) &= x \circ \{z \circ (y^{-1})^{-1}\}^{-1}, \end{aligned} \quad (3.13)$$

we have to show that

$$(x \circ y^{-1}) \circ z^{-1} = x \circ \{z \circ (y^{-1})^{-1}\}^{-1}. \quad (3.14)$$

We have the following:

$$\begin{aligned} x \circ \{z \circ (y^{-1})^{-1}\}^{-1} &= x \circ (z \circ y)^{-1} \\ &= x \circ (y^{-1} \circ z^{-1}) \\ &= (x \circ y^{-1}) \circ z^{-1}. \end{aligned} \quad (3.15)$$

Hence, if $\Gamma \vdash_{\mathcal{Q}\mathcal{S}} A$, then $\xi(\Gamma) \vdash_{\mathcal{A}\mathcal{G}} \xi(A)$. □

Conversely, we can show that if $\Gamma \vdash_{\mathcal{A}\mathcal{G}} A$, then $\eta(\Gamma) \vdash_{\mathcal{Q}\mathcal{S}} \eta(A)$.

THEOREM 3.11. For $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{A}\mathcal{G})$, if $\Gamma \vdash_{\mathcal{A}\mathcal{G}} A$, then $\eta(\Gamma) \vdash_{\mathcal{Q}\mathcal{S}} \eta(A)$.

PROOF. As above, it is sufficient to show that $\eta(A)$ is provable in the theory $\mathcal{Q}\mathcal{S}$ of QS-algebras for every axiom A of the theory of Abelian groups.

For the case of (G1), we have to show that

$$(x * (0 * y)) * (0 * z) = x * (0 * (y * (0 * z))) \quad (3.16)$$

because $\eta((x \circ y) \circ z) = (x * (0 * y)) * (0 * z)$ and $\eta(x \circ (y \circ z)) = x * (0 * (y * (0 * z)))$. By the proposition above, we have

$$\begin{aligned} (x * (0 * y)) * (0 * z) &= x * ((0 * z) * (0 * (0 * y))) \\ &= x * ((0 * z) * y) \\ &= x * (0 * (y * (0 * z))). \end{aligned} \quad (3.17)$$

Other cases are proved easily, so we omit their proofs.

The theorem can be proved completely. □

Moreover, it follows from Lemma 3.5 and Corollary 3.6 that we have $\vdash_{\mathcal{Q}\mathcal{S}} \eta\xi(t) = t$ for every term $t \in \mathcal{T}(\mathcal{Q}\mathcal{S})$. For the case of Abelian groups, it is easy to prove that $\vdash_{\mathcal{A}\mathcal{G}} \xi\eta(s) = s$ for every term $s \in \mathcal{T}(\mathcal{A}\mathcal{G})$. Thus we have the following.

THEOREM 3.12. For these maps,

- (1) $\Gamma \vdash_{\mathcal{Q}\mathcal{S}} \eta\xi(A)$ if and only if $\Gamma \vdash_{\mathcal{Q}\mathcal{S}} A$ for all $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{Q}\mathcal{S})$,
- (2) $\Gamma \vdash_{\mathcal{A}\mathcal{G}} \xi\eta(A)$ if and only if $\Gamma \vdash_{\mathcal{A}\mathcal{G}} A$ for all $\Gamma \cup \{A\} \in \mathcal{F}(\mathcal{A}\mathcal{G})$.

4. Some properties. In this section, we prove other properties of QS -algebras, especially properties about the G -part and *mediality*. That is,

- (1) the G -part $G(X)$ of a QS -algebra X is a normal subgroup generated by the class of all elements of order 2 of X ;
- (2) every QS -algebra X is *medial*, that is, it satisfies the condition

$$(x * y) * (z * u) = (x * z) * (y * u) \quad (4.1)$$

for all elements $x, y, z, u \in X$.

Let X be a QS -algebra. A subset $G(X) = \{x \in X \mid 0 * x = x\}$ is called a G -part of X . For the G -part of X , we have the following results.

PROPOSITION 4.1. *If $x, y \in G(X)$, then $x * y = y * x$.*

PROOF. Suppose $x, y \in G(X)$. Since $0 * x = x$ and $0 * y = y$, we have

$$\begin{aligned} x * y &= (0 * x) * (0 * y) \\ &= (0 * (0 * y)) * x \quad (\text{by (QS3)}) \\ &= y * x. \end{aligned} \quad (4.2) \quad \square$$

PROPOSITION 4.2. *If $x, y \in G(X)$, then $x * y \in G(X)$.*

PROOF. Suppose that $x, y \in G(X)$. It follows that $(0 * (x * y)) * (x * y) = (y * x) * (x * y) = (x * y) * (x * y) = 0$ by Lemma 3.5. Thus we have $0 * (x * y) = x * y$, that is, $x * y \in G(X)$. $\square$

Since any QS -algebra X may be considered as an Abelian group, Proposition 4.2 implies that $G(X)$ is a (normal) subgroup of X . Moreover, since $x^2 = x \cdot x = x * (0 * x) = x * x = 0$ for $x \in G(X)$, every nonunit element in $G(X)$ is of order 2. Hence, we can conclude that the G -part $G(X)$ is the normal subgroup generated by the class of all elements of order 2. It is easy to show that $G(X) = \{x \in X \mid x \text{ is of order 2}\} \cup \{0\}$.

For the statement (2) above, in [1, Theorem 3.6], it is proved that a QS -algebra X is medial if and only if the condition $x * (y * z) = (x * y) * (0 * z)$ holds for all $x, y, z \in X$. On the other hand, by Lemma 3.7, we have $(x * y) * (0 * z) = z * (0 * (x * y)) = z * (y * x)$. Thus X is medial if and only if the condition $x * (y * z) = z * (y * x)$ holds for all $x, y, z \in X$. By using this characterization of mediality, we will prove the following.

THEOREM 4.3. *Every QS -algebra is medial.*

PROOF. It is sufficient to show that $x * (y * z) = z * (y * x)$ holds for all $x, y, z \in X$. Since

$$\begin{aligned} x * (y * z) &= 0 * (0 * (x * (y * z))) \quad (\text{by Corollary 3.6}) \\ &= 0 * ((y * z) * x) \quad (\text{by Lemma 3.5}) \\ &= 0 * ((y * x) * z) \quad (\text{by (QS3)}) \\ &= z * (y * x) \quad (\text{by Lemma 3.5}), \end{aligned} \quad (4.3)$$

it follows that X is medial. $\square$

5. Application. Let $V = \{x, y, z, \dots\}$ be a set of variables and $\mathbf{0}$ a constant. We define a term and equation as follows:

- (1) $\mathbf{0}$ is a term;
- (2) each variable in V is a term;
- (3) if s, t are terms, then $s * t$ is also a term;
- (4) if s, t are terms, then $s = t$ is an equation.

Thus, for example, $\mathbf{0}, \mathbf{0} * x, x * (\mathbf{0} * y), x * y$ are terms and thus $\mathbf{0} = \mathbf{0} * x, x * (\mathbf{0} * y) = x * y$ are equations. By $t(x, y, \dots)$ we mean a term whose variables are in $\{x, y, \dots\}$. We say that an equation $s(x, y, \dots) = t(x, y, \dots)$ is satisfied in a QS-algebra X when for all elements $a, b, \dots \in X$, we have $u^X(a, b, \dots) = v^X(a, b, \dots)$. In particular, an equation $t(x, y, \dots) = \mathbf{0}$ is said to be satisfied in X if $t^X(a, b, \dots) = 0$ for all elements $a, b, \dots \in X$. In the following, by $t(a, b, \dots)$ we mean an element $t^X(a, b, \dots)$ which is an interpretation of a term $t(x, y, \dots)$ in X , that is, $t(a, b, \dots)$ is an abbreviation of $t^X(a, b, \dots)$.

We also define a condition (C) which plays an important role to develop our theory:

(C) for all x and for all y , there exists $t(x, y)$ such that

$$(\mathbf{0} * (\mathbf{0} * x)) * (\mathbf{0} * (\mathbf{0} * y)) = \mathbf{0} * (\mathbf{0} * t(x, y)), \quad t(x, x) = \mathbf{0}. \quad (5.1)$$

By using condition (C), we have the following theorem which shows the relation between Q-algebras and QS-algebras.

THEOREM 5.1. *Let $(X; *, 0)$ be a Q-algebra. $(X; *, 0)$ satisfies condition (C) if and only if $(X^*; *, 0)$ is a QS-algebra, where $X^* = \{0 * (0 * a) \mid a \in X\}$.*

PROOF. *If part.* For all $u, v \in X^*$, there exist $a, b \in X$ such that $u = 0 * (0 * a)$, $v = 0 * (0 * b)$. It follows from condition (C) that $u * v \in X^*$ and that X^* is a subalgebra of X . Hence, X^* is a Q-algebra. We define $u \cdot v = u * (0 * v)$ and $u^{-1} = 0 * u$. Since $u, v, 0 \in X^*$, we have $u \cdot v \in X^*$. Moreover for this operation, we can show that

- (i) $u \cdot v = v \cdot u$,
- (ii) $u \cdot 0 = 0 \cdot u = u$,
- (iii) $u \cdot (0 * u) = (0 * u) \cdot u = 0$,
- (iv) $(u \cdot v) \cdot w = u \cdot (v \cdot w)$.

For the sake of simplicity, we only prove the case of (iv). Before doing so, we note the following result: $(u \cdot v) \cdot w = (u \cdot w) \cdot v$ for all $u, v, w \in X^*$. Because

$$\begin{aligned} (u \cdot v) \cdot w &= (u \cdot v) * (0 * w) \\ &= (u * (0 * v)) * (0 * w) \\ &= (u * (0 * w)) * (0 * v) \\ &= (u \cdot w) \cdot v, \end{aligned} \quad (5.2)$$

it follows from the result that $(u \cdot v) \cdot w = (v \cdot u) \cdot w = (v \cdot w) \cdot u = u \cdot (v \cdot w)$.

Thus the above means that $(X^*; \cdot, -1, 0)$ is an Abelian group. For this group, if we define $u \circ v = u \cdot (0 * v)$, then $(X^*; \circ, 0)$ is a QS-algebra. Clearly we have $u \circ v = u \cdot (0 * v) = u * (0 * (0 * v)) = u * v$ for all $u, v \in X^*$. That is, $(X^*; *, 0)$ is a QS-algebra.

Only if part. Conversely, we suppose that $(X^*; *, 0)$ is a QS-algebra. For all $u, v \in X^*$, there exist $a, b \in X$ such that $u = 0 * (0 * a)$, $v = 0 * (0 * b)$. Since $u * v \in X^*$, $u * v$ has to have a form of $0 * (0 * t(a, b))$. This means that X^* satisfies the condition

$$\forall x \forall y \exists t(x, y) \quad (0 * (0 * x)) * (0 * (0 * y)) = 0 * (0 * t(x, y)). \quad (5.3)$$

It is obvious that $t(a, a) = 0$ for all $a \in X$, that is, $t(x, x) = 0$. Thus X^* satisfies condition (C). $\square$

We consider some cases of $t(x, y)$ as corollaries to the theorem. First of all, let $t(x, y)$ be a form of $x * y$, that is, $t(x, y) = x * y$. In this case, condition (C) has a form of

$$(0 * (0 * x)) * (0 * (0 * y)) = 0 * (0 * (x * y)). \quad (5.4)$$

For a map $f : X \rightarrow X^*$ defined by $f(x) = 0 * (0 * x)$, since $(x, y) \in \text{Ker } f$ if and only if $0 * (0 * x) = 0 * (0 * y)$, we have that $X / \text{Ker } f$, the quotient Q -algebra modulo $\text{Ker } f$, is isomorphic to X^* , that is, $X / \text{Ker } f \cong X^*$. Hence, we have the following.

COROLLARY 5.2. *If $f : X \rightarrow X^*$ is a map defined by $f(x) = 0 * (0 * x)$, then $X / \text{Ker } f \cong X^*$.*

We define a term $t^n(x, y)$ for all nonnegative integers n as follows:

$$\begin{aligned} t^0(x, y) &= 0 * (x * y), \\ t^n(x, y) &= t^{n-1}(x, y) * (0 * (0 * (x * y))). \end{aligned} \quad (5.5)$$

In this case, the corresponding condition (C_n) is

$$(0 * (0 * x)) * (0 * (0 * y)) = 0 * (0 * t^n(x, y)). \quad (5.6)$$

We now have the following result as to condition (C_n) .

COROLLARY 5.3. *Let X be a QS-algebra. If X satisfies condition (C_n) , then X^* is an Abelian group in which every element has order at most $(n + 2)$.*

PROOF. For condition (C_n) , if we take $y = 0$, then we have $0 * (0 * x) = 0 * (0 * t^n(x, 0))$, that is,

$$0 * (0 * x) = 0 * [0 * \{((0 * x) * (0 * (0 * x))) * (0 * (0 * x)) * \cdots * (0 * (0 * x))\}]. \quad (5.7)$$

Since any element $u \in X^*$ has a form of $0 * (0 * a)$ for some element $a \in X$, it follows from (C_n) that

$$\begin{aligned} u &= 0 * [0 * \{((0 * a) * (0 * (0 * a))) * (0 * (0 * a)) * \cdots * (0 * (0 * a))\}] \\ &= 0 * [0 * \{((\cdots ((0 * a) * u) * u) * \cdots) * u\}] \\ &= ((\cdots ((0 * a) * u) * u) * \cdots) * u \\ &= ((\cdots ((0 * u) * u) * u) * \cdots) * u. \end{aligned} \quad (5.8)$$

On the other hand, since $(0 * u) * u = (0 * u) * (0 * (0 * u)) = u^{-1} \cdot u^{-1}$ in the Abelian group X^* , we have

$$u = (u^{-1} \cdot u^{-1}) \cdot u^{-1} \cdots u^{-1} = (u^{-1})^{n+1} = u^{-(n+1)} \quad (5.9)$$

and hence $u^{n+2} = 0$. This means that each element of X^* has order at most $n + 2$. $\square$

As the last case, we suppose $t(x, y) = (x * y) * (y * x)$. Since condition (C) is

$$(0 * (0 * x)) * (0 * (0 * y)) = 0 * (0 * ((x * y) * (y * x))) \quad (5.10)$$

in this case, if we take $y = 0$, then we have the condition

$$0 * (0 * x) = 0 * (0 * (x * (0 * x))). \quad (5.11)$$

This implies that

$$0 * (0 * a) = 0 * (0 * (a * (0 * a))) \quad (5.12)$$

for all $a \in X$. In particular, any element $u \in X^*$ satisfies the condition. Hence, since $u = 0 * (0 * u)$ for all elements $u \in X^*$, we have in the Abelian group X^*

$$u = u * (0 * u) = u \cdot u. \quad (5.13)$$

This means that every element of X^* is an idempotent. But $u \cdot uu$ means $u = 0$ and $X^* = \{0\}$ is trivial.

COROLLARY 5.4. *Let X be a Q -algebra. If X satisfies the condition*

$$(0 * (0 * x)) * (0 * (0 * y)) = 0 * (0 * ((x * y) * (y * x))), \quad (5.14)$$

then X^ is an Abelian group in which every element is an idempotent.*

REFERENCES

- [1] S. S. Ahn and H. S. Kim, *On QS-algebras*, J. Chungcheong Math. Soc. **12** (1999), 33–41.
- [2] M. Kondo and Y. B. Jun, *The class of B-algebras coincides with the class of groups*, Sci. Math. Jpn. **57** (2003), no. 1, 197–199.
- [3] J. Neggers, S. S. Ahn, and H. S. Kim, *On Q-algebras*, Int. J. Math. Math. Sci. **27** (2001), no. 12, 749–757.
- [4] J. Neggers and H. S. Kim, *On B-algebras*, Mat. Vesnik **54** (2002), no. 1-2, 21–29.

Michiro Kondo: School of Information Environment, Tokyo Denki University, Inzai 270-1382, Japan

E-mail address: kondo@sie.dendai.ac.jp