CONTINUITY FOR SOME MULTILINEAR OPERATORS OF INTEGRAL OPERATORS ON TRIEBEL-LIZORKIN SPACES

LIU LANZHE

Received 15 March 2003

The continuity for some multilinear operators related to certain fractional singular integral operators on Triebel-Lizorkin spaces is obtained. The operators include Calderon-Zygmund singular integral operator and fractional integral operator.

2000 Mathematics Subject Classification: 42B20, 42B25.

1. Introduction. Let T be a Calderon-Zygmund singular integral operator; a well-known result of Coifman et al. (see [6]) states that the commutator $[b, T] = T(bf) - bTf$ (where $b \in BMO$) is bounded on $L^p(R^n)$ for $1 < p < \infty$; Chanillo (see [1]) proves a similar result when T is replaced by the fractional integral operator; in [10, 11], these results on the Triebel-Lizorkin spaces and the case $b \in \text{Lip } \beta$ (where $\text{Lip } \beta$ is the homogeneous Lipschitz space) are obtained. The main purpose of this paper is to discuss the continuity for some multilinear operators related to certain fractional singular integral operators on the Triebel-Lizorkin spaces. In fact, we will establish the continuity on the Triebel-Lizorkin spaces for the multilinear operators only under certain conditions on the size of the operators. As to the applications, the continuity for the multilinear operators related to the Calderon-Zygmund singular integral operator and fractional integral operator on the Triebel-Lizorkin spaces is obtained.

2. Notations and results. Throughout this paper, Q will denote a cube of R^n with side parallel to the axes, and for a cube Q , let $f_Q = |Q|^{-1} \int_Q f(x) dx$ and $f^\#(x) = \sup_{x \in Q} |Q|^{-1} \int_Q |f(y) - f_Q| dy$. For $1 \leq r < \infty$ and $0 \leq \delta < n$, let

$$M_{\delta,r}(f)(x) = \sup_{x \in Q} \left(\frac{1}{|Q|^{1-r\delta/n}} \int_Q |f(y)|^r dy \right)^{1/r}; \quad (2.1)$$

we denote $M_{\delta,r}(f) = M_r(f)$ if $\delta = 0$, which is the Hardy-Littlewood maximal function when $r = 1$. For $\beta > 0$ and $p > 1$, let $\dot{F}_p^{\beta,\infty}$ be the homogeneous Triebel-Lizorkin space; the Lipschitz space $\dot{\Lambda}_\beta$ is the space of functions f such that

$$\|f\|_{\dot{\Lambda}_\beta} = \sup_{\substack{x, h \in R^n \\ h \neq 0}} \frac{|\Delta_h^{[\beta]+1} f(x)|}{|h|^\beta} < \infty, \quad (2.2)$$

where Δ_h^k denotes the k th difference operator (see [11]).

We are going to consider the fractional singular integral operator as follows.

DEFINITION 2.1. Let $T : S \rightarrow S'$ be a linear operator. T is called a fractional singular integral operator if there exists a locally integrable function $K(x, y)$ on $R^n \times R^n$ such that

$$T(f)(x) = \int K(x, y)f(y)dy \quad (2.3)$$

for every bounded and compactly supported function f . Let m be a positive integer and A a function on R^n . Denote that

$$R_{m+1}(A; x, y) = A(x) - \sum_{|\alpha| \leq m} \frac{1}{\alpha!} D^\alpha A(y) (x - y)^\alpha. \quad (2.4)$$

The multilinear operator related to fractional singular integral operator T is defined by

$$T_A(f)(x) = \int \frac{R_{m+1}(A; x, y)}{|x - y|^m} K(x, y) f(y) dy. \quad (2.5)$$

Note that when $m = 0$, T_A is just the commutator of T and A while when $m > 0$, it is nontrivial generalizations of the commutators. It is well known that multilinear operators are of great interest in harmonic analysis and have been widely studied by many authors (see [2, 3, 4, 5, 7, 8, 14]). The main purpose of this paper is to study the continuity for the multilinear operator on Triebel-Lizorkin spaces. We will prove the following theorem in Section 3.

THEOREM 2.2. Let $0 < \beta < 1$ and $D^\alpha A \in \dot{\Lambda}_\beta$ for $|\alpha| = m$. Suppose T is the fractional singular integral operator such that T is bounded from $L^p(R^n)$ to $L^q(R^n)$ for $0 \leq \delta < n$, $1 < p < n/\delta$, and $1/p - 1/q = \delta/n$. If T satisfies the size condition

$$|T_A(f)(x) - T_A(f)(x_0)| \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} |Q|^{\beta/n} M_{\delta,1}(f)(x) \quad (2.6)$$

for any cube $Q = Q(x_0, l)$ with $\text{supp } f \subset (2Q)^c$, $x \in Q$, and some $0 \leq \delta < n$, then

- (a) T_A maps $L^p(R^n)$ continuously into $\dot{F}_q^{\beta, \infty}(R^n)$ for $0 \leq \delta < n$, $1 < p < n/\delta$, and $1/p - 1/q = \delta/n$;
- (b) T_A maps $L^p(R^n)$ continuously into $L^q(R^n)$ for $0 \leq \delta < n - \beta$, $1 < p < n/(\delta + \beta)$, and $1/p - 1/q = (\delta + \beta)/n$.

From the theorem, we get the following corollary.

COROLLARY 2.3. Fix $\varepsilon > 0$, $0 < \beta < \min(1, \varepsilon)$, $\delta \geq 0$, and $D^\alpha A \in \dot{\Lambda}_\beta$ for $|\alpha| = m$. Let K be a locally integrable function on $R^n \times R^n$ satisfying

$$\begin{aligned} |K(x, y)| &\leq C|x - y|^{-n+\delta}, \\ |K(y, x) - K(z, x)| &\leq C|y - z|^\varepsilon |x - z|^{-n-\varepsilon+\delta} \end{aligned} \quad (2.7)$$

if $2|y - z| \leq |x - z|$. Denote that (2.3) holds and denote the multilinear operator of T by (2.5) for every bounded and compactly supported function f and $x \in (\text{supp } f)^c$. Suppose T is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$ for $0 \leq \delta < n$, $1 < p < n/\delta$, and $1/q = 1/p - \delta/n$. Then

- (a) T_A maps $L^p(\mathbb{R}^n)$ continuously into $\dot{F}_q^{\beta, \infty}(\mathbb{R}^n)$ for $0 \leq \delta < n$, $1 < p < n/\delta$, and $1/p - 1/q = \delta/n$;
- (b) T_A maps $L^p(\mathbb{R}^n)$ continuously into $L^q(\mathbb{R}^n)$ for $0 \leq \delta < n - \beta$, $1 < p < n/(\delta + \beta)$, and $1/p - 1/q = (\delta + \beta)/n$.

3. Proof of Theorem 2.2. To prove the theorem, we need the following lemmas.

LEMMA 3.1 [11]. For $0 < \beta < 1$ and $1 < p < \infty$,

$$\begin{aligned} \|f\|_{\dot{F}_p^{\beta, \infty}} &\approx \left\| \sup_Q \frac{1}{|Q|^{1+\beta/n}} \int_Q |f(x) - f_Q| dx \right\|_{L^p} \\ &\approx \left\| \sup_{c \in Q} \frac{1}{|Q|^{1+\beta/n}} \int_Q |f(x) - c| dx \right\|_{L^p}. \end{aligned} \quad (3.1)$$

LEMMA 3.2 [11]. For $0 < \beta < 1$ and $1 \leq p \leq \infty$,

$$\begin{aligned} \|f\|_{\dot{\Lambda}_\beta} &\approx \sup_Q \frac{1}{|Q|^{1+\beta/n}} \int_Q |f(x) - f_Q| dx \\ &\approx \sup_Q \frac{1}{|Q|^{\beta/n}} \left(\frac{1}{|Q|} \int_Q |f(x) - f_Q|^p dx \right)^{1/p}. \end{aligned} \quad (3.2)$$

LEMMA 3.3 [1]. Suppose that $1 \leq r < p < n/\delta$ and $1/q = 1/p - \delta/n$. Then $\|M_{\delta, r}(f)\|_{L^q} \leq C\|f\|_{L^p}$.

LEMMA 3.4 [5]. Let A be a function on $\mathbb{R}^n$ and $D^\alpha A \in L^q(\mathbb{R}^n)$ for $|\alpha| = m$ and some $q > n$. Then

$$|R_m(A; x, y)| \leq C|x - y|^m \sum_{|\alpha|=m} \left(\frac{1}{|\tilde{Q}(x, y)|} \int_{\tilde{Q}(x, y)} |D^\alpha A(z)|^q dz \right)^{1/q}, \quad (3.3)$$

where $\tilde{Q}(x, y)$ is the cube centered at x and having side length $5\sqrt{n}|x - y|$.

PROOF OF THEOREM 2.2. Fix a cube $Q = Q(x_0, l)$ and $\tilde{x} \in Q$. Let $\tilde{Q} = 5\sqrt{n}Q$ and $\tilde{A}(x) = A(x) - \sum_{|\alpha|=m} (1/\alpha!)(D^\alpha A)_{\tilde{Q}} x^\alpha$. Then $R_m(A; x, y) = R_m(\tilde{A}; x, y)$ and $D^\alpha \tilde{A} = D^\alpha A - (D^\alpha A)_{\tilde{Q}}$ for $|\alpha| = m$. We write, for $f_1 = f\chi_{\tilde{Q}}$ and $f_2 = f\chi_{\mathbb{R}^n \setminus \tilde{Q}}$,

$$\begin{aligned} T_A(f)(x) &= \int_{\mathbb{R}^n} \frac{R_{m+1}(A; x, y)}{|x - y|^m} K(x, y) f(y) dy \\ &= \int_{\mathbb{R}^n} \frac{R_{m+1}(A; x, y)}{|x - y|^m} K(x, y) f_2(y) dy \\ &\quad + \int_{\mathbb{R}^n} \frac{R_m(\tilde{A}; x, y)}{|x - y|^m} K(x, y) f_1(y) dy \\ &\quad - \sum_{|\alpha|=m} \frac{1}{\alpha!} \int_{\mathbb{R}^n} \frac{K(x, y)(x - y)^\alpha}{|x - y|^m} D^\alpha \tilde{A}(y) f_1(y) dy, \end{aligned} \quad (3.4)$$

then

$$\begin{aligned}
 & |T_A(f)(x) - T_A(f_2)(x_0)| \\
 & \leq \left| T\left(\frac{R_m(\tilde{A}; x, \cdot)}{|x - \cdot|^m} f_1\right)(x) \right| + \sum_{|\alpha|=m} \frac{1}{\alpha!} \left| T\left(\frac{(x - \cdot)^\alpha}{|x - \cdot|^m} D^\alpha \tilde{A} f_1\right)(x) \right| \\
 & + |T_A(f_2)(x) - T_A(f_2)(x_0)| := I(x) + II(x) + III(x).
 \end{aligned} \tag{3.5}$$

Thus,

$$\begin{aligned}
 & \frac{1}{|Q|^{1+\beta/n}} \int_Q |T_A(f)(x) - T_A(f_2)(x_0)| dx \\
 & \leq \frac{1}{|Q|^{1+\beta/n}} \int_Q I(x) dx + \frac{1}{|Q|^{1+\beta/n}} \int_Q II(x) dx + \frac{1}{|Q|^{1+\beta/n}} \int_Q III(x) dx \\
 & := I + II + III.
 \end{aligned} \tag{3.6}$$

Now, we estimate I , II , and III , respectively. First, for $x \in Q$ and $y \in \tilde{Q}$, using Lemmas 3.4 and 3.2, we get

$$\begin{aligned}
 R_m(\tilde{A}; x, y) & \leq C|x - y|^m \sum_{|\alpha|=m} \sup_{x \in \tilde{Q}} |D^\alpha A(x) - (D^\alpha A)_{\tilde{Q}}| \\
 & \leq C|x - y|^m |Q|^{\beta/n} \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta}.
 \end{aligned} \tag{3.7}$$

Thus, taking r, s such that $1 \leq r < p$ and $1/s = 1/r - \delta/n$, by (L^r, L^s) boundedness of T and Hölder's inequality, we obtain

$$\begin{aligned}
 I & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} \frac{1}{|Q|} \int_Q |T(f_1)(x)| dx \\
 & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} \|T(f_1)\|_{L^s} |Q|^{-1/s} \\
 & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} \|f_1\|_{L^r} |Q|^{-1/s} \\
 & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} \left(\frac{1}{|\tilde{Q}|^{1-r\delta/n}} \int_{\tilde{Q}} |f(y)|^r dy \right)^{1/r} \\
 & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} M_{\delta, r}(f)(\tilde{x}).
 \end{aligned} \tag{3.8}$$

Second, using the inequality (see [11])

$$\|(D^\alpha A - (D^\alpha A)_{\tilde{Q}}) f \chi_{\tilde{Q}}\|_{L^r} \leq C|Q|^{1/s+\beta/n} \|D^\alpha A\|_{\dot{\lambda}_\beta} M_{\delta, r}(f)(x), \tag{3.9}$$

we gain

$$\begin{aligned}
 II &\leq \frac{C}{|Q|^{1+\beta/n}} \sum_{|\alpha|=m} \left\| T \left((D^\alpha A - (D^\alpha A)_{\tilde{Q}}) f \chi_{2Q} \right) \right\|_{L^s} |Q|^{1-1/s} \\
 &\leq C |Q|^{-\beta/n-1/s} \sum_{|\alpha|=m} \left\| (D^\alpha A - (D^\alpha A)_{\tilde{Q}}) f \chi_{2Q} \right\|_{L^r} \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} M_{\delta,r}(f)(\tilde{x}).
 \end{aligned} \tag{3.10}$$

For *III*, using the size condition of T , we have

$$III \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} M_{\delta,1}(f)(\tilde{x}). \tag{3.11}$$

We now put these estimates together, and taking the supremum over all Q such that $\tilde{x} \in Q$, and using Lemmas 3.1 and 3.3, we obtain

$$\|T_A(f)\|_{F_q^{\beta,\infty}} \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} \|f\|_{L^p}. \tag{3.12}$$

This completes the proof of (a).

For (b), by the same argument as in the proof of (a), we have

$$\frac{1}{|Q|} \int_Q |T_A(f)(y) - T_A(f_2)(x_0)| dy \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} (M_{\delta+\beta,r}(f) + M_{\delta+\beta,1}(f)). \tag{3.13}$$

Thus, we get the sharp estimate of T_A as follows:

$$(T_A(f))^\# \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} (M_{\delta+\beta,r}(f) + M_{\delta+\beta,1}(f)). \tag{3.14}$$

Now, using Lemma 3.3, we gain

$$\begin{aligned}
 \|T_A(f)\|_{L^q} &\leq C \|(T_A(f))^\#\|_{L^q} \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} (\|M_{\delta+\beta,r}(f)\|_{L^q} + \|M_{\delta+\beta,1}(f)\|_{L^q}) \leq C \|f\|_{L^p}.
 \end{aligned} \tag{3.15}$$

This completes the proof of (b) and the theorem. $\square$

PROOF OF COROLLARY 2.3. It suffices to verify that T satisfies the size condition in Theorem 2.2.

Suppose $\text{supp } f \subset \tilde{Q}^c$ and $x \in Q = Q(x_0, l)$. We write

$$\begin{aligned}
 & T_A(f)(x) - T_A(f)(x_0) \\
 &= \int_{R^n} \left[\frac{K(x, y)}{|x - y|^m} - \frac{K(x_0, y)}{|x_0 - y|^m} \right] R_m(\tilde{A}; x, y) f(y) dy \\
 &+ \int_{R^n} \frac{K(x_0, y) f(y)}{|x_0 - y|^m} [R_m(\tilde{A}; x, y) - R_m(\tilde{A}; x_0, y)] dy \\
 &- \sum_{|\alpha|=m} \frac{1}{\alpha!} \int_{R^n} \left(\frac{K(x, y)(x - y)^\alpha}{|x - y|^m} - \frac{K(x_0, y)(x_0 - y)^\alpha}{|x_0 - y|^m} \right) D^\alpha \tilde{A}(y) f(y) dy \\
 &:= I_1 + I_2 + I_3.
 \end{aligned} \tag{3.16}$$

By Lemma 3.4 and the following inequality, for $b \in \dot{\Lambda}_\beta$:

$$|b(x) - b_Q| \leq \frac{1}{|Q|} \int_Q \|b\|_{\dot{\Lambda}_\beta} |x - y|^\beta dy \leq \|b\|_{\dot{\Lambda}_\beta} (|x - x_0| + l)^\beta, \tag{3.17}$$

we get

$$|R_m(\tilde{A}; x, y)| \leq \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} (|x - y| + l)^{m+\beta}. \tag{3.18}$$

Note that by $|x - y| \sim |x_0 - y|$ for $x \in Q$ and $y \in R^n \setminus \tilde{Q}$, we obtain, by the condition of K ,

$$\begin{aligned}
 |I_1| &\leq C \int_{R^n \setminus \tilde{Q}} \left(\frac{|x - x_0|}{|x_0 - y|^{m+n+1-\delta}} + \frac{|x - x_0|^\varepsilon}{|x_0 - y|^{m+n+\varepsilon-\delta}} \right) |R_m(\tilde{A}; x, y)| |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} \sum_{k=0}^{\infty} \int_{2^{k+1}\tilde{Q} \setminus 2^k\tilde{Q}} \left(\frac{|x - x_0|}{|x_0 - y|^{n+1-\delta-\beta}} + \frac{|x - x_0|^\varepsilon}{|x_0 - y|^{n+\varepsilon-\delta-\beta}} \right) |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} |Q|^{\beta/n} \sum_{k=1}^{\infty} (2^{k(\beta-1)} + 2^{k(\beta-\varepsilon)}) \frac{1}{|2^k\tilde{Q}|^{1-\delta/n}} \int_{2^k\tilde{Q}} |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} |Q|^{\beta/n} M_{\delta,1}(f)(x).
 \end{aligned} \tag{3.19}$$

For I_2 , by the formula (see [3])

$$R_m(\tilde{A}; x, y) - R_m(\tilde{A}; x_0, y) = \sum_{|\eta|<m} \frac{1}{\eta!} R_{m-|\eta|}(D^\eta \tilde{A}; x, x_0) (x - y)^\eta \tag{3.20}$$

and [Lemma 3.4](#), we get

$$\begin{aligned}
 |I_2| &\leq C \int_{R^n \setminus \tilde{Q}} \frac{|R_m(\tilde{A}; x, y) - R_m(\tilde{A}; x_0, y)|}{|x_0 - y|^{m+n-\delta}} |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} \sum_{k=0}^{\infty} \int_{2^{k+1}\tilde{Q} \setminus 2^k\tilde{Q}} \frac{|x - x_0|^{\beta+1}}{|x_0 - y|^{n+1-\delta}} |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} |Q|^{\beta/n} \sum_{k=1}^{\infty} 2^{-k} \frac{1}{|2^k\tilde{Q}|^{1-\delta/n}} \int_{2^k\tilde{Q}} |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} |Q|^{\beta/n} M_{\delta,1}(f)(x).
 \end{aligned} \tag{3.21}$$

For I_3 , similar to the estimates of I_1 , we obtain

$$\begin{aligned}
 |I_3| &\leq C \int_{R^n \setminus \tilde{Q}} \left(\frac{|x - x_0|}{|x_0 - y|^{n+1-\delta}} + \frac{|x - x_0|^\varepsilon}{|x_0 - y|^{n+\varepsilon-\delta}} \right) |D^\alpha A(y) - (D^\alpha A)_{\tilde{Q}}| |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} |Q|^{\beta/n} \sum_{k=1}^{\infty} (2^{k(\beta-1)} + 2^{k(\beta-\varepsilon)}) \frac{1}{|2^k\tilde{Q}|^{1-\delta/n}} \int_{2^k\tilde{Q}} |f(y)| dy \\
 &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\lambda}_\beta} |Q|^{\beta/n} M_{\delta,1}(f)(x).
 \end{aligned} \tag{3.22}$$

Thus (2.6) holds. This completes the proof of the corollary. $\square$

4. Applications. In this section, we will apply [Theorem 2.2](#) and [Corollary 2.3](#) to some particular operators such as the Calderon-Zygmund singular integral operator and fractional integral operator.

APPLICATION 1 (Calderon-Zygmund singular integral operator). Let T be the Calderon-Zygmund operator defined by (2.3) (see [9, 12, 13]); the multilinear operator related to T is defined by

$$T_A f(x) = \int \frac{R_{m+1}(A; x, y)}{|x - y|^m} K(x, y) f(y) dy. \tag{4.1}$$

Then it is easy to see that T satisfies the conditions in [Corollary 2.3](#) with $\delta = 0$; thus T_A is bounded from $L^p(R^n)$ to $\dot{F}_p^{\beta,\infty}(R^n)$ for $D^\alpha A \in \dot{\lambda}_\beta$, $|\alpha| = m$, $0 < \beta < 1$, $1 < p < \infty$ and from $L^p(R^n)$ to $L^q(R^n)$ for $D^\alpha A \in \dot{\lambda}_\beta$, $|\alpha| = m$, $0 < \beta < 1$, $1 < p < n/\beta$, and $1/p - 1/q = \beta/n$.

APPLICATION 2 (fractional integral operator with rough kernel). For $0 \leq \delta < n$, let T_δ be the fractional integral operator with rough kernel defined by (see [7, 8, 14])

$$T_\delta f(x) = \int_{R^n} \frac{\Omega(x - y)}{|x - y|^{n-\delta}} f(y) dy; \tag{4.2}$$

the multilinear operator related to T_δ is defined by

$$T_\delta^A f(x) = \int_{\mathbb{R}^n} \frac{R_{m+1}(A; x, y)}{|x - y|^{m+n-\delta}} \Omega(x - y) f(y) dy, \quad (4.3)$$

where Ω is homogeneous of degree zero on $\mathbb{R}^n$, $\int_{S^{n-1}} \Omega(x') d\sigma(x') = 0$, and $\Omega \in \text{Lip}_\gamma(S^{n-1})$ for $0 < \gamma \leq 1$, that is, there exists a constant $M > 0$ such that for any $x, y \in S^{n-1}$, $|\Omega(x) - \Omega(y)| \leq M|x - y|^\gamma$. Then T_δ satisfies the conditions in [Corollary 2.3](#). In fact, for $\text{supp } f \subset (2Q)^c$ and $x \in Q = Q(x_0, l)$, by the condition of Ω , we have (see [\[14\]](#))

$$\left| \frac{\Omega(x - y)}{|x - y|^{n-\delta}} - \frac{\Omega(x_0 - y)}{|x_0 - y|^{n-\delta}} \right| \leq C \left(\frac{|x - x_0|}{|x_0 - y|^{n+1-\delta}} + \frac{|x - x_0|^\gamma}{|x_0 - y|^{n+\gamma-\delta}} \right); \quad (4.4)$$

thus, similar to the proof of [Corollary 2.3](#), we get

$$\begin{aligned} & |T_\delta^A(f)(x) - T_\delta^A(f)(x_0)| \\ & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} \sum_{k=0}^{\infty} \int_{2^{k+1}\tilde{Q} \setminus 2^k\tilde{Q}} \left(\frac{|x - x_0|}{|x_0 - y|^{n+1-\delta-\beta}} + \frac{|x - x_0|^\gamma}{|x_0 - y|^{n+\gamma-\delta-\beta}} \right) |f(y)| dy \\ & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} |Q|^{\beta/n} \sum_{k=1}^{\infty} (2^{k(\beta-1)} + 2^{k(\beta-\gamma)}) \frac{1}{|2^k\tilde{Q}|^{1-\delta/n}} \int_{2^k\tilde{Q}} |f(y)| dy \\ & \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} |Q|^{\beta/n} M_{\delta,1}(f)(x). \end{aligned} \quad (4.5)$$

Therefore, T_δ^A is bounded from $L^p(\mathbb{R}^n)$ to $\dot{F}_q^{\beta,\infty}(\mathbb{R}^n)$ for $D^\alpha A \in \dot{\Lambda}_\beta$, $|\alpha| = m$, $0 < \beta < \gamma$, $1 < p < n/\beta$ and from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$ for $D^\alpha A \in \dot{\Lambda}_\beta$, $|\alpha| = m$, $0 < \beta < 1$, $1 < p < n/(\delta + \beta)$, and $1/p - 1/q = (\delta + \beta)/n$.

REFERENCES

- [1] S. Chanillo, *A note on commutators*, Indiana Univ. Math. J. **31** (1982), no. 1, 7-16.
- [2] W. Chen, *A Besov estimate for multilinear singular integrals*, Acta Math. Sin. (Engl. Ser.) **16** (2000), no. 4, 613-626.
- [3] J. Cohen, *A sharp estimate for a multilinear singular integral in $\mathbb{R}^n$* , Indiana Univ. Math. J. **30** (1981), no. 5, 693-702.
- [4] J. Cohen and J. Gosselin, *On multilinear singular integrals on $\mathbb{R}^n$* , Studia Math. **72** (1982), no. 3, 199-223.
- [5] ———, *A BMO estimate for multilinear singular integrals*, Illinois J. Math. **30** (1986), no. 3, 445-464.
- [6] R. Coifman, R. Rochberg, and G. Weiss, *Factorization theorems for Hardy spaces in several variables*, Ann. of Math. (2) **103** (1976), no. 3, 611-635.
- [7] Y. Ding, *A note on multilinear fractional integrals with rough kernel*, Adv. Math. (China) **30** (2001), no. 3, 238-246.
- [8] Y. Ding and S. Z. Lu, *Weighted boundedness for a class of rough multilinear operators*, Acta Math. Sin. (Engl. Ser.) **17** (2001), no. 3, 517-526.
- [9] J. García-Cuerva and J. L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Mathematics Studies, vol. 116, North-Holland Publishing, Amsterdam, 1985.

- [10] S. Janson, *Mean oscillation and commutators of singular integral operators*, Ark. Mat. **16** (1978), no. 2, 263–270.
- [11] M. Paluszyński, *Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss*, Indiana Univ. Math. J. **44** (1995), no. 1, 1–17.
- [12] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, New Jersey, 1993.
- [13] A. Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Pure and Applied Mathematics, vol. 123, Academic Press, Florida, 1986.
- [14] P. Zhang, *Some problems related to fractional integrals and the Marcinkiewicz integrals*, Ph.D. thesis, Beijing Normal University, Beijing, 2001.

Liu Lanzhe: College of Mathematics and Computer, Changsha University of Science and Technology, Changsha 410077, China

E-mail address: lanzhe@263.net