UNIT-CIRCLE-PRESERVING MAPPINGS

SOON-MO JUNG and BYUNGBAE KIM

Received 6 April 2004

We prove that if a one-to-one mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($n \geq 2$) preserves the unit circles, then f is a linear isometry up to translation.

2000 Mathematics Subject Classification: 51K05.

1. Introduction. Let X and Y be normed spaces. A mapping $f : X \rightarrow Y$ is called an isometry if f satisfies the equality

$$\|f(x) - f(y)\| = \|x - y\| \quad (1.1)$$

for all $x, y \in X$. A distance $r > 0$ is said to be preserved (conserved) by a mapping $f : X \rightarrow Y$ if

$$\|f(x) - f(y)\| = r \quad \forall x, y \in X \text{ with } \|x - y\| = r. \quad (1.2)$$

If f is an isometry, then every distance $r > 0$ is conserved by f , and vice versa. We can now raise a question whether each mapping that preserves certain distances is an isometry. Indeed, Aleksandrov [1] had raised a question whether a mapping $f : X \rightarrow X$ preserving a distance $r > 0$ is an isometry, which is now known to us as the Aleksandrov problem. Without loss of generality, we may assume $r = 1$ when X is a normed space (see [16]).

Beckman and Quarles [2] solved the Aleksandrov problem for finite-dimensional real Euclidean spaces $X = \mathbb{R}^n$ (see also [3, 4, 5, 6, 7, 8, 11, 12, 13, 14, 15, 17, 18, 19, 20]).

THEOREM 1.1 (Beckman and Quarles). *If a mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($2 \leq n < \infty$) preserves a distance $r > 0$, then f is a linear isometry up to translation.*

Recently, Zaks [25] proved the rational analogues of the Beckman-Quarles theorem. Indeed, he assumes that $n = 4k(k+1)$ for some $k \geq 1$ or $n = 2m^2 - 1$ for some $m \geq 3$, and he proves that if a mapping $f : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$ preserves the unit distance, then f is an isometry (see also [21, 22, 23, 24]).

It seems interesting to investigate whether the “distance $r > 0$ ” in the Beckman-Quarles theorem can be replaced by some properties characterized by “geometrical figures” without loss of its validity.

In [9], the first author proved that if a one-to-one mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($n \geq 2$) maps every regular triangle (quadrilateral or hexagon) of side length $a > 0$ onto a figure of

FIGURE 2.1

the same type with side length $b > 0$, then there exists a linear isometry $I: \mathbb{R}^n \rightarrow \mathbb{R}^n$ up to translation such that

$$f(x) = \frac{b}{a}I(x). \quad (1.3)$$

Furthermore, the first author proved that if a one-to-one mapping $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ maps every unit circle onto a unit circle, then f is a linear isometry up to translation (see [10]).

In this connection, we will extend the result of [10] to the n -dimensional cases; more precisely, we prove in this paper that if a one-to-one mapping $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($n \geq 2$) maps every unit circle onto a unit circle, then f is a linear isometry up to translation.

2. Preliminaries. We start with any two distinct points a and b in $\mathbb{R}^n$ with the distance between the two less than 2. Let their distance be

$$2c = 2\sin\varphi_0 \quad \text{with } 0 < \varphi_0 < \frac{\pi}{2}, \quad 0 < c < 1. \quad (2.1)$$

Given such two distinct points whose distance is less than 2, we can choose a coordinate $(y_1, \dots, y_n)$ for $\mathbb{R}^n$ such that

$$a = (0, \dots, 0, \sin\varphi_0), \quad b = (0, \dots, 0, -\sin\varphi_0). \quad (2.2)$$

Let the $(n-2)$ -dimensional unit sphere contained in the space orthogonal to the y_n -direction be

$$Y = \{(y_1, \dots, y_{n-1}, 0) \mid y_1^2 + \dots + y_{n-1}^2 = 1\}. \quad (2.3)$$

If we call the center of any unit circle passing through the two points (a and b) o' and the origin of the coordinate o , then the vector $\overrightarrow{oo'}$ is perpendicular to the y_n -axis and its length must be $\cos\varphi_0$ and therefore $\overrightarrow{oo'} \in \tilde{Y} = \cos\varphi_0 Y$, see Figure 2.1. It means that any unit circle passing through the points a and b has its center in $\tilde{Y} = \cos\varphi_0 Y$. Let T be the set of union of all the unit circles passing through the points a and b . More precisely, if we define the following set:

$$T = \{(\cos\varphi + \cos\varphi_0)y + (0, \dots, 0, \sin\varphi) \mid y \in Y, \quad 0 \leq \varphi < 2\pi\}, \quad (2.4)$$

then it is clear that this is the set of union of all the unit circles which are centered at $\cos \varphi_0 \gamma$ for each fixed $\gamma \in Y$ and which pass through a and b when $\varphi = \pi \mp \varphi_0$ (see Figure 2.1).

The intersection of T and the $\gamma_1\text{-}\gamma_n$ plane consists of two circles, say C_1 (when $\gamma_1 = 1$, i.e., $\gamma = (1, 0, \dots, 0)$) and C_2 (when $\gamma_1 = -1$, i.e., $\gamma = (-1, 0, \dots, 0)$, see Figure 2.1). In the following contexts, we will consider the cases $\gamma_1 = 1$ and -1 in connection with T as the circles C_1 and C_2 , respectively. Call S_1 the $(n-1)$ -dimensional unit sphere containing the circle C_1 . If we let the center of C_1 be O and the center of S_1 be $\tilde{O}$, then it is obvious that $O = \tilde{O}$.

(To see this, choose any point $A \in C_1$ and its antipodal point B in C_1 . Then, by the definition of the antipodal points that they lie exactly the opposite with respect to the center of the circle C_1 whose center is at O , and because they are of the same length 1, we have the following condition that

$$\overrightarrow{OA} = -\overrightarrow{OB}, \quad \overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB} = 2\overrightarrow{OB}. \quad (2.5)$$

On the other hand, we have, since the two points A and B lie also on the unit sphere S_1 with its center at $\tilde{O}$,

$$2 = |\overrightarrow{AB}| = |\overrightarrow{A\tilde{O}} + \overrightarrow{\tilde{O}B}| \leq |\overrightarrow{A\tilde{O}}| + |\overrightarrow{\tilde{O}B}| = 1 + 1 = 2. \quad (2.6)$$

Therefore, by the Cauchy-Schwarz inequality, $\overrightarrow{A\tilde{O}}$ is a positive multiple of $\overrightarrow{\tilde{O}B}$, which means $\overrightarrow{A\tilde{O}} = \overrightarrow{\tilde{O}B}$ because their lengths are both 1. So,

$$\overrightarrow{AB} = \overrightarrow{A\tilde{O}} + \overrightarrow{\tilde{O}B} = 2\overrightarrow{\tilde{O}B}, \quad (2.7)$$

and therefore $\tilde{O} = O$.)

Now, we first show that S_1 and T intersect only at C_1 . To make computation simpler we use a new coordinate x for $\mathbb{R}^n$, where

$$x = \gamma - (\cos \varphi_0, 0, \dots, 0). \quad (2.8)$$

In this coordinate (see Figure 2.2), S_1 becomes the unit sphere S centered at the origin,

$$S_1 = S = \{(x_1, \dots, x_n) \mid x_1^2 + \dots + x_n^2 = 1\}, \quad (2.9)$$

$$T = \{x = (\cos \varphi + \cos \varphi_0)\gamma + (0, \dots, 0, \sin \varphi) - (\cos \varphi_0, 0, \dots, 0) \mid \gamma \in Y, 0 \leq \varphi < 2\pi\}. \quad (2.10)$$

With the help of this coordinate we show the following lemma.

LEMMA 2.1. $T \cap S_1 = C_1$.

FIGURE 2.2

PROOF. If any element in T has distance 1 from the origin of the x -coordinate, then we have

$$\begin{aligned}
 1 &= [(\cos \varphi + \cos \varphi_0)y_1 - \cos \varphi_0]^2 + (\cos \varphi + \cos \varphi_0)^2 y_2^2 \\
 &\quad + \cdots + (\cos \varphi + \cos \varphi_0)^2 y_{n-1}^2 + \sin^2 \varphi \\
 &= (\cos \varphi + \cos \varphi_0)^2 - 2 \cos \varphi_0 (\cos \varphi + \cos \varphi_0) y_1 + \cos^2 \varphi_0 + \sin^2 \varphi \\
 &= 1 + 2 \cos^2 \varphi_0 (1 - y_1) + 2 \cos \varphi_0 \cos \varphi (1 - y_1).
 \end{aligned} \tag{2.11}$$

Therefore, we have

$$0 = 2 \cos \varphi_0 (1 - y_1) (\cos \varphi + \cos \varphi_0). \tag{2.12}$$

With $y_1 = 1$, T in (2.10) represents the unit circle C_1 in the x_1 - x_n plane. If

$$\cos \varphi = -\cos \varphi_0, \quad \text{i.e., } \varphi = \pi \mp \varphi_0, \tag{2.13}$$

then it follows from (2.10) that

$$T = \{x = (-\cos \varphi_0, 0, \dots, 0, \pm \sin \varphi_0)\} = \{a, b\} \tag{2.14}$$

which also belong to C_1 . □

Now, consider, as in Figure 2.3, the origin e and $\tilde{e} = (-2, 0, \dots, 0)$ in the x -coordinate and the unit circle C_1 passing through e and $\tilde{e}$ in the x_1 - x_n plane. Choose a point $d \in C_1$, $d \notin \{e, \tilde{e}\}$. We parameterize all the unit circles passing through the points e and d . We assume the x_n -coordinate of d is negative.

By triangle inequality, the distance between e and d is less than 2, say $2 \sin \varphi_0$, with $0 < \varphi_0 < \pi/2$. Choose a new coordinate y for $\mathbb{R}^n$ and consider two points

$$e' = (0, \dots, 0, \sin \varphi_0), \quad d' = (0, \dots, 0, -\sin \varphi_0), \tag{2.15}$$

(see Figure 2.4).

FIGURE 2.3

FIGURE 2.4

To get a parameterization of the unit circles passing through e and d , we consider the mapping M defined by

$$x = My = \begin{bmatrix} \cos \varphi_0 & 0 & \cdots & 0 & \sin \varphi_0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \\ -\sin \varphi_0 & 0 & \cdots & 0 & \cos \varphi_0 \end{bmatrix} [y + (\cos \varphi_0, 0, \dots, 0)]^t - (1, 0, \dots, 0)^t. \quad (2.16)$$

This transformation M is an isometry (since it is a composition of a rotation and translations) and sends

$$\{y = (0, \dots, 0, \pm \sin \varphi_0)\} = \{e', d'\} \quad (2.17)$$

to

$$\{x = (0, \dots, 0), x = (\cos(-2\varphi_0) - 1, 0, \dots, 0, \sin(-2\varphi_0))\} = \{e, d\} \quad (2.18)$$

and therefore it sends any unit circle passing through e' and d' to a unit circle passing through e and d .

Therefore, by comparing Figure 2.4 with Figure 2.1 and considering (2.4), all the unit circles passing through e and d can be parameterized as

$$\{x = M\gamma \mid \gamma = (\cos \varphi + \cos \varphi_0)\gamma' + (0, \dots, 0, \sin \varphi), \gamma' \in Y, 0 \leq \varphi < 2\pi\}. \quad (2.19)$$

With the help of this parameterization, we are ready to show the following lemma.

LEMMA 2.2. *For $d \in C_1$, $d \notin \{e, \tilde{e}\}$, any unit circle in $\mathbb{R}^n$, which passes through d and e, has some point whose x_1 -coordinate is positive, except the circle C_1 .*

PROOF. Without loss of generality, we can assume the x_n -coordinate of d is negative. Note that with $\varphi = \pi \mp \varphi_0$ in (2.19), $\gamma = (0, \dots, 0, \pm \sin \varphi_0)$ are the points e' or d' in the γ -coordinate and further $\varphi = \pi \mp \varphi_0$ means that

$$x = (0, \dots, 0) = e, \quad x = (\cos(-2\varphi_0) - 1, 0, \dots, 0, \sin(-2\varphi_0)) = d \quad (2.20)$$

in the x -coordinate, regardless of $\gamma' \in Y$. Any unit circle passing through e and d is given as $x = M\gamma$ with γ given as in (2.19), that is,

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \\ x_n \end{bmatrix} = \begin{bmatrix} \cos \varphi_0 & 0 & \cdots & 0 & \sin \varphi_0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \\ -\sin \varphi_0 & 0 & \cdots & 0 & \cos \varphi_0 \end{bmatrix} \begin{bmatrix} (\cos \varphi + \cos \varphi_0)\gamma'_1 + \cos \varphi_0 \\ (\cos \varphi + \cos \varphi_0)\gamma'_2 \\ \vdots \\ (\cos \varphi + \cos \varphi_0)\gamma'_{n-1} \\ \sin \varphi \end{bmatrix} - \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}. \quad (2.21)$$

The first coordinate is

$$x_1 = \cos \varphi_0 (\cos \varphi + \cos \varphi_0) \gamma'_1 + \cos^2 \varphi_0 + \sin \varphi_0 \sin \varphi - 1. \quad (2.22)$$

We show that for $\gamma'_1 \neq -1$ ($\gamma'_1 = -1$ means the circle C'_1 in the γ -coordinate and the circle C_1 in the x -coordinate, see Figure 2.4), there is always some φ near $\pi - \varphi_0$ (i.e., near the point e) such that the above x_1 becomes positive.

Let

$$\theta = (\pi - \varphi_0) - \varphi = \pi - (\varphi + \varphi_0), \quad (2.23)$$

and so

$$\varphi = \pi - (\theta + \varphi_0). \quad (2.24)$$

Then, the above is

$$\begin{aligned}
 x_1 &= -\cos \varphi_0 \cos(\theta + \varphi_0) y'_1 + \cos^2 \varphi_0 (1 + y'_1) + \sin \varphi_0 \sin(\theta + \varphi_0) - 1 \\
 &= -\cos \varphi_0 [\cos \theta \cos \varphi_0 - \sin \theta \sin \varphi_0] y'_1 + \sin \varphi_0 [\sin \theta \cos \varphi_0 + \cos \theta \sin \varphi_0] \\
 &\quad - 1 + \cos^2 \varphi_0 (1 + y'_1) \\
 &= \sin \theta \sin \varphi_0 \cos \varphi_0 (1 + y'_1) + \cos \theta \sin^2 \varphi_0 - \cos \theta \cos^2 \varphi_0 y'_1 \\
 &\quad - 1 + \cos^2 \varphi_0 (1 + y'_1) \\
 &= \sin \theta \sin \varphi_0 \cos \varphi_0 (1 + y'_1) + \cos \theta - \cos \theta \cos^2 \varphi_0 (1 + y'_1) \\
 &\quad - [1 - \cos^2 \varphi_0 (1 + y'_1)] \\
 &= \sin \theta \sin \varphi_0 \cos \varphi_0 (1 + y'_1) - [1 - \cos^2 \varphi_0 (1 + y'_1)] (1 - \cos \theta).
 \end{aligned} \tag{2.25}$$

$\theta = 0$ ($\varphi = \pi - \varphi_0$) means the intersection point e and the above x_1 becomes 0 as it should. Assume

$$\theta \neq 0 \quad (-\pi - \varphi_0 < \theta < 0, \quad 0 < \theta \leq \pi - \varphi_0). \tag{2.26}$$

Then, x_1 is positive if and only if

$$\sin \theta \sin \varphi_0 \cos \varphi_0 (1 + y'_1) > [1 - \cos^2 \varphi_0 (1 + y'_1)] (1 - \cos \theta), \tag{2.27}$$

that is,

$$\frac{\sin \theta}{1 - \cos \theta} > \frac{1 - \cos^2 \varphi_0 (1 + y'_1)}{\sin \varphi_0 \cos \varphi_0 (1 + y'_1)} \tag{2.28}$$

(recall $y'_1 \neq -1$ and $0 < \varphi_0 < \pi/2$). In other words, the x_1 -coordinate is positive if and only if

$$\cot \frac{\theta}{2} > \frac{1 - \cos^2 \varphi_0 (1 + y'_1)}{\sin \varphi_0 \cos \varphi_0 (1 + y'_1)}. \tag{2.29}$$

Therefore, for $y'_1 \neq -1$ (i.e., except the circle C_1), the x_1 -coordinate is positive for small enough $\theta > 0$. $\square$

3. Main theorem. In the previous section, we introduced all preliminary lemmas for the main result of this paper. Now, we prove our main theorem.

THEOREM 3.1. *If a one-to-one mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ maps every unit circle onto a unit circle, then f is a linear isometry up to translation.*

PROOF. We show f preserves the distance 2. Suppose the distance between $a = f(A)$ and $b = f(B)$ is less than 2, while the distance between A and B is 2—see [Figure 3.1](#). Then, we show it leads to a contradiction.

Let the distance between a and b be $2c$ ($0 < c < 1$). Choose any unit circle C passing through A and B and let $f(C) = C_1$. Choose a coordinate for a and b as in [Figure 3.1](#) such that C_1 lies in the x_1 - x_n plane and

$$a = (-1 - \sqrt{1 - c^2}, 0, \dots, 0, c), \quad b = (-1 - \sqrt{1 - c^2}, 0, \dots, 0, -c). \tag{3.1}$$

FIGURE 3.1

Let

$$e = (0, \dots, 0), \quad \tilde{e} = (-2, 0, \dots, 0). \quad (3.2)$$

Let $f(E) = e$ and $\tilde{E}$ the antipodal point (in C) of E and let $f(\tilde{E}) = d$. Let the union of all the unit circles passing through a and b be T and the $(n-1)$ -dimensional unit sphere passing through A and B be S and the $(n-1)$ -dimensional unit sphere passing through e and $\tilde{e}$ be S_1 .

Then, it is clear that any point P on S ($P \notin \{A, B\}$) lies in some unit circle determined by the three points A , B , and P . To see this, if we call O the common center of C and S , and let

$$\langle \overrightarrow{OP}, \overrightarrow{OA} \rangle = \sin \varphi_0 \quad \left(-\frac{\pi}{2} < \varphi_0 < \frac{\pi}{2} \right), \quad (3.3)$$

then the unit circle determined by these three points is parameterized as

$$\overrightarrow{OV}(\varphi) = \cos \varphi \left(\frac{\overrightarrow{OP} - \sin \varphi_0 \overrightarrow{OA}}{\cos \varphi_0} \right) + \sin \varphi \overrightarrow{OA} \quad (-\pi < \varphi \leq \pi). \quad (3.4)$$

Note that

$$\left\{ \left(\frac{\overrightarrow{OP} - \sin \varphi_0 \overrightarrow{OA}}{\cos \varphi_0} \right), \overrightarrow{OA} \right\} \quad (3.5)$$

are orthonormal to each other and

$$\begin{aligned} \overrightarrow{OV}(\varphi_0) &= \overrightarrow{OP}, & \overrightarrow{OV}\left(\frac{\pi}{2}\right) &= \overrightarrow{OA}, \\ \overrightarrow{OV}\left(-\frac{\pi}{2}\right) &= -\overrightarrow{OA} = \overrightarrow{OB}. \end{aligned} \quad (3.6)$$

Since the image of this unit circle lies in T , it follows that the image of the whole S under f lies in T .

It is also obvious that the x_1 -coordinate of any point in T is nonpositive. (Note that the center of any unit circle passing through a and b has coordinate

$$\sqrt{1-c^2}y - (1 + \sqrt{1-c^2}, 0, \dots, 0) \quad \text{for some } y \in Y, \quad (3.7)$$

(see (2.4)) and the distance between this center and any $x = (x_1, \dots, x_n)$ is

$$\sqrt{(x_1 + 1 + \sqrt{1-c^2}(1-y_1))^2 + \dots} \quad (3.8)$$

and because

$$\sqrt{1-c^2}(1-y_1) \geq 0, \quad (3.9)$$

positive x_1 makes the distance larger than 1, which means that if $x_1 > 0$, we have $x \notin T$.)

Now, if $d = \tilde{e}$, then the image of any unit circle passing through E and $\tilde{E}$ lies in both T and S_1 . However, by Lemma 2.1, $T \cap S_1 = C_1$ and this fact contradicts the injectivity of f .

On the other hand, if $d \neq \tilde{e}$, the image of any unit circle, except the circle C , passing through E and $\tilde{E}$ is a unit circle passing through e and d . This unit circle is not C_1 since f is one-to-one, and by Lemma 2.2 it cannot stay completely in T , a contradiction.

Consequently, f preserves the distance 2. According to the well-known theorem of Beckman and Quarles, f is a linear isometry up to translation. $\square$

ACKNOWLEDGMENT. This work was supported by Korea Research Foundation Grant KRF-2003-015-C00023.

REFERENCES

- [1] A. D. Aleksandrov, *Mappings of families of sets*, Soviet Math. Dokl. **11** (1970), 116–120.
- [2] F. S. Beckman and D. A. Quarles Jr., *On isometries of Euclidean spaces*, Proc. Amer. Math. Soc. **4** (1953), 810–815.
- [3] W. Benz, *Isometrien in normierten Räumen*, Aequationes Math. **29** (1985), no. 2-3, 204–209 (German).
- [4] ———, *An elementary proof of the theorem of Beckman and Quarles*, Elem. Math. **42** (1987), no. 1, 4–9.
- [5] W. Benz and H. Berens, *A contribution to a theorem of Ulam and Mazur*, Aequationes Math. **34** (1987), no. 1, 61–63.
- [6] R. L. Bishop, *Characterizing motions by unit distance invariance*, Math. Mag. **46** (1973), 148–151.
- [7] K. Cieśliński and T. M. Rassias, *On some properties of isometric mappings*, Facta Univ. Ser. Math. Inform. (1992), no. 7, 107–115.
- [8] D. Greenwell and P. D. Johnson, *Functions that preserve unit distance*, Math. Mag. **49** (1976), no. 2, 74–79.
- [9] S.-M. Jung, *Mappings preserving some geometrical figures*, Acta Math. Hungar. **100** (2003), no. 1-2, 167–175.
- [10] ———, *Mappings preserving unit circles in $\mathbb{R}^2$* , Octagon Math. Mag. **11** (2003), 450–453.
- [11] B. Mielnik and T. M. Rassias, *On the Aleksandrov problem of conservative distances*, Proc. Amer. Math. Soc. **116** (1992), no. 4, 1115–1118.
- [12] P. S. Modenov and A. S. Parkhomenko, *Geometric Transformations. Vol. 1: Euclidean and Affine Transformations*, Academic Press, New York, 1965.

- [13] T. M. Rassias, *Is a distance one preserving mapping between metric spaces always an isometry?* Amer. Math. Monthly **90** (1983), 200.
- [14] ———, *Some remarks on isometric mappings*, Facta Univ. Ser. Math. Inform. (1987), no. 2, 49–52.
- [15] ———, *Mappings that preserve unit distance*, Indian J. Math. **32** (1990), no. 3, 275–278.
- [16] ———, *Properties of isometries and approximate isometries*, Recent Progress in Inequalities (Niš, 1996) (G. V. Milovanovic, ed.), Math. Appl., vol. 430, Kluwer Academic Publishers, Dordrecht, 1998, pp. 341–379.
- [17] T. M. Rassias and P. Šemrl, *On the Mazur-Ulam theorem and the Aleksandrov problem for unit distance preserving mappings*, Proc. Amer. Math. Soc. **118** (1993), no. 3, 919–925.
- [18] T. M. Rassias and C. S. Sharma, *Properties of isometries*, J. Nat. Geom. **3** (1993), no. 1, 1–38.
- [19] E. M. Schröder, *Eine Ergänzung zum Satz von Beckman and Quarles*, Aequationes Math. **19** (1979), no. 1, 89–92 (German).
- [20] C. G. Townsend, *Congruence-preserving mappings*, Math. Mag. **43** (1970), 37–38.
- [21] A. Tyszk, *A discrete form of the Beckman-Quarles theorem*, Amer. Math. Monthly **104** (1997), no. 8, 757–761.
- [22] ———, *Discrete versions of the Beckman-Quarles theorem*, Aequationes Math. **59** (2000), no. 1-2, 124–133.
- [23] ———, *A discrete form of the Beckman-Quarles theorem for rational eight-space*, Aequationes Math. **62** (2001), no. 1-2, 85–93.
- [24] J. Zaks, *A discrete form of the Beckman-Quarles theorem for rational spaces*, J. Geom. **72** (2001), no. 1-2, 199–205.
- [25] ———, *The Beckman-Quarles theorem for rational spaces*, Discrete Math. **265** (2003), no. 1–3, 311–320.

Soon-Mo Jung: Mathematics Section, College of Science and Technology, Hong-Ik University,
339-701 Chochiwon, Korea

E-mail address: smjung@wow.hongik.ac.kr

Byungbae Kim: Mathematics Section, College of Science and Technology, Hong-Ik University,
339-701 Chochiwon, Korea

E-mail address: bkim@wow.hongik.ac.kr