INEQUALITY FOR RICCI CURVATURE OF CERTAIN SUBMANIFOLDS IN LOCALLY CONFORMAL ALMOST COSYMPLECTIC MANIFOLDS

DAE WON YOON

Received 23 June 2004

We establish inequalities between the Ricci curvature and the squared mean curvature, and also between the k -Ricci curvature and the scalar curvature for a slant, semi-slant, and bi-slant submanifold in a locally conformal almost cosymplectic manifold with arbitrary codimension.

1. Preliminaries

Let $\widetilde{M}$ be a $(2m+1)$ -dimensional almost contact manifold with almost contact structure (φ, ξ, η) , that is, a global vector field ξ , a $(1, 1)$ tensor field φ , and a 1-form η on $\widetilde{M}$ such that $\varphi^2 X = -X + \eta(X)\xi$, $\eta(\xi) = 1$ for any vector field X on $\widetilde{M}$. We consider a product manifold $\widetilde{M} \times \mathbb{R}$, where $\mathbb{R}$ denotes a real line. Then a vector field on $\widetilde{M} \times \mathbb{R}$ is given by $(X, f(d/dt))$, where X is a vector field tangent to $\widetilde{M}$, t the coordinate of $\mathbb{R}$, and f a function on $\widetilde{M} \times \mathbb{R}$. We define a linear map J on the tangent space of $\widetilde{M} \times \mathbb{R}$ by $J(X, f(d/dt)) = (\varphi X - f\xi, \eta(X)(d/dt))$. Then we have $J^2 = -I$, and hence J is an almost complex structure on $\widetilde{M} \times \mathbb{R}$. The manifold $\widetilde{M}$ is said to be *normal* (see [6]) if the almost complex structure J is integrable (i.e., J arises from a complex structure on $\widetilde{M} \times \mathbb{R}$). Let g be a Riemannian metric on $\widetilde{M}$ compatible with (φ, ξ, η) , that is, $g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y)$ for any vector fields X and Y tangent to $\widetilde{M}$. Thus, the manifold $\widetilde{M}$ is almost contact metric, and (φ, ξ, η, g) is its almost contact metric structure. Clearly, we have $\eta(X) = g(X, \xi)$ for any vector field X tangent to $\widetilde{M}$. Let Φ denote the fundamental 2-form of $\widetilde{M}$ defined by $\Phi(X, Y) = g(\varphi X, Y)$ for any vector fields X and Y tangent to $\widetilde{M}$. The manifold $\widetilde{M}$ is said to be *almost cosymplectic* if the forms η and Φ are closed. That is, $d\eta = 0$ and $d\Phi = 0$, where d is the operator of exterior differentiation. If $\widetilde{M}$ is almost cosymplectic and normal, then it is called *cosymplectic* (see [1]). It is well known that the almost contact metric manifold is cosymplectic if and only if $\tilde{\nabla}\varphi$ vanishes identically, where $\tilde{\nabla}$ is the Levi-Civita connection on $\widetilde{M}$. An almost contact metric manifold $\widetilde{M}$ is a locally conformal almost cosymplectic manifold if and only if there exists a 1-form ω such that $d\Phi = 2\omega \wedge \Phi$, $d\eta = \omega \wedge \eta$, and $d\omega = 0$.

On the other hand, it is wellknown that the Riemannian curvature tensor $\tilde{R}$ on a locally conformal almost cosymplectic manifold $\widetilde{M}$ ($m \geq 2$) of pointwise constant φ -sectional

curvature c satisfies (see[6])

$$\begin{aligned}
 & g(\tilde{R}(X, Y)Z, W) \\
 &= \frac{c - 3f^2}{4} \{g(X, W)g(Y, Z) - g(X, Z)g(Y, W)\} \\
 &+ \frac{c + f^2}{4} \{g(X, \varphi W)g(Y, \varphi Z) - g(X, \varphi Z)g(Y, \varphi W) - 2g(X, \varphi Y)g(Z, \varphi W)\} \\
 &- \left(\frac{c + f^2}{4} + f' \right) \{g(X, W)\eta(Y)\eta(Z) - g(X, Z)\eta(Y)\eta(W) + g(Y, Z)\eta(X)\eta(W) \\
 &\quad - g(Y, W)\eta(X)\eta(Z)\}, \quad X, Y, Z, W \in T_p M,
 \end{aligned} \tag{1.1}$$

where f is the function such that $\omega = f\eta$, $f' = \xi f$.

In [5], Lotta has introduced the following notion of slant submanifolds into almost contact metric manifolds. A submanifold M tangent to ξ in locally conformal almost cosymplectic manifold $\tilde{M}$ is said to be *slant* if for any $p \in M$ and any $X \in T_p M$, linearly independent of ξ , the angle between φX and $T_p M$ is a constant $\theta \in [0, \pi/2]$, called the *slant angle* of M in $\tilde{M}$. Invariant and anti-invariant submanifolds of $\tilde{M}$ are slant submanifolds with slant angles $\theta = 0$ and $\theta = \pi/2$, respectively.

We say that a submanifold M tangent to ξ is a *bi-slant* submanifold in $\tilde{M}$ if there exist two orthogonal distributions $\mathcal{D}_1$ and $\mathcal{D}_2$ on M such that

- (1) TM admits the orthogonal direct decomposition $TM = \mathcal{D}_1 \oplus \mathcal{D}_2 \oplus \{\xi\}$;
- (2) for any $i = 1, 2$, $\mathcal{D}_i$ is slant distribution with slant angle θ_i .

On the other hand, CR -submanifolds of $\tilde{M}$ are bi-slant submanifolds with $\theta_1 = 0$, $\theta_2 = \pi/2$.

Let $2d_1 = \dim \mathcal{D}_1$ and $2d_2 = \dim \mathcal{D}_2$.

Remark 1.1. If either d_1 or d_2 vanishes, the bi-slant submanifold is a slant submanifold. Thus, slant submanifolds are particular cases of bi-slant submanifolds.

A submanifold M tangent to ξ is called a *semi-slant* submanifold in $\tilde{M}$ if there exist two orthogonal distributions $\mathcal{D}_1$ and $\mathcal{D}_2$ on M such that

- (1) TM admits the orthogonal direct decomposition $TM = \mathcal{D}_1 \oplus \mathcal{D}_2 \oplus \{\xi\}$;
- (2) the distribution $\mathcal{D}_1$ is an invariant distribution, that is, $\varphi(\mathcal{D}_1) = \mathcal{D}_1$;
- (3) the distribution $\mathcal{D}_2$ is slant with angle $\theta \neq 0$.

Remark 1.2. The invariant distribution of a semi-slant submanifold is a slant distribution with zero angle. Thus, it is obvious that, in fact, semi-slant submanifolds are particular cases of bi-slant submanifolds.

- (1) If $d_2 = 0$, then M is an invariant submanifold.
- (2) If $d_1 = 0$ and $\theta = \pi/2$, then M is an anti-invariant submanifold.

For the other properties and examples of slant, bi-slant, and semi-slant submanifolds in an almost contact metric manifold, we refer to [2, 3].

Let M be an n -dimensional submanifold of a locally conformal almost cosymplectic manifold $\tilde{M}$ equipped with a Riemannian metric g . The Gauss and Weingarten formulas

are given, respectively, by

$$\tilde{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad \tilde{\nabla}_X N = -A_N X + \nabla_X^\perp N, \quad (1.2)$$

for all $X, Y \in TM$ and $N \in T^\perp M$, where $\tilde{\nabla}$, ∇ , and $\nabla^\perp$ are the Riemannian, induced Riemannian, and induced normal connections in $\tilde{M}$, M , and the normal bundle $T^\perp M$ of M , respectively, and h is the second fundamental form related to the shape operator A by $g(h(X, Y), N) = g(A_N X, Y)$. Also, let R be the Riemannian curvature tensor of M . Then the equation of Gauss is given by

$$\tilde{R}(X, Y, Z, W) = R(X, Y, Z, W) + g(h(X, W), h(Y, Z)) - g(h(X, Z), h(Y, W)), \quad (1.3)$$

for any vectors X, Y, Z, W tangent to M .

For any vector X tangent to M , we put $\varphi X = PX + FX$, where PX and FX are the tangential and the normal components of φX , respectively. Given an orthonormal basis $\{e_1, \dots, e_n\}$ of M , we define the squared norm of P by

$$\|P\|^2 = \sum_{i,j=1}^n g^2(Pe_i, e_j) \quad (1.4)$$

and the mean curvature vector $H(p)$ at $p \in M$ is given by $H = (1/n) \sum_{i=1}^n h(e_i, e_i)$.

We put

$$h_{ij}^r = g(h(e_i, e_j), e_r), \quad \|h\|^2 = \sum_{i,j=1}^n g(h(e_i, e_j), h(e_i, e_j)), \quad (1.5)$$

where $\{e_{n+1}, \dots, e_{2m+1}\}$ is an orthonormal basis of $T_p^\perp M$ and $r = n+1, \dots, 2m+1$. A submanifold M in $\tilde{M}$ is called *totally geodesic* if the second fundamental form vanishes identically and *totally umbilical* if there is a real number λ such that $h(X, Y) = \lambda g(X, Y)H$ for any tangent vectors X, Y on M .

For an n -dimensional Riemannian manifold M , we denote by $K(\pi)$ the sectional curvature of M associated with a plane section $\pi \subset T_p M$, $p \in M$. For an orthonormal basis $\{e_1, \dots, e_n\}$ of the tangent space $T_p M$, the scalar curvature τ is defined by

$$\tau = \sum_{i < j} K_{ij}, \quad (1.6)$$

where K_{ij} denotes the sectional curvature of the 2-plane section spanned by e_i and e_j .

Suppose that L is a k -plane section of $T_p M$ and X a unit vector in L . We choose an orthonormal basis $\{e_1, \dots, e_k\}$ of L such that $e_1 = X$. Define the Ricci curvature Ric_L of L at X by

$$\text{Ric}_L(X) = K_{12} + \dots + K_{1k}. \quad (1.7)$$

We simply called such a curvature a *k-Ricci curvature*. The scalar curvature τ of the k -plane section L is given by

$$\tau(L) = \sum_{1 \leq i < j \leq k} K_{ij}. \quad (1.8)$$

For each integer k , $2 \leq k \leq n$, the Riemannian invariant Θ_k on an n -dimensional Riemannian manifold M is defined by

$$\Theta_k(p) = \frac{1}{k-1} \inf_{L, X} \text{Ric}_L(X), \quad p \in M, \quad (1.9)$$

where L runs over all k -plane sections in $T_p M$ and X runs over all unit vectors in L .

Recall that for a submanifold M in a Riemannian manifold, the relative null space of M at a point $p \in M$ is defined by

$$N_p = \{X \in T_p M \mid h(X, Y) = 0 \ \forall Y \in T_p M\}. \quad (1.10)$$

2. Ricci curvature and squared mean curvature

Chen established a sharp relationship between the Ricci curvature and the squared mean curvature for submanifolds in real space forms (see [4]). We prove similar inequalities for slant, bi-slant, and semi-slant submanifolds in a locally conformal almost cosymplectic manifold $\widetilde{M}$. We consider submanifolds M tangent to ξ .

THEOREM 2.1. *Let M be an n -dimensional θ -slant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, the following hold.*

(1) *For each unit vector $X \in T_p M$ orthogonal to ξ ,*

$$\text{Ric}(X) \leq \frac{1}{4} \left\{ (n-1)(c-3f^2) + \frac{3}{2}(c+f^2) \cos^2 \theta - 4 \left(\frac{c+f^2}{4} + f' \right) + n^2 \|H\|^2 \right\}. \quad (2.1)$$

(2) *If $H(p) = 0$, then a unit tangent vector X orthogonal to ξ at p satisfies the equality case of (2.1) if and only if $X \in N_p$.*

(3) *The equality case of (2.1) holds identically for all unit tangent vectors orthogonal to ξ at p if and only if p is a totally geodesic point.*

Proof. (1) Let $X \in T_p M$ be a unit tangent vector at p orthogonal to ξ . We choose an orthonormal basis $e_1, \dots, e_n = \xi, e_{n+1}, \dots, e_{2m+1}$, such that $e_1, \dots, e_n$ are tangent to M at p with $e_1 = X$. Then, from the equation of Gauss, we have

$$\begin{aligned} n^2 \|H\|^2 &= 2\tau + \|h\|^2 - \frac{n(n-1)(c-3f^2)}{4} \\ &\quad - \frac{3(n-1)(c+f^2)}{4} \cos^2 \theta + 2(n-1) \left(\frac{c+f^2}{4} + f' \right). \end{aligned} \quad (2.2)$$

From (2.2), we get

$$\begin{aligned}
 n^2 \|H\|^2 &= 2\tau + \sum_{r=n+1}^{2m+1} \left[(h_{11}^r)^2 + (h_{22}^r + \cdots + h_{nn}^r)^2 + 2 \sum_{1 \leq i < j \leq n} (h_{ij}^r)^2 \right] \\
 &\quad - 2 \sum_{r=n+1}^{2m+1} \sum_{2 \leq i < j \leq n} h_{ii}^r h_{jj}^r - \frac{n(n-1)(c-3f^2)}{4} \\
 &\quad - \frac{3(n-1)(c+f^2)}{4} \cos^2 \theta + 2(n-1) \left(\frac{c+f^2}{4} + f' \right) \\
 &= 2\tau + \frac{1}{2} \sum_{r=n+1}^{2m+1} \left[(h_{11}^r + h_{22}^r + \cdots + h_{nn}^r)^2 + (h_{11}^r - h_{22}^r - \cdots - h_{nn}^r)^2 \right] \\
 &\quad + 2 \sum_{r=n+1}^{2m+1} \sum_{1 \leq i < j \leq n} (h_{ij}^r)^2 - 2 \sum_{r=n+1}^{2m+1} \sum_{2 \leq i < j \leq n} h_{ii}^r h_{jj}^r \\
 &\quad - \frac{n(n-1)(c-3f^2)}{4} - \frac{3(n-1)(c+f^2)}{4} \cos^2 \theta + 2(n-1) \left(\frac{c+f^2}{4} + f' \right). \tag{2.3}
 \end{aligned}$$

By using the equation of Gauss, we have

$$\begin{aligned}
 \sum_{2 \leq i < j \leq n} K_{ij} &= \sum_{r=n+1}^{2m+1} \sum_{2 \leq i < j \leq n} \left[h_{ii}^r h_{jj}^r - (h_{ij}^r)^2 \right] + \frac{(n-1)(n-2)(c-3f^2)}{8} \\
 &\quad + \frac{3(n-2)(c+f^2)}{8} \cos^2 \theta + \frac{1}{2} \left(\frac{c+f^2}{4} + f' \right) (-2n+4). \tag{2.4}
 \end{aligned}$$

Substituting (2.4) in (2.3), we get

$$\frac{1}{2} n^2 \|H\|^2 \geq 2 \operatorname{Ric}(X) - \frac{(n-1)(c-3f^2)}{2} - \frac{3(c+f^2)}{4} \cos^2 \theta + 2 \left(\frac{c+f^2}{4} + f' \right), \tag{2.5}$$

or equivalently (2.1).

(2) Assume that $H(P) = 0$. Equality holds in (2.1) if and only if

$$\begin{aligned}
 h_{12}^r &= \cdots = h_{1n}^r = 0, \\
 h_{11}^r &= h_{22}^r + \cdots + h_{nn}^r, \quad r \in \{n+1, \dots, 2m+1\}. \tag{2.6}
 \end{aligned}$$

Then $h_{1j}^r = 0$ for all $j \in \{1, \dots, n\}$, $r \in \{n+1, \dots, 2m+1\}$, that is, $X \in N_p$.

(3) Then equality case of (2.1) holds for all unit tangent vectors orthogonal to ξ at p if and only if

$$\begin{aligned}
 h_{ij}^r &= 0, \quad i \neq j, r \in \{n+1, \dots, 2m+1\}, \\
 h_{11}^r + \cdots + h_{nn}^r - 2h_{ii}^r &= 0, \quad i \in \{1, \dots, n\}, r \in \{n+1, \dots, 2m+1\}. \tag{2.7}
 \end{aligned}$$

In this case, it follows that p is a totally geodesic point. The converse is trivial. $\square$

THEOREM 2.2. Let M be an n -dimensional bi-slant submanifold satisfying $g(X, \varphi Y) = 0$, for any $X \in \mathcal{D}_1$ and any $Y \in \mathcal{D}_2$, tangent to ξ in a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, the following hold.

(1) For each unit vector $X \in T_p M$ orthogonal to ξ and if

(i) X is tangent to $\mathcal{D}_1$,

$$\text{Ric}(X) \leq \frac{1}{4} \left\{ (n-1)(c-3f^2) + \frac{3}{2}(c+f^2) \cos^2 \theta_1 - 4 \left(\frac{c+f^2}{4} + f' \right) + n^2 \|H\|^2 \right\}, \quad (2.8)$$

and if

(ii) X is tangent to $\mathcal{D}_2$,

$$\text{Ric}(X) \leq \frac{1}{4} \left\{ (n-1)(c-3f^2) + \frac{3}{2}(c+f^2) \cos^2 \theta_2 - 4 \left(\frac{c+f^2}{4} + f' \right) + n^2 \|H\|^2 \right\}. \quad (2.9)$$

(2) If $H(p) = 0$, then a unit tangent vector X orthogonal to ξ at p satisfies the equality case of (2.8) and (2.9) if and only if $X \in N_p$.

(3) The equality case of (2.8) and (2.9) holds identically for all unit tangent vectors orthogonal to ξ at p if and only if p is a totally geodesic point.

Proof. (1) Let $X \in T_p M$ be a unit tangent vector at p orthogonal to ξ . We choose an orthonormal basis $e_1, \dots, e_n = \xi, e_{n+1}, \dots, e_{2m+1}$ such that $e_1, \dots, e_n$ are tangent to M at p with $e_1 = X$. Then, from the equation of Gauss, we have

$$\begin{aligned} n^2 \|H\|^2 &= 2\tau + \|h\|^2 - \frac{n(n-1)(c-3f^2)}{4} \\ &\quad - \frac{6(c+f^2)}{4} (d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2) + 2(n-1) \left(\frac{c+f^2}{4} + f' \right), \end{aligned} \quad (2.10)$$

where $2d_1 = \dim \mathcal{D}_1$ and $2d_2 = \dim \mathcal{D}_2$.

From (2.10), we get

$$\begin{aligned} n^2 \|H\|^2 &= 2\tau + \sum_{r=n+1}^{2m+1} \left[(h_{11}^r)^2 + (h_{22}^r + \dots + h_{nn}^r)^2 + 2 \sum_{1 \leq i < j \leq n} (h_{ij}^r)^2 \right] \\ &\quad - 2 \sum_{r=n+1}^{2m+1} \sum_{2 \leq i < j \leq n} h_{ii}^r h_{jj}^r - \frac{n(n-1)(c-3f^2)}{4} \\ &\quad - \frac{6(c+f^2)}{4} (d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2) + 2(n-1) \left(\frac{c+f^2}{4} + f' \right) \end{aligned}$$

$$\begin{aligned}
&= 2\tau + \frac{1}{2} \sum_{r=n+1}^{2m+1} \left[(h_{11}^r + h_{22}^r + \cdots + h_{nn}^r)^2 + (h_{11}^r - h_{22}^r - \cdots - h_{nn}^r)^2 \right] \\
&\quad + 2 \sum_{r=n+1}^{2m+1} \sum_{1 \leq i < j \leq n} (h_{ij}^r)^2 - 2 \sum_{r=n+1}^{2m+1} \sum_{2 \leq i < j \leq n} h_{ii}^r h_{jj}^r - \frac{n(n-1)(c-3f^2)}{4} \\
&\quad - \frac{6(c+f^2)}{4} (d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2) + 2(n-1) \left(\frac{c+f^2}{4} + f' \right).
\end{aligned} \tag{2.11}$$

We distinguish two cases.

(i) If X is tangent to $\mathcal{D}_1$, then we have

$$\begin{aligned}
\sum_{2 \leq i < j \leq n} K_{ij} &= \sum_{r=n+1}^{2m+1} \sum_{2 \leq i < j \leq n} \left[h_{ii}^r h_{jj}^r - (h_{ij}^r)^2 \right] + \frac{(n-1)(n-2)(c-3f^2)}{8} \\
&\quad + \frac{c+f^2}{8} [6(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2) - 3 \cos^2 \theta_1] + \frac{1}{2} \left(\frac{c+f^2}{4} + f' \right) (-2n+4).
\end{aligned} \tag{2.12}$$

Substituting (2.12) in (2.11), one gets

$$\frac{1}{2} n^2 \|H\|^2 \geq 2 \operatorname{Ric}(X) - \frac{(n-1)(c-3f^2)}{2} - \frac{3(c+f^2)}{4} \cos^2 \theta_1 + 2 \left(\frac{c+f^2}{4} + f' \right), \tag{2.13}$$

which is equivalent to (2.8).

(ii) If X is tangent to $\mathcal{D}_2$, then we have

$$\begin{aligned}
\sum_{2 \leq i < j \leq n} K_{ij} &= \sum_{r=n+1}^{2m+1} \sum_{2 \leq i < j \leq n} \left[h_{ii}^r h_{jj}^r - (h_{ij}^r)^2 \right] + \frac{(n-1)(n-2)(c-3f^2)}{8} \\
&\quad + \frac{c+f^2}{8} [6(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2) - 3 \cos^2 \theta_2] + \frac{1}{2} \left(\frac{c+f^2}{4} + f' \right) (-2n+4).
\end{aligned} \tag{2.14}$$

Substituting (2.14) in (2.11), one gets

$$\frac{1}{2} n^2 \|H\|^2 \geq 2 \operatorname{Ric}(X) - \frac{(n-1)(c-3f^2)}{2} - \frac{3(c+f^2)}{4} \cos^2 \theta_2 + 2 \left(\frac{c+f^2}{4} + f' \right), \tag{2.15}$$

which is equivalent to (2.9).

(2) Assume that $H(p) = 0$. Equality holds in (2.8) and (2.9) if and only if

$$\begin{aligned}
h_{12}^r &= \cdots = h_{1n}^r = 0, \\
h_{11}^r &= h_{22}^r + \cdots + h_{nn}^r, \quad r \in \{n+1, \dots, 2m+1\}.
\end{aligned} \tag{2.16}$$

Then $h_{ij}^r = 0$ for all $j \in \{1, \dots, n\}$, $r \in \{n+1, \dots, 2m+1\}$, that is, $X \in N_p$.

(3) Then equality case of (2.8) and (2.9) holds for all unit tangent vectors orthogonal to ξ at p if and only if

$$\begin{aligned} h_{ij}^r &= 0, \quad i \neq j, \quad r \in \{n+1, \dots, 2m+1\}, \\ h_{11}^r + \dots + h_{nn}^r - 2h_{ii}^r &= 0, \quad i \in \{1, \dots, n\}, \quad r \in \{n+1, \dots, 2m+1\}. \end{aligned} \quad (2.17)$$

In this case, it follows that p is a totally geodesic point. The converse is trivial. $\square$

COROLLARY 2.3. *Let M be an n -dimensional semi-slant submanifold in a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\tilde{M}$. Then, the following hold.*

(1) *For each unit vector $X \in T_p M$ orthogonal to ξ and if*

(i) *X is tangent to $\mathcal{D}_1$,*

$$\text{Ric}(X) \leq \frac{1}{4} \left\{ (n-1)(c-3f^2) - 4 \left(\frac{c+f^2}{4} + f' \right) + n^2 \|H\|^2 \right\}, \quad (2.18)$$

and if

(ii) *X is tangent to $\mathcal{D}_2$,*

$$\text{Ric}(X) \leq \frac{1}{4} \left\{ (n-1)(c-3f^2) + \frac{3}{2}(c+f^2) \cos^2 \theta - 4 \left(\frac{c+f^2}{4} + f' \right) + n^2 \|H\|^2 \right\}. \quad (2.19)$$

(2) *If $H(p) = 0$, then a unit tangent vector X orthogonal to ξ at p satisfies the equality case of (2.18) and (2.19) if and only if $X \in N_p$.*

(3) *The equality case of (2.18) and (2.19) holds identically for all unit tangent vectors orthogonal to ξ at p if and only if p is a totally geodesic point.*

COROLLARY 2.4. *Let M be an n -dimensional invariant submanifold in a $(2m+1)$ -dimensional cosymplectic space form $\tilde{M}(c)$. Then, the following hold.*

(1) *For each unit vector $X \in T_p M$ orthogonal to ξ ,*

$$\text{Ric}(X) \leq \frac{1}{4} \left\{ (n-1)(c-3f^2) + \frac{3}{2}(c+f^2) - 4 \left(\frac{c+f^2}{4} + f' \right) + n^2 \|H\|^2 \right\}. \quad (2.20)$$

(2) *If $H(p) = 0$, then a unit tangent vector X orthogonal to ξ at p satisfies the equality case of (2.20) if and only if $X \in N_p$.*

(3) *The equality case of (2.20) holds identically for all unit tangent vectors orthogonal to ξ at p if and only if p is a totally geodesic point.*

COROLLARY 2.5. *Let M be an n -dimensional anti-invariant submanifold in a $(2m+1)$ -dimensional cosymplectic space form $\tilde{M}(c)$. Then, the following hold.*

(1) *For each unit vector $X \in T_p M$ orthogonal to ξ ,*

$$\text{Ric}(X) \leq \frac{1}{4} \left\{ (n-1)(c-3f^2) - 4 \left(\frac{c+f^2}{4} + f' \right) + n^2 \|H\|^2 \right\}. \quad (2.21)$$

(2) *If $H(p) = 0$, then a unit tangent vector X orthogonal to ξ at p satisfies the equality case of (2.21) if and only if $X \in N_p$.*

- (3) The equality case of (2.21) holds identically for all unit tangent vectors orthogonal to ξ at p if and only if p is a totally geodesic point.

3. k -Ricci curvature and squared mean curvature

In this section, we prove relationship between the k -Ricci curvature and the squared mean curvature for slant, bi-slant, and semi-slant submanifolds in a locally conformal almost cosymplectic manifold $\widetilde{M}$. We state an inequality between the scalar curvature and the squared mean curvature for submanifolds M tangent to the vector field ξ .

THEOREM 3.1. *Let M be an n -dimensional θ -slant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then,*

$$\|H\|^2 \geq \frac{2\tau}{n(n-1)} - \frac{1}{4n} \left[n(c-3f^2) + 3(c+f^2)\cos^2\theta - 8\left(\frac{c+f^2}{4} + f'\right) \right], \quad (3.1)$$

equality holding at a point $p \in M$ if and only if p is a totally umbilical point.

Proof. Let p be a point of M . We choose an orthonormal basis $\{e_1, e_2, \dots, e_n = \xi\}$ for the tangent space $T_p M$ and $\{e_{n+1}, \dots, e_{2m+1}\}$ for the normal space $T_p^\perp M$ at p such that the normal vector e_{n+1} is in the direction of the mean curvature vector and $e_1, e_2, \dots, e_n$ diagonalize the shape operator A_{n+1} . Then, we have

$$A_{n+1} = \begin{pmatrix} a_1 & 0 & 0 & \dots & 0 \\ 0 & a_2 & 0 & \dots & 0 \\ 0 & 0 & a_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & a_n \end{pmatrix}, \quad (3.2)$$

$$A_r = (h_{ij}^r), \quad \sum_{i=1}^n h_{ii}^r = 0, \quad n+2 \leq r \leq 2m+1.$$

From the equation of Gauss,

$$\begin{aligned} n^2 \|H\|^2 &= 2\tau + \sum_{i=1}^n a_i^2 + \sum_{r=n+2}^{2m+1} \sum_{i,j=1}^n (h_{ij}^r)^2 - \frac{n(n-1)(c-3f^2)}{4} \\ &\quad - \frac{3(n-1)(c+f^2)}{4} \cos^2\theta + 2(n-1) \left(\frac{c+f^2}{4} + f' \right). \end{aligned} \quad (3.3)$$

On the other hand,

$$\sum_{i < j} (a_i - a_j)^2 = (n-1) \sum_{i=1}^n a_i^2 - 2 \sum_{i < j} a_i a_j. \quad (3.4)$$

Therefore, from the above equation, we have

$$n^2 \|H\|^2 = \left(\sum_{i=1}^n a_i \right)^2 = \sum_{i=1}^n a_i^2 + 2 \sum_{i < j} a_i a_j \leq n \sum_{i=1}^n a_i^2. \quad (3.5)$$

Combining (3.3) and (3.5),

$$\begin{aligned} n(n-1) \|H\|^2 \geq 2\tau + \sum_{r=n+2}^{2m+1} \sum_{i,j=1}^n (h_{ij}^r)^2 - \frac{n(n-1)(c-3f^2)}{4} \\ - \frac{3(n-1)(c+f^2)}{4} \cos^2 \theta + 2(n-1) \left(\frac{c+f^2}{4} + f' \right), \end{aligned} \quad (3.6)$$

which implies inequality (3.1). If the equality sign of (3.1) holds at a point $p \in M$, then from (3.4) and (3.6) we get $A_r = 0$ ($r = n+2, \dots, 2m+1$) and $a_1 = \dots = a_n$. Consequently, p is a totally umbilical point. The converse is trivial. $\square$

THEOREM 3.2. *Let M be an n -dimensional bi-slant submanifold satisfying $g(X, \phi Y) = 0$, for any $X \in \mathcal{D}_1$ and any $Y \in \mathcal{D}_2$, tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then,*

$$\begin{aligned} \|H\|^2 \geq \frac{2\tau}{n(n-1)} - \frac{1}{4n(n-1)} \left[n(n-1)(c-3f^2) + 6(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2)(c+f^2) \right. \\ \left. - 8(n-1) \left(\frac{c+f^2}{4} + f' \right) \right], \end{aligned} \quad (3.7)$$

where $2d_1 = \dim \mathcal{D}_1$ and $2d_2 = \dim \mathcal{D}_2$.

THEOREM 3.3. *Let M be an n -dimensional semi-slant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then,*

$$\begin{aligned} \|H\|^2 \geq \frac{2\tau}{n(n-1)} - \frac{1}{4n(n-1)} \left[n(n-1)(c-3f^2) + 6(d_1 + d_2 \cos^2 \theta)(c+f^2) \right. \\ \left. - 8(n-1) \left(\frac{c+f^2}{4} + f' \right) \right], \end{aligned} \quad (3.8)$$

where $2d_1 = \dim \mathcal{D}_1$ and $2d_2 = \dim \mathcal{D}_2$.

THEOREM 3.4. *Let M be an n -dimensional θ -slant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, for any integer k ($2 \leq k \leq n$) and any point $p \in M$,*

$$\|H\|^2 \geq \Theta_k(p) - \frac{1}{4n} \left[n(c-3f^2) + 3(c+f^2) \cos^2 \theta - 8 \left(\frac{c+f^2}{4} + f' \right) \right]. \quad (3.9)$$

Proof. Let $\{e_1, \dots, e_n\}$ be an orthonormal basis of $T_p M$. Denote by $L_{i_1 \dots i_k}$ the k -plane section spanned by $e_{i_1}, \dots, e_{i_k}$. It follows from (1.7) and (1.8) that

$$\begin{aligned}\tau(L_{i_1 \dots i_k}) &= \frac{1}{2} \sum_{i \in \{i_1, \dots, i_k\}} \text{Ric}_{L_{i_1 \dots i_k}}(e_i), \\ \tau(p) &= \frac{1}{\binom{n-2}{k-2}} \sum_{1 \leq i_1 < \dots < i_k \leq n} \tau(L_{i_1 \dots i_k}).\end{aligned}\quad (3.10)$$

Combining (1.9) and (3.10), we obtain

$$\tau(p) \geq \frac{n(n-1)}{2} \Theta_k(p). \quad (3.11)$$

Therefore, by using (3.1) and (3.11), we can obtain the inequality in Theorem 3.4. $\square$

THEOREM 3.5. *Let M be an n -dimensional bi-slant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, for any integer k ($2 \leq k \leq n$) and any point $p \in M$,*

$$\begin{aligned}\|H\|^2 \geq \Theta_k(p) - \frac{1}{4n(n-1)} &\left[n(n-1)(c-3f^2) + 6(d_1 \cos^2 \theta_1 + d_2 \cos^2 \theta_2)(c+f^2) \right. \\ &\left. - 8(n-1) \left(\frac{c+f^2}{4} + f' \right) \right],\end{aligned}\quad (3.12)$$

where $2d_1 = \dim \mathcal{D}_1$ and $2d_2 = \dim \mathcal{D}_2$.

THEOREM 3.6. *Let M be an n -dimensional semi-slant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, for any integer k ($2 \leq k \leq n$) and any point $p \in M$,*

$$\begin{aligned}\|H\|^2 \geq \Theta_k(p) - \frac{1}{4n(n-1)} &\left[n(n-1)(c-3f^2) + 6(d_1 + d_2 \cos^2 \theta)(c+f^2) \right. \\ &\left. - 8(n-1) \left(\frac{c+f^2}{4} + f' \right) \right],\end{aligned}\quad (3.13)$$

where $2d_1 = \dim \mathcal{D}_1$ and $2d_2 = \dim \mathcal{D}_2$.

COROLLARY 3.7. *Let M be an n -dimensional invariant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, for any integer k ($2 \leq k \leq n$) and any point $p \in M$,*

$$\|H\|^2 \geq \Theta_k(p) - \frac{1}{4n} \left[n(c-3f^2) + 3(c+f^2) - 8 \left(\frac{c+f^2}{4} + f' \right) \right]. \quad (3.14)$$

COROLLARY 3.8. *Let M be an n -dimensional anti-invariant submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, for any integer k ($2 \leq k \leq n$) and any point $p \in M$,*

$$\|H\|^2 \geq \Theta_k(p) - \frac{1}{4n} \left[n(c - 3f^2) - 8 \left(\frac{c+f^2}{4} + f' \right) \right]. \quad (3.15)$$

COROLLARY 3.9. *Let M be an n -dimensional contact CR-submanifold tangent to ξ into a $(2m+1)$ -dimensional locally conformal almost cosymplectic manifold $\widetilde{M}$. Then, for any integer k ($2 \leq k \leq n$) and any point $p \in M$,*

$$\|H\|^2 \geq \Theta_k(p) - \frac{1}{4n(n-1)} \left[n(n-1)(c - 3f^2) + 6d_1(c + f^2) - 8(n-1) \left(\frac{c+f^2}{4} + f' \right) \right]. \quad (3.16)$$

References

- [1] D. E. Blair, *Contact Manifolds in Riemannian Geometry*, Lecture Notes in Mathematics, vol. 509, Springer, Berlin, 1976.
- [2] J. L. Cabrerizo, A. Carriazo, L. M. Fernández, and M. Fernández, *Semi-slant submanifolds of a Sasakian manifold*, *Geom. Dedicata* **78** (1999), no. 2, 183–199.
- [3] ———, *Slant submanifolds in Sasakian manifolds*, *Glasg. Math. J.* **42** (2000), no. 1, 125–138.
- [4] B.-Y. Chen, *Relations between Ricci curvature and shape operator for submanifolds with arbitrary codimensions*, *Glasg. Math. J.* **41** (1999), no. 1, 33–41.
- [5] A. Lotta, *Slant submanifolds in contact geometry*, *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)* **39** (1996), no. 1-4, 183–198.
- [6] Z. Olszak, *Locally conformal almost cosymplectic manifolds*, *Colloq. Math.* **57** (1989), no. 1, 73–87.

Dae Won Yoon: Department of Mathematics Education and Research Institute of Natural Science (RINS), Gyeongsang National University, 900 Gazwa-dong, Jinju 660-701, South Korea
E-mail address: dwyoon@gsnu.ac.kr