ON ABSOLUTE MATRIX SUMMABILITY METHODS

H. S. ÖZARSLAN AND H. N. ÖĞDÜK

Received 27 October 2004 and in revised form 1 August 2005

We have proved a theorem on $|T, p_n|_k$ summability methods. This theorem includes a known theorem.

1. Introduction

Let $\sum a_n$ be a given infinite series with partial sums (s_n) . By (w_n^δ) , we denote the n th Cesàro means of order $\delta (\delta > -1)$ of the sequence (s_n) . The series $\sum a_n$ is said to be summable $|C, \delta|_k$, $k \geq 1$, if (see [3])

$$\sum_{n=1}^{\infty} n^{k-1} |w_n^\delta - w_{n-1}^\delta|^k < \infty. \quad (1.1)$$

In the special case for $\delta = 1$, $|C, \delta|_k$ summability reduces to $|C, 1|_k$ summability.

Let (p_n) be a sequence of positive numbers such that

$$P_n = \sum_{v=0}^n p_v \longrightarrow \infty \quad \text{as } n \longrightarrow \infty, \quad (P_{-i} = p_{-i} = 0, i \geq 1). \quad (1.2)$$

The sequence-to-sequence transformation

$$\vartheta_n = \frac{1}{P_n} \sum_{v=0}^n p_v s_v \quad (1.3)$$

defines the sequence (ϑ_n) of the $(\tilde{N}, p_n)$ means of the sequence (s_n) , generated by the sequence of coefficients (p_n) (see [4]). The series $\sum a_n$ is said to be summable $|\tilde{N}, p_n|_k$, $k \geq 1$, if (see [1])

$$\sum_{n=1}^{\infty} \left(\frac{P_n}{p_n} \right)^{k-1} |\vartheta_n - \vartheta_{n-1}|^k < \infty. \quad (1.4)$$

If we take $p_n = 1$ for all values of n , then $|\tilde{N}, p_n|_k$ summability is the same as $|C, 1|_k$ summability.

Given a normal matrix $T = (t_{nk})$, we associate two lower semimatrices $\bar{T} = (\bar{t}_{nk})$ and $\hat{T} = (\hat{t}_{nk})$ as follows:

$$\begin{aligned} \bar{t}_{nk} &= \sum_{i=k}^n t_{ni}, \quad n, k = 0, 1, \dots, \\ \hat{t}_{00} &= \bar{t}_{00} = t_{00}, \quad \hat{t}_{nk} = \bar{t}_{nk} - \bar{t}_{n-1,k}, \quad n = 1, 2, \dots \end{aligned} \quad (1.5)$$

It may be noted that $\bar{T}$ and $\hat{T}$ are the well-known matrices of series-to-sequence and series-to-series transformations, respectively. Then, we have

$$\begin{aligned} T_n(s) &= \sum_{v=0}^n t_{nv} s_v = \sum_{v=0}^n \bar{t}_{nv} a_v, \\ \bar{\Delta} T_n(s) &= \sum_{v=0}^n \hat{t}_{nv} a_v. \end{aligned} \quad (1.6)$$

The series $\sum a_n$ is said to be summable $|T, p_n|_k$, $k \geq 1$, if (see [5])

$$\sum_{n=1}^{\infty} \left(\frac{P_n}{p_n} \right)^{k-1} |\bar{\Delta} T_n(s)|^k < \infty. \quad (1.7)$$

In the special case, for $t_{nv} = p_v/P_n$, $|T, p_n|_k$ summability is the same as $|\bar{N}, p_n|_k$ summability.

2. The main result

The object of this paper is to prove the following theorem.

THEOREM 2.1. *Let $k \geq 1$. Let (s_n) be a bounded sequence and suppose that (λ_n) is a sequence such that*

$$\begin{aligned} \sum_{n=0}^m \left(\frac{P_n}{p_n} \right)^{k-1} |\lambda_n|^k |t_{nn}|^k &= O(1) \quad \text{as } m \rightarrow \infty, \\ \sum_{n=0}^m |\Delta \lambda_n| &= O(1) \quad \text{as } m \rightarrow \infty. \end{aligned} \quad (2.1)$$

If

$$\frac{1}{|t_{nn}|} \sum_{v=0}^{n-1} |\Delta_v(\hat{t}_{nv})| = O(1) \quad \text{as } n \rightarrow \infty, \quad (2.2)$$

$$\sum_{n=v+1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} |\Delta_v \hat{t}_{nv}| |t_{nn}|^{k-1} = O \left(\left(\frac{P_v}{p_v} \right)^{k-1} |t_{vv}|^k \right) \quad \text{as } m \rightarrow \infty, \quad (2.3)$$

$$\frac{1}{|t_{nn}|} \sum_{v=0}^{n-1} |\Delta\lambda_v| |\hat{t}_{n,v+1}| = O(1) \quad \text{as } n \rightarrow \infty, \quad (2.4)$$

$$\sum_{n=v+1}^{m+1} \left(\frac{P_n}{p_n}\right)^{k-1} |\hat{t}_{n,v+1}| |t_{nn}|^{k-1} = O(1) \quad \text{as } m \rightarrow \infty, \quad (2.5)$$

then the series $\sum a_n \lambda_n$ is summable $|T, p_n|_k$.

Proof. Let (y_n) be the T -transform of the series $\sum a_n \lambda_n$. Then we have, by (1.6),

$$Y_n = y_n - y_{n-1} = \sum_{v=0}^n \hat{t}_{nv} a_v \lambda_v. \quad (2.6)$$

Since $\hat{t}_{nn} = t_{nn}$, by Abel's transformation, we get that

$$\begin{aligned} Y_n &= \sum_{v=0}^{n-1} \Delta_v (\hat{t}_{nv} \lambda_v) s_v + \hat{t}_{nn} \lambda_n s_n \\ &= \sum_{v=0}^{n-1} \Delta \lambda_v \hat{t}_{n,v+1} s_v + \sum_{v=0}^{n-1} \lambda_v \Delta_v (\hat{t}_{nv}) s_v + s_n t_{nn} \lambda_n \\ &= Y_n(1) + Y_n(2) + Y_n(3). \end{aligned} \quad (2.7)$$

Using Minkowski's inequality, it is sufficient to show that

$$\sum_{n=1}^{\infty} \left(\frac{P_n}{p_n}\right)^{k-1} |Y_n(r)|^k < \infty \quad \text{for } r = 1, 2, 3. \quad (2.8)$$

Since (s_n) is bounded, when $k > 1$, applying Hölder's inequality with indices k and k' , where $1/k + 1/k' = 1$, we have that

$$\begin{aligned} \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n}\right)^{k-1} |Y_n(1)|^k &\leq \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n}\right)^{k-1} \left\{ \sum_{v=0}^{n-1} |\Delta \lambda_v| |\hat{t}_{n,v+1}| |s_v| \right\}^k \\ &= O(1) \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n}\right)^{k-1} \sum_{v=0}^{n-1} |\Delta \lambda_v| |\hat{t}_{n,v+1}| |t_{nn}|^{k-1} \\ &\quad \times \left\{ \frac{1}{|t_{nn}|} \sum_{v=0}^{n-1} |\Delta \lambda_v| |\hat{t}_{n,v+1}| \right\}^{k-1} \\ &= O(1) \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n}\right)^{k-1} \sum_{v=0}^{n-1} |\Delta \lambda_v| |\hat{t}_{n,v+1}| |t_{nn}|^{k-1} \\ &= O(1) \sum_{v=0}^m |\Delta \lambda_v| \sum_{n=v+1}^{m+1} \left(\frac{P_n}{p_n}\right)^{k-1} |\hat{t}_{n,v+1}| |t_{nn}|^{k-1} \\ &= O(1) \sum_{v=0}^m |\Delta \lambda_v| = O(1) \quad \text{as } m \rightarrow \infty, \end{aligned} \quad (2.9)$$

by virtue of the hypothesis of Theorem 2.1.

Again using Hölder's inequality, we have

$$\begin{aligned}
 \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} |Y_n(2)|^k &\leq \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} \left\{ \sum_{v=0}^{n-1} |\lambda_v| |\Delta_v \hat{t}_{nv}| |s_v| \right\}^k \\
 &= O(1) \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} \sum_{v=0}^{n-1} |\lambda_v|^k |\Delta_v \hat{t}_{nv}| |t_{nn}|^{k-1} \\
 &\quad \times \left\{ \frac{1}{|t_{nn}|} \sum_{v=0}^{n-1} |\Delta_v \hat{t}_{nv}| \right\}^{k-1} \\
 &= O(1) \sum_{n=1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} \sum_{v=0}^{n-1} |\lambda_v|^k |\Delta_v \hat{t}_{nv}| |t_{nn}|^{k-1} \\
 &= O(1) \sum_{v=0}^m |\lambda_v|^k \sum_{n=v+1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} |\Delta_v \hat{t}_{nv}| |t_{nn}|^{k-1} \\
 &= O(1) \sum_{v=0}^m \left(\frac{P_v}{p_v} \right)^{k-1} |\lambda_v|^k |t_{vv}|^k = O(1) \quad \text{as } m \rightarrow \infty,
 \end{aligned} \tag{2.10}$$

by virtue of the hypothesis of Theorem 2.1.

Finally, we have that

$$\sum_{n=1}^m \left(\frac{P_n}{p_n} \right)^{k-1} |Y_n(3)|^k = O(1) \sum_{n=1}^m \left(\frac{P_n}{p_n} \right)^{k-1} |t_{nn}|^k |\lambda_n|^k = O(1) \quad \text{as } m \rightarrow \infty, \tag{2.11}$$

by virtue of the hypothesis of Theorem 2.1.

Therefore, we get that

$$\sum_{n=1}^m \left(\frac{P_n}{p_n} \right)^{k-1} |Y_n(r)|^k = O(1) \quad \text{as } m \rightarrow \infty, \text{ for } r = 1, 2, 3. \tag{2.12}$$

This completes the proof of Theorem 2.1. $\square$

3. An application

Now we will prove the following corollary.

COROLLARY 3.1 (see [2]). *Let $k \geq 1$. If the sequence (s_n) is bounded and (λ_n) is a sequence such that*

$$\begin{aligned}
 \sum_{n=1}^m \frac{P_n}{p_n} |\lambda_n|^k &= O(1) \quad \text{as } m \rightarrow \infty, \\
 \sum_{n=1}^m |\Delta \lambda_n| &= O(1) \quad \text{as } m \rightarrow \infty,
 \end{aligned} \tag{3.1}$$

then the series $\sum a_n \lambda_n$ is summable $|\tilde{N}, p_n|_k$.

Proof. In Theorem 2.1, let $t_{nv} = p_v/P_n$. Then to prove the corollary, it is sufficient to show that the conditions of Theorem 2.1 are satisfied.

If $t_{nn} = p_n/P_n$, (2.1) are automatically satisfied.

Since

$$\begin{aligned}
 \Delta_v \hat{t}_{nv} &= \hat{t}_{nv} - \hat{t}_{n,v+1} \\
 &= \bar{t}_{nv} - \bar{t}_{n-1,v} - \bar{t}_{n,v+1} + \bar{t}_{n-1,v+1} \\
 &= \sum_{i=v}^n t_{ni} - \sum_{i=v}^{n-1} t_{n-1,i} - \sum_{i=v+1}^n t_{ni} + \sum_{i=v+1}^{n-1} t_{n-1,i} \\
 &= \frac{1}{P_n} \sum_{i=v}^n p_i - \frac{1}{P_{n-1}} \sum_{i=v}^{n-1} p_i - \frac{1}{P_n} \sum_{i=v+1}^n p_i + \frac{1}{P_{n-1}} \sum_{i=v+1}^{n-1} p_i \\
 &= -\frac{p_n p_v}{P_n P_{n-1}},
 \end{aligned} \tag{3.2}$$

we get

$$\frac{1}{|t_{nn}|} \sum_{v=0}^{n-1} |\Delta_v \hat{t}_{nv}| = \frac{P_n}{P_n} \sum_{v=0}^{n-1} \frac{p_n p_v}{P_n P_{n-1}} = O(1) \quad \text{as } n \rightarrow \infty. \tag{3.3}$$

Thus condition (2.2) is satisfied.

Using $\Delta_v \hat{t}_{nv}$ and t_{nn} ,

$$\begin{aligned}
 \sum_{n=v+1}^{m+1} \left(\frac{P_n}{P_n} \right)^{k-1} |\Delta_v \hat{t}_{nv}| |t_{nn}|^{k-1} &= \sum_{n=v+1}^{m+1} \left(\frac{P_n}{P_n} \right)^{k-1} \frac{p_n p_v}{P_n P_{n-1}} \left(\frac{p_n}{P_n} \right)^{k-1} \\
 &= p_v \sum_{n=v+1}^{m+1} \frac{p_n}{P_n P_{n-1}} = \frac{p_v}{P_v} \\
 &= \left(\frac{P_v}{P_v} \right)^{k-1} |t_{vv}|^k \quad \text{as } m \rightarrow \infty,
 \end{aligned} \tag{3.4}$$

condition (2.3) is satisfied.

Since

$$\begin{aligned}
 \hat{t}_{nv} &= \bar{t}_{nv} - \bar{t}_{n-1,v} = \sum_{i=v}^n t_{ni} - \sum_{i=v}^{n-1} t_{n-1,i} \\
 &= \frac{1}{P_n} \sum_{i=v}^n p_i - \frac{1}{P_{n-1}} \sum_{i=v}^{n-1} p_i \\
 &= P_{v-1} \left(-\frac{1}{P_n} + \frac{1}{P_{n-1}} \right) = P_{v-1} \frac{p_n}{P_n P_{n-1}},
 \end{aligned}$$

$$\begin{aligned}
\frac{1}{|t_{nn}|} \sum_{v=0}^{n-1} |\Delta\lambda_v| |\hat{t}_{n,v+1}| &= \frac{P_n}{p_n} \sum_{v=0}^{n-1} |\Delta\lambda_v| P_v \frac{p_n}{P_n P_{n-1}} \\
&= \frac{1}{P_{n-1}} \sum_{v=0}^{n-1} |\Delta\lambda_v| P_v = O(1) \sum_{v=0}^{n-1} |\Delta\lambda_v| = O(1) \quad \text{as } n \rightarrow \infty,
\end{aligned}
\tag{3.5}$$

and condition (2.4) is satisfied.

Finally,

$$\begin{aligned}
\sum_{n=v+1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} |\hat{t}_{n,v+1}| |t_{nn}|^{k-1} &= \sum_{n=v+1}^{m+1} \left(\frac{P_n}{p_n} \right)^{k-1} \frac{P_v p_n}{P_n P_{n-1}} \left(\frac{p_n}{P_n} \right)^{k-1} \\
&= P_v \sum_{n=v+1}^{m+1} \frac{p_n}{P_n P_{n-1}} = O(1) \quad \text{as } m \rightarrow \infty,
\end{aligned}
\tag{3.6}$$

so condition (2.5) is satisfied.

This completes the proof of the corollary. $\square$

References

- [1] H. Bor, *On two summability methods*, Math. Proc. Cambridge Philos. Soc. **97** (1985), no. 1, 147–149.
- [2] ———, *A note on $|\overline{N}, p_n|_k$ summability factors*, Rend. Mat. Appl. (7) **12** (1992), no. 4, 937–942 (1993).
- [3] T. M. Flett, *On an extension of absolute summability and some theorems of Littlewood and Paley*, Proc. London Math. Soc. (3) **7** (1957), 113–141.
- [4] G. H. Hardy, *Divergent Series*, Clarendon Press, Oxford University Press, New York, 1949.
- [5] W. T. Sulaiman, *Inclusion theorems for absolute matrix summability methods of an infinite series. IV*, Indian J. Pure Appl. Math. **34** (2003), no. 11, 1547–1557.

H. S. Özarıslan: Department of Mathematics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey

E-mail address: seyhan@erciyes.edu.tr

H. N. Öğdük: Department of Mathematics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey

E-mail address: nogduk@erciyes.edu.tr