APPLICATION OF AN INTEGRAL FORMULA TO CR-SUBMANIFOLDS OF COMPLEX HYPERBOLIC SPACE

JIN SUK PAK AND HYANG SOOK KIM

Received 24 May 2004 and in revised form 2 March 2005

The purpose of this paper is to study n -dimensional compact CR-submanifolds of complex hyperbolic space $\text{CH}^{(n+p)/2}$, and especially to characterize geodesic hypersphere in $\text{CH}^{(n+1)/2}$ by an integral formula.

1. Introduction

Let $\bar{M}$ be a complex space form of constant holomorphic sectional curvature c and let M be an n -dimensional CR-submanifold of $(n-1)$ CR-dimension in $\bar{M}$. Then M has an almost contact metric structure (F, U, u, g) (see Section 2) induced from the canonical complex structure of $\bar{M}$. Hence on an n -dimensional CR-submanifold of $(n-1)$ CR-dimension, we can consider two structures, namely, almost contact structure F and a submanifold structure represented by second fundamental form A . In this point of view, many differential geometers have classified M under the conditions concerning those structures (cf. [3, 5, 8, 9, 10, 11, 12, 14, 15, 16]). In particular, Montiel and Romero [12] have classified real hypersurfaces M of complex hyperbolic space $\text{CH}^{(n+1)/2}$ which satisfy the commutativity condition

(C)

$$FA = AF \quad (1.1)$$

by using the S^1 -fibration $\pi : H_1^{n+2} \rightarrow \text{CH}^{(n+1)/2}$ of the anti-de Sitter space H_1^{n+2} over $\text{CH}^{(n+1)/2}$, and obtained Theorem 4.1 stated in Section 2. We notice that among the model spaces in Theorem 4.1, the geodesic hypersphere is only compact.

In this paper, we will investigate n -dimensional compact CR-submanifold of $(n-1)$ CR-dimension in complex hyperbolic space and provide a characterization of the geodesic hypersphere, which is equivalent to condition (C), by using the following integral formula established by Yano [17, 18]:

$$\int_M \text{div} \{ (\nabla_X X) - (\text{div} X)X \} * 1 = \int_M \left\{ \text{Ric}(X, X) + \frac{1}{2} \| \mathcal{L}_X g \|^2 - \| \nabla X \|^2 - (\text{div} X)^2 \right\} * 1 = 0, \quad (1.2)$$

where X is an arbitrary vector field tangent to M . Our results of the paper are complex hyperbolic versions of those in [6, 15].

2. Preliminaries

Let M be an n -dimensional CR-submanifold of $(n - 1)$ CR-dimension isometrically immersed in a complex space form $\bar{M}^{(n+p)/2}(c)$. Denoting by $(J, \bar{g})$ the Kähler structure of $\bar{M}^{(n+p)/2}(c)$, it follows by definition (cf. [5, 6, 8, 9, 13, 16]) that the maximal J -invariant subspace

$$\mathcal{D}_x := T_x M \cap J T_x M \quad (2.1)$$

of the tangent space $T_x M$ of M at each point x in M has constant dimension $(n - 1)$. So there exists a unit vector field U_1 tangent to M such that

$$\mathcal{D}_x^\perp = \text{Span} \{U_1\}, \quad \forall x \in M, \quad (2.2)$$

where $\mathcal{D}_x^\perp$ denotes the subspace of $T_x M$ complementary orthogonal to $\mathcal{D}_x$. Moreover, the vector field ξ_1 defined by

$$\xi_1 := J U_1 \quad (2.3)$$

is normal to M and satisfies

$$J T M \subset T M \oplus \text{Span} \{\xi_1\}. \quad (2.4)$$

Hence we have, for any tangent vector field X and for a local orthonormal basis $\{\xi_1, \xi_\alpha\}_{\alpha=2, \dots, p}$ of normal vectors to M , the following decomposition in tangential and normal components:

$$JX = FX + u^1(X)\xi_1, \quad (2.5)$$

$$J\xi_\alpha = -U_\alpha + P\xi_\alpha, \quad \alpha = 1, \dots, p. \quad (2.6)$$

Since the structure $(J, \bar{g})$ is Hermitian and $J^2 = -I$, we can easily see from (2.5) and (2.6) that F and P are skew-symmetric linear endomorphisms acting on $T_x M$ and $T_x M^\perp$, respectively, and that

$$g(FU_\alpha, X) = -u^1(X)\bar{g}(\xi_1, P\xi_\alpha), \quad (2.7)$$

$$g(U_\alpha, U_\beta) = \delta_{\alpha\beta} - \bar{g}(P\xi_\alpha, P\xi_\beta), \quad (2.8)$$

where $T_x M^\perp$ denotes the normal space of M at x and g the metric on M induced from $\bar{g}$. Furthermore, we also have

$$g(U_\alpha, X) = u^1(X)\delta_{1\alpha}, \quad (2.9)$$

and consequently,

$$g(U_1, X) = u^1(X), \quad U_\alpha = 0, \quad \alpha = 2, \dots, p. \quad (2.10)$$

Next, applying J to (2.5) and using (2.6) and (2.10), we have

$$F^2X = -X + u^1(X)U_1, \quad u^1(X)P\xi_1 = -u^1(FX)\xi_1, \quad (2.11)$$

from which, taking account of the skew-symmetry of P and (2.7),

$$u^1(FX) = 0, \quad FU_1 = 0, \quad P\xi_1 = 0. \quad (2.12)$$

Thus (2.6) may be written in the form

$$J\xi_1 = -U_1, \quad J\xi_\alpha = P\xi_\alpha, \quad \alpha = 2, \dots, p. \quad (2.13)$$

These equations tell us that (F, g, U_1, u^1) defines an almost contact metric structure on M (cf. [5, 6, 8, 9, 16]), and consequently, $n = 2m + 1$ for some integer m .

We denote by $\bar{\nabla}$ and ∇ the Levi-Civita connection on $\tilde{M}^{(n+p)/2}(c)$ and M , respectively. Then the Gauss and Weingarten formulas are given by

$$\bar{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad (2.14)$$

$$\bar{\nabla}_X \xi_\alpha = -A_\alpha X + \nabla_X^\perp \xi_\alpha, \quad \alpha = 1, \dots, p, \quad (2.15)$$

for any vector fields X, Y tangent to M . Here $\nabla^\perp$ denotes the normal connection induced from $\bar{\nabla}$ in the normal bundle $TM^\perp$ of M , and h and A_α the second fundamental form and the shape operator corresponding to ξ_α , respectively. It is clear that h and A_α are related by

$$h(X, Y) = \sum_{\alpha=1}^p g(A_\alpha X, Y) \xi_\alpha. \quad (2.16)$$

We put

$$\nabla_X^\perp \xi_\alpha = \sum_{\beta=1}^p s_{\alpha\beta}(X) \xi_\beta. \quad (2.17)$$

Then $(s_{\alpha\beta})$ is the skew-symmetric matrix of connection forms of $\nabla^\perp$.

Now, using (2.14), (2.15), and (2.17), and taking account of the Kähler condition $\bar{\nabla}J = 0$, we differentiate (2.5) and (2.6) covariantly and compare the tangential and normal parts. Then we can easily find that

$$(\nabla_X F)Y = u^1(Y)A_1X - g(A_1Y, X)U_1, \quad (2.18)$$

$$(\nabla_X u^1)(Y) = g(FA_1X, Y), \quad (2.19)$$

$$\nabla_X U_1 = FA_1X, \quad (2.20)$$

$$g(A_\alpha U_1, X) = - \sum_{\beta=2}^p s_{1\beta}(X) \bar{g}(P\xi_\beta, \xi_\alpha), \quad \alpha = 2, \dots, p, \quad (2.21)$$

for any X, Y tangent to M .

In the rest of this paper, we suppose that the distinguished normal vector field ξ_1 is parallel with respect to the normal connection $\nabla^\perp$. Hence (2.17) gives

$$s_{1\alpha} = 0, \quad \alpha = 2, \dots, p, \quad (2.22)$$

which, together with (2.21), yields

$$A_\alpha U_1 = 0, \quad \alpha = 2, \dots, p. \quad (2.23)$$

On the other hand, the ambient manifold $\bar{M}^{(n+p)/2}(c)$ is of constant holomorphic sectional curvature c and consequently, its Riemannian curvature tensor $\bar{R}$ satisfies

$$\bar{R}_{\bar{X}\bar{Y}}\bar{Z} = \frac{c}{4} \{ \bar{g}(\bar{Y}, \bar{Z})\bar{X} - \bar{g}(\bar{X}, \bar{Z})\bar{Y} + \bar{g}(J\bar{Y}, \bar{Z})J\bar{X} - \bar{g}(J\bar{X}, \bar{Z})J\bar{Y} - 2\bar{g}(J\bar{X}, \bar{Y})J\bar{Z} \} \quad (2.24)$$

for any $\bar{X}, \bar{Y}, \bar{Z}$ tangent to $\bar{M}^{(n+p)/2}(c)$ (cf. [1, 2, 4, 19]). So, the equations of Gauss and Codazzi imply that

$$\begin{aligned} R_{XY}Z &= \frac{c}{4} \{ g(Y, Z)X - g(X, Z)Y + g(FY, Z)FX - g(FX, Z)FY - 2g(FX, Y)FZ \} \\ &\quad + \sum_{\alpha} \{ g(A_\alpha Y, Z)A_\alpha X - g(A_\alpha X, Z)A_\alpha Y \}, \end{aligned} \quad (2.25)$$

$$(\nabla_X A_1)Y - (\nabla_Y A_1)X = \frac{c}{4} \{ g(X, U_1)FY - g(Y, U_1)FX - 2g(FX, Y)U_1 \}, \quad (2.26)$$

for any X, Y, Z tangent to M with the aid of (2.22), where R denotes the Riemannian curvature tensor of M . Moreover, (2.11) and (2.25) yield

$$\text{Ric}(X, Y) = \frac{c}{4} \{ (n+2)g(X, Y) - 3u^1(X)u^1(Y) \} + \sum_{\alpha} \{ (\text{tr} A_\alpha)g(A_\alpha X, Y) - g(A_\alpha^2 X, Y) \}, \quad (2.27)$$

$$\rho = \frac{c}{4} (n+3)(n-1) + n^2 \|\mu\|^2 - \sum_{\alpha} \text{tr} A_\alpha^2, \quad (2.28)$$

where Ric and ρ denote the Ricci tensor and the scalar curvature, respectively, and

$$\mu = \frac{1}{n} \sum_{\alpha} (\text{tr} A_\alpha) \xi_\alpha \quad (2.29)$$

is the mean curvature vector (cf. [1, 2, 4, 19]).

3. Codimension reduction of CR-submanifolds of $\text{CH}^{(n+p)/2}$

Let M be an n -dimensional CR-submanifold of $(n-1)$ CR-dimension in a complex hyperbolic space $\text{CH}^{(n+p)/2}$ with constant holomorphic sectional curvature $c = -4$.

Applying the integral formula (1.2) to the vector field U_1 , we have

$$\int_M \left\{ \text{Ric}(U_1, U_1) + \frac{1}{2} \|\mathcal{L}_{U_1} g\|^2 - \|\nabla U_1\|^2 - (\text{div} U_1)^2 \right\} * 1 = 0. \quad (3.1)$$

Now we take an orthonormal basis $\{U_1, e_a, e_{a^*}\}_{a=1, \dots, (n-1)/2}$ of tangent vectors to M such that

$$e_{a^*} := Fe_a, \quad a = 1, \dots, \frac{n-1}{2}. \quad (3.2)$$

Then it follows from (2.11) and (2.20) that

$$\operatorname{div} U_1 = \operatorname{tr}(FA_1) = \sum_{a=1}^{(n-1)/2} \{g(FA_1 e_a, e_a) + g(FA_1 e_{a^*}, e_{a^*})\} = 0. \quad (3.3)$$

Also, using (2.20), we have

$$\|\nabla U_1\|^2 = g(FA_1 U_1, FA_1 U_1) + \sum_{a=1}^{(n-1)/2} \{g(FA_1 e_a, FA_1 e_a) + g(FA_1 e_{a^*}, FA_1 e_{a^*})\}, \quad (3.4)$$

from which, together with (2.11) and (2.12), we can easily obtain

$$\|\nabla U_1\|^2 = \operatorname{tr} A_1^2 - \|A_1 U_1\|^2. \quad (3.5)$$

Furthermore, (2.20) yields

$$(\mathcal{L}_{U_1} g)(X, Y) = g(\nabla_X U_1, Y) + g(\nabla_Y U_1, X) = g((FA_1 - A_1 F)X, Y), \quad (3.6)$$

and consequently,

$$\|\mathcal{L}_{U_1} g\|^2 = \|FA_1 - A_1 F\|^2. \quad (3.7)$$

On the other hand, (2.27) and (2.28) with $c = -4$ yield

$$\operatorname{Ric}(U_1, U_1) = -(n-1) + u^1(A_1 U_1)(\operatorname{tr} A_1) - \|A_1 U_1\|^2, \quad (3.8)$$

$$\operatorname{tr}(A_1^2) = -\rho - (n+3)(n-1) + n^2 \|\mu\|^2 - \sum_{\alpha=2}^p \operatorname{tr} A_\alpha^2. \quad (3.9)$$

Substituting (3.3), (3.5), (3.7), (3.8), and (3.9) into (3.1), we have

$$\begin{aligned} \int_M \left\{ \frac{1}{2} \|FA_1 - A_1 F\|^2 + \operatorname{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 \right. \\ \left. + \|A_1 U_1\|^2 + (n+3)(n-1) + \sum_{\alpha=2}^p \operatorname{tr} A_\alpha^2 \right\} * 1 = 0, \end{aligned} \quad (3.10)$$

or equivalently,

$$\int_M \left\{ \frac{1}{2} \|FA_1 - A_1 F\|^2 + u^1(A_1 U_1)(\operatorname{tr} A_1) - \operatorname{tr} A_1^2 - (n-1) \right\} * 1 = 0. \quad (3.11)$$

Thus we have the following lemma.

LEMMA 3.1. *Let M be an n -dimensional compact orientable CR-submanifold of $(n-1)$ CR-dimension in a complex hyperbolic space $\mathbb{CH}^{(n+p)/2}$. If the distinguished normal vector field ξ_1 is parallel with respect to the normal connection and if the inequality*

$$\text{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 + \|A_1 U_1\|^2 + (n+3)(n-1) \geq 0 \quad (3.12)$$

holds on M , then

$$A_1 F = F A_1 \quad (3.13)$$

and $A_\alpha = 0$ for $\alpha = 2, \dots, p$.

COROLLARY 3.2. *Let M be a compact orientable real hypersurface of $\mathbb{CH}^{(n+1)/2}$ over which the inequality*

$$\text{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 + \|A_1 U_1\|^2 + (n+3)(n-1) \geq 0 \quad (3.14)$$

holds. Then M satisfies the commutativity condition (C).

Combining Lemma 3.1 and the codimension reduction theorem proved in [7, Theorem 3.2, page 126], we have the following theorem.

THEOREM 3.3. *Let M be an n -dimensional compact orientable CR-submanifold of $(n-1)$ CR-dimension in a complex hyperbolic space $\mathbb{CH}^{(n+p)/2}$. If the distinguished normal vector field ξ_1 is parallel with respect to the normal connection and if the inequality*

$$\text{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 + \|A_1 U_1\|^2 + (n+3)(n-1) \geq 0 \quad (3.15)$$

holds on M , then there exists a totally geodesic complex hyperbolic space $\mathbb{CH}^{(n+1)/2}$ immersed in $\mathbb{CH}^{(n+p)/2}$ such that $M \subset \mathbb{CH}^{(n+1)/2}$. Moreover M satisfies the commutativity condition (C) as a real hypersurface of $\mathbb{CH}^{(n+1)/2}$.

Proof. Let

$$N_0(x) := \{\eta \in T_x M^\perp \mid A_\eta = 0\} \quad (3.16)$$

and let $H_0(x)$ be the maximal holomorphic subspace of $N_0(x)$, that is,

$$H_0(x) = N_0(x) \cap JN_0(x). \quad (3.17)$$

Then, by means of Lemma 3.1,

$$H_0(x) = N_0(x) = \text{Span}\{\xi_2, \dots, \xi_p\}. \quad (3.18)$$

Hence, the orthogonal complement $H_1(x)$ of $H_0(x)$ in $TM^\perp$ is $\text{Span}\{\xi_1\}$ and so, $H_1(x)$ is invariant under the parallel translation with respect to the normal connection and $\dim H_1(x) = 1$ at any point $x \in M$. Thus, applying the codimension reduction theorem in [4] proved by Kawamoto, we verify that there exists a totally geodesic complex hyperbolic space $\mathbb{CH}^{(n+1)/2}$ immersed in $\mathbb{CH}^{(n+p)/2}$ such that $M \subset \mathbb{CH}^{(n+1)/2}$. Therefore, M can

be regarded as a real hypersurface of $\text{CH}^{(n+1)/2}$ which is totally geodesic in $\text{CH}^{(n+p)/2}$. Tentatively, we denote $\text{CH}^{(n+1)/2}$ by M' , and by i_1 we denote the immersion of M into M' , and by i_2 the totally geodesic immersion of M' into $\text{CH}^{(n+p)/2}$. Then it is clear from (2.14) that

$$\nabla'_{i_1 X} i_1 Y = i_1 \nabla_X Y + h'(X, Y) = i_1 \nabla_X Y + g(A'X, Y)\xi', \quad (3.19)$$

where ∇' is the induced connection on M' from that of $\text{CH}^{(n+p)/2}$, h' the second fundamental form of M in M' , and A' the corresponding shape operator to a unit normal vector field ξ' to M in M' . Since $i = i_2 \circ i_1$ and M' is totally geodesic in $\text{CH}^{(n+p)/2}$, we can easily see that (2.15) and (3.19) imply that

$$\xi_1 = i_2 \xi', \quad A_1 = A'. \quad (3.20)$$

Since M' is a holomorphic submanifold of $\text{CH}^{(n+p)/2}$, for any X in TM ,

$$J i_2 X = i_2 J' X \quad (3.21)$$

is valid, where J' is the induced Kähler structure on M' . Thus it follows from (2.5) that

$$\begin{aligned} J i X &= J i_2 \circ i_1 X = i_2 J' i_1 X = i_2 (i_1 F' X + u'(X)\xi') \\ &= i F' X + u'(X) i_2 \xi' = i F' X + u'(X)\xi_1 \end{aligned} \quad (3.22)$$

for any vector field X tangent to M . Comparing this equation with (2.5), we have $F = F'$ and $u^1 = u'$, which, together with Lemma 3.1, implies that

$$A' F' = F' A'. \quad (3.23)$$

□

4. An integral formula on the model space $M_{2p+1, 2q+1}^h(r)$

We first explain the model hypersurfaces of complex hyperbolic space due to Montiel and Romero for later use (for the details, see [12]).

Consider the complex $(n+3)/2$ -space $C_1^{(n+3)/2}$ endowed with the pseudo-Euclidean metric g_0 given by

$$g_0 = -dz_0 d\bar{z}_0 + \sum_{j=1}^m dz_j d\bar{z}_j, \quad \left(m+1 := \frac{n+3}{2}\right), \quad (4.1)$$

where $\bar{z}_k$ denotes the complex conjugate of z_k .

On $C_1^{(n+3)/2}$, we define

$$F(z, w) = -z_0 \bar{w}_0 + \sum_{k=1}^m z_k \bar{w}_k. \quad (4.2)$$

Put

$$H_1^{n+2} = \{z = (z_0, z_1, \dots, z_m) \in C_1^{(n+3)/2} : \langle z, z \rangle = -1\}, \quad (4.3)$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product on $C_1^{(n+3)/2}$ induced from g_0 . Then it is known that H_1^{n+2} , together with the induced metric, is a pseudo-Riemannian manifold of constant sectional curvature -1 , which is known as an anti-de Sitter space. Moreover, H_1^{n+2} is a principal S^1 -bundle over $\text{CH}^{(n+1)/2}$ with projection $\pi: H_1^{n+2} \rightarrow \text{CH}^{(n+1)/2}$ which is a Riemannian submersion with fundamental tensor J and time-like totally geodesic fibers.

Given p, q integers with $2p + 2q = n - 1$ and $r \in R$ with $0 < r < 1$, we denote by $M_{2p+1, 2q+1}(r)$ the Lorentz hypersurface of H_1^{n+2} defined by the equations

$$-|z_0|^2 + \sum_{k=1}^m |z_k|^2 = -1, \quad r \left(-|z_0|^2 + \sum_{k=1}^p |z_k|^2 \right) = - \sum_{k=p+1}^m |z_k|^2, \quad (4.4)$$

where $z = (z_0, z_1, \dots, z_m) \in C_1^{(n+3)/2}$. In fact, $M_{2p+1, 2q+1}(r)$ is isometric to the product

$$H_1^{2p+1} \left(\frac{1}{r-1} \right) \times S^{2q+1} \left(\frac{r}{1-r} \right), \quad (4.5)$$

where $1/(r-1)$ and $r/(1-r)$ denote the squares of the radii and each factor is embedded in H_1^{n+2} in a totally umbilical way. Since $M_{2p+1, 2q+1}(r)$ is S^1 -invariant, $M_{2p+1, 2q+1}^h(r) := \pi(M_{2p+1, 2q+1}(r))$ is a real hypersurface of $\text{CH}^{(n+1)/2}$ which is complete and satisfies the condition (C).

As already mentioned in Section 1, Montiel and Romero [12] have classified real hypersurfaces M of $\text{CH}^{(n+1)/2}$ which satisfy the condition (C) and obtained the following classification theorem.

THEOREM 4.1. *Let M be a complete real hypersurface of $\text{CH}^{(n+1)/2}$ which satisfies the condition (C). Then there exist the following possibilities.*

- (1) M has three constant principal curvatures $\tanh \theta$, $\coth \theta$, $2 \coth 2\theta$ with multiplicities $2p$, $2q$, 1 , respectively, $2p + 2q = n - 1$. Moreover, M is congruent to $M_{2p+1, 2q+1}^h(\tanh^2 \theta)$.
- (2) M has two constant principal curvatures λ_1, λ_2 with multiplicities $n - 1$ and 1 , respectively. (i) If $\lambda_1 > 1$, then $\lambda_1 = \coth \theta$, $\lambda_2 = 2 \coth 2\theta$ with $\theta > 0$, and M is congruent to a geodesic hypersphere $M_{1, n}^h(\tanh^2 \theta)$. (ii) If $\lambda_1 < 1$, then $\lambda_1 = \tanh \theta$, $\lambda_2 = 2 \coth 2\theta$ with $\theta > 0$, and M is congruent to $M_{n, 1}^h(\tanh^2 \theta)$. (iii) If $\lambda_1 = 1$, then $\lambda_2 = 2$ and M is congruent to a horosphere.

Combining Corollary 3.2 and Theorem 4.1, we have the following theorem.

THEOREM 4.2. *Let M be a compact orientable real hypersurface of $\text{CH}^{(n+1)/2}$ over which the inequality*

$$\text{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 + \|A_1 U_1\|^2 + (n+3)(n-1) \geq 0 \quad (4.6)$$

holds. Then M is congruent to a geodesic hypersphere $M_{1, n}^h(r)$ in $\text{CH}^{(n+1)/2}$.

Combining Theorems 3.3 and 4.2, we have the following theorem.

THEOREM 4.3. *Let M be an n -dimensional compact orientable CR-submanifold of $(n-1)$ CR-dimension in a complex hyperbolic space $\text{CH}^{(n+p)/2}$. If the distinguished normal vector field ξ_1 is parallel with respect to the normal connection and if the inequality*

$$\text{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 + \|A_1 U_1\|^2 + (n+3)(n-1) \geq 0 \quad (4.7)$$

holds on M , then M is congruent to a geodesic hypersphere $M_{1,n}^h(\tanh^2 \theta)$ in $\text{CH}^{(n+1)/2}$.

Remark 4.4. As already shown in (3.10) and (3.11), the equality

$$\begin{aligned} \text{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 + \|A_1 U_1\|^2 + (n+3)(n-1) \\ = u^1(A_1 U_1)(\text{tr} A_1) - \text{tr} A_1^2 - (n-1) \end{aligned} \quad (4.8)$$

holds on M . On the other hand, the geodesic hypersphere $M_{1,n}^h(\tanh^2 \theta)$ in Theorem 4.1 has constant principal curvatures $\coth \theta$ and $2\coth 2\theta$ with multiplicities $n-1$ and 1 , respectively. Hence we can easily verify the equality

$$u^1(A_1 U_1)(\text{tr} A_1) - \text{tr} A_1^2 - (n-1) = 0, \quad (4.9)$$

and consequently,

$$\text{Ric}(U_1, U_1) + \rho - n^2 \|\mu\|^2 + \|A_1 U_1\|^2 + (n+3)(n-1) = 0 \quad (4.10)$$

on $M_{1,n}^h(\tanh^2 \theta)$.

Remark 4.5. If we put $V := \nabla_{U_1} U_1 - (\text{div} U_1)U_1$, then it easily follows from (2.11) that $V = FA_1 U_1$. Taking account of (3.3), (3.5), (3.7), and (3.8), we obtain

$$\text{div} V = \frac{1}{2} \|FA_1 - A_1 F\|^2 + u^1(A_1 U_1)(\text{tr} A_1) - \text{tr} A_1^2 - (n-1). \quad (4.11)$$

Hence if the commutativity condition (C) holds on M , then the vector field V is zero since U_1 is a principal vector of A_1 , and consequently,

$$u^1(A_1 U_1)(\text{tr} A_1) - \text{tr} A_1^2 - (n-1) = 0. \quad (4.12)$$

Thus, on n -dimensional CR-submanifold M of $(n-1)$ CR-dimension in a complex hyperbolic space $\text{CH}^{(n+p)/2}$ over which the commutativity condition C holds, the function $u^1(A_1 U_1)$ cannot be zero at any point of M . A real hypersurface of a complex hyperbolic space $\text{CH}^{(n+p)/2}$ satisfying the commutativity condition (C) cannot be minimal.

Acknowledgment

This work was supported by ABRL Grant Project no. R14-2002-003-01000-0 from Korea Science and Engineering Foundation (KOSEF).

References

- [1] A. Bejancu, *CR submanifolds of a Kaehler manifold. I*, Proc. Amer. Math. Soc. **69** (1978), no. 1, 135–142.
- [2] ———, *Geometry of CR-Submanifolds*, Mathematics and its Applications (East European Series), vol. 23, D. Reidel Publishing, Dordrecht, 1986.
- [3] A. A. Borisenko, *On the global structure of Hopf hypersurfaces in a complex space form*, Illinois J. Math. **45** (2001), no. 1, 265–277.
- [4] B. Y. Chen, *Geometry of Submanifolds*, Marcel Dekker, New York, 1973.
- [5] Y.-W. Choe and M. Okumura, *Scalar curvature of a certain CR-submanifold of complex projective space*, Arch. Math. (Basel) **68** (1997), no. 4, 340–346.
- [6] M. Djorić and M. Okumura, *Certain application of an integral formula to CR-submanifold of complex projective space*, Publ. Math. Debrecen **62** (2003), no. 1-2, 213–225.
- [7] S. Kawamoto, *Codimension reduction for real submanifolds of a complex hyperbolic space*, Rev. Mat. Univ. Complut. Madrid **7** (1994), no. 1, 119–128.
- [8] J.-H. Kwon and J. S. Pak, *CR-submanifolds of $(n-1)$ CR-dimension in a complex projective space*, Saitama Math. J. **15** (1997), 55–65.
- [9] ———, *n -dimensional CR-submanifolds of $(n-1)$ CR-dimension immersed in a complex space form*, Far East J. Math. Sci., Special volume (1999), Part III, 347–360.
- [10] H. B. Lawson Jr., *Rigidity theorems in rank-1 symmetric spaces*, J. Differential Geom. **4** (1970), 349–357.
- [11] S. Montiel, *Real hypersurfaces of a complex hyperbolic space*, J. Math. Soc. Japan **37** (1985), no. 3, 515–535.
- [12] S. Montiel and A. Romero, *On some real hypersurfaces of a complex hyperbolic space*, Geom. Dedicata **20** (1986), no. 2, 245–261.
- [13] R. Nirenberg and R. O. Wells Jr., *Approximation theorems on differentiable submanifolds of a complex manifold*, Trans. Amer. Math. Soc. **142** (1969), 15–35.
- [14] M. Okumura, *On some real hypersurfaces of a complex projective space*, Trans. Amer. Math. Soc. **212** (1975), 355–364.
- [15] ———, *Compact real hypersurfaces of a complex projective space*, J. Differential Geom. **12** (1977), no. 4, 595–598 (1978).
- [16] M. Okumura and L. Vanhecke, *n -dimensional real submanifolds with $(n-1)$ -dimensional maximal holomorphic tangent subspace in complex projective spaces*, Rend. Circ. Mat. Palermo (2) **43** (1994), no. 2, 233–249.
- [17] K. Yano, *On harmonic and Killing vector fields*, Ann. of Math. (2) **55** (1952), 38–45.
- [18] ———, *Integral Formulas in Riemannian Geometry*, Pure and Applied Mathematics, vol. 1, Marcel Dekker, New York, 1970.
- [19] K. Yano and M. Kon, *CR Submanifolds of Kaehlerian and Sasakian Manifolds*, Progress in Mathematics, vol. 30, Birkhäuser Boston, Massachusetts, 1983.

Jin Suk Pak: Department of Mathematics, College of Natural Science, Kyungpook National University, Daegu 702-701, Korea

E-mail address: jspak@knu.ac.kr

Hyang Sook Kim: Department of Computational Mathematics, School of Computer Aided Science, College of Natural Science, Inje University, Kimhae, Kyungnan 621-749, Korea

E-mail address: mathkim@inje.ac.kr